

Title	Pseudo-orbit tracing property and strong transversality of diffeomorphisms on closed manifolds
Author(s)	Sakai, Kazuhiro
Citation	Osaka Journal of Mathematics. 1994, 31(2), p. 373-386
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9551">https://doi.org/10.18910/9551</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## PSEUDO-ORBIT TRACING PROPERTY AND STRONG TRANSVERSALITY OF DIFFEOMORPHISMS ON CLOSED MANIFOLDS

KAZUHIRO SAKAI

(Received February 8, 1993)

### 1. Introduction

We are interested in the dynamical property of a diffeomorphism  $f$  having the pseudo-orbit tracing property of a closed manifold  $M$ . Let  $d$  be a metric for  $M$ . A sequence of points  $\{x_i\}_{i \in \mathbf{Z}}$  of  $M$  is called a  $\delta$ -pseudo-orbit of  $f$  if  $d(f(x_i), x_{i+1}) < \delta$  for  $i \in \mathbf{Z}$ . A sequence  $\{x_i\}_{i \in \mathbf{Z}}$  is said to be  $f$ - $\varepsilon$ -traced by  $y \in M$  if  $d(f^i(y), x_i) < \varepsilon$  for  $i \in \mathbf{Z}$ .

We say that  $f$  has the *pseudo-orbit tracing property* (abbrev. **POTP**) if for every  $\varepsilon > 0$  there is  $\delta > 0$  such that every  $\delta$ -pseudo-orbit of  $f$  can be  $f$ - $\varepsilon$ -traced by some point.

In [5] Robinson proved that every Axiom A diffeomorphism satisfying strong transversality has **POTP**. Thus it will be natural to ask whether **POTP** implies Axiom A and strong transversality. For this problem we have partial results that are answered in [4] for  $\dim M = 2$  and in [7] for  $\dim M = 3$ . However we have no answer for higher dimensions.

Our aim is to prove the following

**Theorem.** *The  $C^1$  interior of all diffeomorphisms having **POTP** of a closed manifold  $M$ ,  $\mathcal{P}(M)$ , coincides with the set of all Axiom A diffeomorphisms satisfying strong transversality.*

We say that  $f$  has the  $C^1$  *uniform pseudo-orbit tracing property* (abbrev.  $C^1$ -**UPOTP**) if there is a  $C^1$  neighborhood  $\mathcal{U}(f)$  of  $f$  with the property that for  $\varepsilon > 0$  there is  $\delta > 0$  such that every  $\delta$ -pseudo-orbit of  $g \in \mathcal{U}(f)$  is  $g$ - $\varepsilon$ -traced by some point. Since every Axiom A diffeomorphism satisfying strong transversality has  $C^1$ -**UPOTP** (see [6, Theorem]), if we establish our theorem, then the following corollary is obtained.

**Corollary.** *The set of all diffeomorphism having  $C^1$ -**UPOTP** is characterized as the set of all Axiom A diffeomorphisms satisfying strong transversality.*

It was proved in [4] that all periodic points of  $f \in \mathcal{P}(M)$  are hyperbolic. From this we can prove that each  $f$  belonging to  $\mathcal{P}(M)$  satisfies Axiom A with no-cycle. Recently it was shown in general by Aoki [1]. Therefore, to conclude our theorem it remains only to prove the following proposition.

**Proposition.** *Every  $f \in \mathcal{P}(M)$  satisfies strong transversality.*

Unfortunately this can not be proved by the techniques mentioned in [4] and [7]. Thus we need a new technique for the proof of the proposition.

## 2. Proof of Proposition

Let  $\text{Diff}(M)$  denote the set of all diffeomorphisms of  $M$  endowed with  $C^1$  topology, and let  $p = f^n(p)$  ( $n > 0$ ) be a hyperbolic periodic point of  $f \in \text{Diff}(M)$ . Even if  $p$  is hyperbolic, when  $\dim M \geq 3$ , it is not easy to construct an  $f^n$ -invariant foliation in a neighborhood of  $p$  that is compatible with the local stable manifold (i.e. the leaf passing through  $p$  is the local stable manifold of  $p$ ). In this paper, by using Franks's lemma we make a new diffeomorphism  $g$  ( $g^n(p) = p$ ), arbitrarily near to  $f$  in  $C^1$  topology, which has a  $g^n$ -invariant compatible foliation in a neighborhood of  $p$  (see lemmas 1 and 2). This foliation will play an essential role in the proof of the proposition.

Let  $f \in \text{Diff}(M)$  satisfy Axiom A with no-cycle. The non-wandering set  $\Omega(f)$  of  $f$  is expressed as a finite disjoint union of basic sets  $\{\Lambda_i(f)\}$ , and for a sufficiently small  $\varepsilon_0 > 0$  and  $x \in \Omega(f)$  there are a local stable manifold  $W_{\varepsilon_0}^s(x, f)$  and a local unstable manifold  $W_{\varepsilon_0}^u(x, f)$ . Let  $\Lambda(f)$  be a basic set of  $f$ . Since  $\dim W_{\varepsilon_0}^s(x, f) = \dim W_{\varepsilon_0}^s(y, f)$  ( $x, y \in \Lambda(f)$ ), we denote by  $\text{Ind } \Lambda(f)$  the dimension of  $W_{\varepsilon_0}^s(x, f)$  for  $x \in \Lambda(f)$ . If  $g \in \text{Diff}(M)$  is  $C^1$  close to  $f$ , then the number of basic sets  $\{\Lambda_i(g)\}$  of  $g$  coincides with that of basic sets  $\{\Lambda_i(f)\}$  since  $f$  is  $\Omega$ -stable.

Put  $B_\varepsilon(x) = \{y \in M \mid d(x, y) \leq \varepsilon\}$  for  $\varepsilon > 0$  and let  $\rho$  be a usual  $C^1$  metric of  $\text{Diff}(M)$ . Then we have the following

**Lemma 1.** *Let  $\varepsilon_0 > 0$  be as above and let  $\Lambda(f)$  be a basic set such that  $1 \leq \text{Ind } \Lambda(f) \leq \dim M - 1$ . Then, for a periodic point  $p \in \Lambda(f)$  ( $f^n(p) = p$ ,  $n > 0$ ), a neighborhood  $\mathcal{U}(f) \subset \text{Diff}(M)$  and a number  $\gamma > 0$  there are  $0 < \varepsilon_1 < \varepsilon_0/2$ ,  $g \in \mathcal{U}(f)$  and a basic set  $\Lambda(g)$  for  $g$  such that*

$$(i) \quad B_{4\varepsilon_1}(f^i(p)) \cap B_{4\varepsilon_1}(f^j(p)) = \phi \text{ for } 0 \leq i \neq j \leq n-1,$$

$$(ii) \quad g(x) = \begin{cases} \exp_{f^{i+1}(p)} \circ D_{f^i(p)} f \circ \exp_{f^i(p)}^{-1}(x) & \text{if } x \in B_{\varepsilon_1}(f^i(p)) \text{ for } 0 \leq i \leq n-1, \\ f(x) & \text{if } x \notin \bigcup_{i=0}^{n-1} B_{4\varepsilon_1}(f^i(p)), \end{cases}$$

(iii)  $g^n(p) = p \in \Lambda(g)$  and  $\rho(W_{\varepsilon_0}^\sigma(p, f), W_{\varepsilon_0}^\sigma(p, g)) < \gamma$  for  $\sigma = s, u$  (i.e. there is a  $C^1$  diffeomorphism  $\xi^\sigma: W_{\varepsilon_0}^\sigma(p, f) \rightarrow W_{\varepsilon_0}^\sigma(p, g)$  such that  $\rho(\xi^\sigma, id) < \gamma$  ( $\sigma = s, u$ )).

Proof. Since  $\Lambda(f)$  is hyperbolic, there is  $e > 0$  such that  $d(f^n(x), f^n(y)) \leq e$  ( $x, y \in \Lambda(f)$  and  $n \in \mathbf{Z}$ ) implies  $x = y$  (see [5]). By  $\Omega$ -stability theorem, there exists a neighborhood  $\mathcal{U}_0(f) \subset \mathcal{U}(f)$  of  $f$  such that for every  $g \in \mathcal{U}_0(f)$  there is a homeomorphism  $h_g$ , which maps  $\Omega(f)$  onto the non-wandering set  $\Omega(g)$  of  $g$ , satisfying

$$\begin{cases} g \circ h_g = h_g \circ f, \\ d(h_g, id|_{\Omega(f)}) < e, \\ \rho(W_{\varepsilon_0}^\sigma(p, f), W_{\varepsilon_0}^\sigma(h_g(p), g)) < \gamma \text{ for } \sigma = s, u. \end{cases}$$

By Franks's lemma [2, lemma 1.1], we can find  $g \in \mathcal{U}_0(f)$  and  $0 < \varepsilon_1 < \varepsilon_0/2$  such that

$$B_{4\varepsilon_1}(f^i(p)) \cap B_{4\varepsilon_1}(f^j(p)) = \emptyset \quad (0 \leq i \neq j \leq n-1) \text{ and}$$

$$g(x) = \begin{cases} \exp_{f^{i+1}(p)} \circ D_{f^i(p)} f \circ \exp_{f^i(p)}^{-1}(x) & \text{if } x \in B_{\varepsilon_1}(f^i(p)) \text{ for } 0 \leq i \leq n-1, \\ f(x) & \text{if } x \notin \bigcup_{i=0}^{n-1} B_{4\varepsilon_1}(f^i(p)), \end{cases}$$

We write  $\Lambda(g) = h_g(\Lambda(f))$  for simplicity. Then  $h_g(p) \in \Lambda(g)$  and  $\text{Ind } \Lambda(f) = \text{Ind } \Lambda(g)$ . Clearly  $g(f^i(p)) = \exp_{f^{i+1}(p)} \circ D_{f^i(p)} f \circ \exp_{f^i(p)}^{-1}(f^i(p)) = f^{i+1}(p)$  for  $0 \leq i \leq n-1$  and so  $g(p) = f(p), g^2(p) = f^2(p), \dots, g^n(p) = f^n(p) = p$ . Since

$$\begin{aligned} d(f^i(h_g^{-1}(p)), f^i(p)) &= d(h_g^{-1}(g^i(p)), f^i(p)) \\ &= d(h_g^{-1}(f^i(p)), f^i(p)) < e \quad (i \in \mathbf{Z}), \end{aligned}$$

we have  $h_g(p) = p$ . Therefore  $\rho(W_{\varepsilon_0}^\sigma(p, f), W_{\varepsilon_0}^\sigma(p, g)) < \gamma$  ( $\sigma = s, u$ ) and  $p \in \Lambda(g)$ .

Since  $f$  satisfies Axiom A, by definition there is a  $Df$ -invariant continuous splitting  $T_{\Omega(f)}M = E^s \oplus E^u$  and a constant  $0 < \lambda < 1$  such that

$\|Df|_{E^s}\| \leq \lambda^m$  and  $\|Df|_{E^u}\| \leq \lambda^m$  for  $m > 0$ . We denote by  $E_x^\sigma$  a fiber of  $E^\sigma$  at  $x \in \Omega(f)$  ( $\sigma = s, u$ ), and put  $E_x^\sigma(\varepsilon) = \{v \in E_x^\sigma \mid \|v\| \leq \varepsilon\}$  for  $\varepsilon > 0$ .

Let  $g \in \text{Diff}(M)$ ,  $p = g^n(p) \in \Lambda(g)$  ( $n > 0$ ) and  $\varepsilon_1 > 0$  be as in lemma 1. Then it is easily checked that for  $0 < \varepsilon \leq \varepsilon_1$ , we have  $\exp_p(E_p^\sigma(\varepsilon)) = W_\varepsilon^\sigma(p, g)$  and  $\dim \exp_p(E_p^\sigma(\varepsilon)) = \dim W_{\varepsilon_0}^\sigma(p, g)$  ( $\sigma = s, u$ ). Fix  $\varepsilon_2$  with  $0 < \varepsilon_2 = \varepsilon_2(g, n) < \varepsilon_1$  such that  $x \in B_{\varepsilon_2}(p)$  implies  $g^i(x) \in B_{\varepsilon_1}(g^i(p))$  for  $0 \leq i \leq n-1$ , and define

$$\tilde{W}_{\varepsilon_2}^s(x, g) = \exp_p \left( E_p^s(\varepsilon_2) + \exp_p^{-1}(x) \right)$$

for  $x \in \exp_p(E_p^u(\varepsilon_2))$ . Then, since  $\bigcup_{v \in E_p^u(\varepsilon_2)} (E_p^s(\varepsilon_2) + v)$  is a foliation defined in a neighborhood of  $O_p \in T_p M$  and since  $\exp_p$  is a local diffeomorphism, we have that  $\{\tilde{W}_{\varepsilon_2}^s(x, g) : x \in \exp_p(E_p^u(\varepsilon_2))\}$  is a foliation defined in a neighborhood of  $p$  in  $M$  such that  $\tilde{W}_{\varepsilon_2}^s(p, g) = W_{\varepsilon_2}^s(p, g)$ .

**Lemma 2.**

- (i)  $\tilde{W}_{\varepsilon_2}^s(x, g)$  is a  $C^1$  manifold and  $\dim \tilde{W}_{\varepsilon_2}^s(x, g) = \dim \tilde{W}_{\varepsilon_2}^s(p, g)$ ,
- (ii)  $g^n(\tilde{W}_{\varepsilon_2}^s(x, g)) \subset \tilde{W}_{\varepsilon_2}^s(g^n(x), g)$  for  $x \in \exp_p(E_p^u(\varepsilon_2)) \cap g^{-n}(\exp_p(E_p^u(\varepsilon_2)))$ ,
- (iii) there exists  $C > 0$  such that if  $\{x, g^n(x), \dots, g^{nk}(x)\} \subset \exp_p(E_p^u(\varepsilon_2))$  for some  $k > 0$ , then  $d(g^{nk}(x), g^{nk}(y)) \leq C\lambda^{nk}d(x, y)$  for  $y \in \tilde{W}_{\varepsilon_2}^s(x, g)$ ,

Proof. Assertion (i) is clear, and (ii) is easily obtained. To show (iii) put  $T_p(\varepsilon_2) = \{v \in T_p M \mid \|v\| \leq \varepsilon_2\}$ . Since  $\exp_p : T_p(\varepsilon_2) \rightarrow M$  and  $\exp_p^{-1} : B_{\varepsilon_2}(p) \rightarrow T_p M$  are into diffeomorphisms there is  $K > 0$  such that

$$d(\exp_p(v), \exp_p(w)) \leq K\|v - w\| \quad (v, w \in T_p(\varepsilon_2)),$$

$$\|\exp_p^{-1}(x) - \exp_p^{-1}(y)\| \leq Kd(x, y) \quad (x, y \in B_{\varepsilon_2}(p)).$$

If  $\{x, g^n(x), \dots, g^{nk}(x)\} \subset \exp_p(E_p^u(\varepsilon_2))$  for some  $k > 0$ , then for  $y \in \tilde{W}_{\varepsilon_2}^s(x, g)$  there is  $v_y \in E_p^s(\varepsilon_2)$  such that  $y = \exp_p(v_y + \exp_p^{-1}(x))$ . Thus we have

$$g^n(y) = \exp_p \left( D_p f^n(v_y) + \exp_p^{-1}(g^n(x)) \right)$$

(since  $D_p f^n(\exp_p^{-1}(x)) = \exp_p^{-1}(g^n(x))$ ), and so

$$\left( D_p f^n \circ \exp_p^{-1} \circ g^n \right) (y) = D_p f^{2n}(v_y) + D_p f^n(\exp_p^{-1}(g^n(x))),$$

from which

$$g^{2n}(y) = \exp(D_p f^{2n}(v_y) + D_p f(\exp_p^{-1}(g^{2n}(x))))$$

Since  $g^n(x) \in B_{\varepsilon_2}(p)$ , we have  $(\exp_p \circ D_p f^n \circ \exp_p^{-1})(g^n(x)) = g^{2n}(x)$ ; i.e.  $D_p f^n(\exp_p^{-1}(g^n(x))) = \exp_p^{-1}(g^{2n}(x))$ . Thus  $g^{2n}(y) = \exp_p(D_p f^{2n}(v_y) + \exp_p^{-1}(g^{2n}(x)))$ . By repetition we have

$$g^{nk}(y) = \exp_p(D_p f^{nk}(v_y) + \exp_p^{-1}(g^{nk}(x)))$$

from which

$$\begin{aligned} d(g^{nk}(x), g^{nk}(y)) &\leq K \|\exp_p^{-1}(g^{nk}(x)) - \exp_p^{-1}(g^{nk}(y))\| \\ &= K \|D_p f^{nk}(v_y)\| \\ &\leq K \lambda^{nk} \|v_y\|. \end{aligned}$$

Clearly,  $\|v_y\| = \|\exp_p^{-1}(x) - \exp_p^{-1}(y)\| \leq K d(x, y)$  since  $\exp_p^{-1}(y) = v_y + \exp_p^{-1}(x)$ . Therefore,  $d(g^{nk}(x), g^{nk}(y)) \leq K^2 \lambda^{nk} d(x, y)$ . Assertion (iii) was proved.

Let  $f$  be as before, and denote by  $W^s(x, f)$  the stable manifold and by  $W^u(x, f)$  the unstable manifold for  $x \in \Omega(f)$  respectively.

**Lemma 3.** *Let  $\Lambda_1(f)$  and  $\Lambda_2(f)$  be two distinct basic sets for  $f$ . Suppose that there are  $p = f^n(p) \in \Lambda_1(f)$  ( $n > 0$ ),  $q \in \Lambda_2(f)$  and  $x \in M \setminus \Omega(f)$  such that  $x \in W^s(p, f) \cap W^u(q, f)$ . Then, for neighborhood  $\mathcal{U}(f) \subset \text{Diff}(M)$  there are  $0 < \varepsilon_1 < \varepsilon_0/2$ ,  $g \in \mathcal{U}(f)$  and two distinct basic sets  $\Lambda_1(g)$  and  $\Lambda_2(g)$  for  $g$  such that*

$$(I) \quad B_{4\varepsilon_2}(f^i(p)) \cap B_{4\varepsilon_2}(f^j(p)) = \emptyset \text{ for } 0 \leq i \neq j \leq n-1,$$

$$(II) \quad g(z) = \begin{cases} \exp_{f^{i+1}(p)} \circ D_{f^i(p)} f \circ \exp_{f^i(p)}^{-1}(z) & \text{if } z \in B_{\varepsilon_1}(f^i(p)) \text{ for } 0 \leq i \leq n-1, \\ f(z) & \text{if } z \notin \bigcup_{i=0}^{n-1} B_{4\varepsilon_1}(f^i(p)), \end{cases}$$

$$(III) \quad \begin{cases} p = g^n(p) \in \Lambda_1(g), \quad q \in \Lambda_2(g), \\ x \in W^s(p, g) \cap W^u(q, g), \\ T_x W^s(p, g) = T_x W^s(p, f) \text{ and } T_x W^u(q, g) = T_x W^u(q, f). \end{cases}$$

Proof. Fix  $\mathcal{U}(f) \subset \text{Diff}(M)$ . By lemma 1, for any  $\gamma > 0$  there are

$0 < \varepsilon_1 < \varepsilon_0/2$ ,  $g \in \mathcal{U}(f)$  and a basic set  $\Lambda_1(g)$  satisfying properties (i), (ii) and (iii) of lemma 1. Put  $\Lambda_2(g) = \Lambda_2(f)$ . Then  $q \in \Lambda_2(g)$ . Since  $\gamma$  is arbitrarily small, by (iii) there are a new diffeomorphism  $\tilde{g} \in \mathcal{U}(f)$  and a small neighborhood  $U(x)$  of  $x$  such that  $\tilde{g}(y) = g(y)$  for all  $y \notin U(x)$  and such that

$$\begin{cases} x \in W^s(p, \tilde{g}) \cap W^u(q, \tilde{g}), \\ T_x W^s(p, \tilde{g}) = T_x W^s(p, f), \\ T_x W^s(q, \tilde{g}) = T_x W^s(q, f), \end{cases}$$

For simplicity we identify  $\tilde{g}$  with  $g$ . Thus (I), (II) and (III) are concluded.


Let  $g \in \mathcal{U}(f)$ ,  $p = g^n(p) \in \Lambda_1(g)$  and  $\varepsilon_1 > 0$  be as in lemma 3 and suppose that  $\dim M - \text{Ind } \Lambda_1(f) \geq 2$ . Take  $0 < \varepsilon_2 \leq \varepsilon_1$  be as in lemma 2, and fix  $\alpha > 0$  such that  $D_p f|_{E_p^u(\alpha)} \subset E_p^u(\varepsilon_2)$ . Put  $D^u(p) = \exp_p(E_p^u(\alpha))$ . Then we have

$$\begin{aligned} d(\tilde{W}_{\varepsilon_2}^s(g^{2n}(F^u(p, g)), g), \tilde{W}_{\varepsilon_2}^s(F^u(p, g), g)) &> 0, \\ d(\tilde{W}_{\varepsilon_2}^s(F^u(p, g), g), \tilde{W}_{\varepsilon_2}^s(g^{-2n}(D^u(p)), g)) &> 0 \end{aligned}$$

where

$$(1) \quad F^u(p, g) = \overline{D^u(p)} \setminus \overline{g^{-n}(D^u(p))}$$

is a fundamental domain of  $W_{\varepsilon_2}^u(p, g)$  (recall that  $\exp_p(E_p^u(\varepsilon)) = W_\varepsilon^u(p, g)$  for  $0 < \varepsilon \leq \varepsilon_2$ ).



Let  $G$  be a linear subspace of  $E_p^u$  such that  $1 \leq \dim G < \dim E_p^u$  and write  $B_r^u(E) = B_r(E) \cap \exp_p(E_p^u(\varepsilon_2))$  for a subset  $E$  of  $M$ . Then we can find  $0 < r_0 < \varepsilon_2$  such that

$$(2) \quad F^u(p, g) \setminus B_{r_0}^u(\exp_p(G \cap E_p^u(\varepsilon_2)) \cap F^u(p, g)) \neq \emptyset$$

for every  $G$ . Since

$$\begin{aligned} r'_0 &= d(\tilde{W}_{\varepsilon_2}^s(g^{2n}(F^u(p, g)), g), \tilde{W}_{\varepsilon_2}^s(F^u(p, g), g)) > 0, \\ r''_0 &= d(\tilde{W}_{\varepsilon_2}^s(F^u(p, g), g), \tilde{W}_{\varepsilon_2}^s(g^{-2n}(D^u(p)), g)) > 0, \end{aligned}$$

we define a positive number  $r_1 = \frac{1}{4} \min\{r_0, r'_0, r''_0\}$ .

Put

$$\Gamma(p) = \bigcup_{y \in \exp_p(E_p^u(\varepsilon_2))} \tilde{W}_{\varepsilon_2}^s(y, g).$$

Then, for any  $z \in \Gamma(p)$ , we can find only one point  $y \in \exp_p(E_p^u(\varepsilon_2))$  such that  $z \in \tilde{W}_{\varepsilon_2}^s(y, g)$ , and so we write

$$(3) \quad \pi(z) = y.$$

Then  $\pi: \Gamma(p) \rightarrow \exp_p(E_p^u(\varepsilon_2))$  is differentiable and which plays an essential role in the proof of the proposition. For  $z \in \Gamma(p) \setminus W_{\varepsilon_2}^s(p, g)$ , there is an integer  $N_z > 0$  such that  $g^{ni}(\pi(z)) \in D^u(p)$  for  $0 \leq i \leq N_z$  (especially  $g^{nN_z}(\pi(z)) \in F^u(p, g)$ ) and  $g^{n(N_z+1)}(\pi(z)) \notin D^u(p)$ .

**Lemma 4.** *Under the above notations, there is  $0 < \varepsilon_3 < r_1$  such that  $\text{diam } \pi(B_{\varepsilon_3}(g^{nN_z}(z))) < r_1$  for every  $z \in \left(\bigcup_{y \in W_{\varepsilon_3}^u(p, g)} \tilde{W}_{\varepsilon_3}^s(y, g)\right) \setminus W_{\varepsilon_3}^s(p, g)$ .*

*Proof.* If this is false, for  $k > 0$  there are

$$z_k \in \left(\bigcup_{y \in W_{\frac{1}{k}}^u(p, g)} \tilde{W}_{\frac{1}{k}}^s(y, g)\right) \setminus W_{\frac{1}{k}}^s(p, g)$$

and  $N_k = N_{z_k} > 0$  such that  $\text{diam } \pi(B_{\frac{1}{k}}(g^{nN_k}(z_k))) \geq r_1$ . Since  $z_k \in \tilde{W}_{\frac{1}{k}}^s(\pi(z_k), g)$ , we have  $N_k \rightarrow \infty$  as  $k \rightarrow \infty$  (because of  $\pi(z_k) \in W_{\frac{1}{k}}^u(p, g)$ ). From  $g^{ni}(\pi(z_k)) \in D^u(p) \subset \exp_p(E_p^u(\varepsilon_2))$  for  $0 \leq i \leq N_k$ , we have

$$d(g^{nN_k}(\pi(z_k)), g^{nN_k}(z_k)) \leq C\lambda^{nN_k}d(\pi(z_k), z_k) \rightarrow 0 \text{ as } k \rightarrow \infty$$

by lemma 2 (iii).

For  $k > 0$  there are  $w_k, w'_k \in \exp_p(E_p^u(\varepsilon_2)), v_k \in \tilde{W}_{\varepsilon_2}^s(w_k, g) \cap B_{\frac{1}{k}}(g^{nN_k}(z_k))$  and  $v'_k \in \tilde{W}_{\varepsilon_2}^s(w'_k, g) \cap B_{\frac{1}{k}}(g^{nN_k}(z_k))$  such that  $d(w_k, w'_k) \geq r_1$ . If  $w_k \rightarrow w$  and  $w'_k \rightarrow w'$  ( $k \rightarrow \infty$ ), then  $w, w' \in \exp_p(E_p^u(\varepsilon_2))$  and  $d(w, w') \geq r_1$ . When  $v_k \rightarrow v$  and  $v'_k \rightarrow v'$  as  $k \rightarrow \infty$ , we have  $v = v' \in \exp_p(E_p^u(\varepsilon_2))$  by the properties

$$\begin{cases} g^{nN_k}(\pi(z_k)) \in \exp_p(E_p^u(\varepsilon_2)), \\ d(g^{nN_k}(\pi(z_k)), g^{nN_k}(z_k)) \rightarrow 0 \text{ as } k \rightarrow \infty, \\ d(v_k, g^{nN_k}(z_k)) < \frac{1}{k} \text{ and } d(v'_k, g^{nN_k}(z_k)) < \frac{1}{k}. \end{cases}$$

Since  $\tilde{W}_{\varepsilon_2}^s(y, g)$  ( $y \in \exp_p(E_p^u(\varepsilon_2))$ ) is continuous with respect to  $y$ , we have  $v \in \tilde{W}_{\varepsilon_2}^s(w, g)$ . Thus  $v = w$  since  $\tilde{W}_{\varepsilon_2}^s(w, g) \cap \exp_p(E_p^u(\varepsilon_2))$  is a single point and  $v, w \in \exp_p(E_p^u(\varepsilon_2))$ . In this way we get  $w = v = v' = w'$ , thus contradicting.

We are in a position to prove the proposition. Hereafter let  $\dim M \geq 4$  and  $f \in \mathcal{P}(M)$ . Notice that  $f$  satisfies Axiom A with no-cycle.

Fix  $x \in M \setminus \Omega(f)$ . Then there are distinct basic sets  $\Lambda_i(f)$  ( $i = 1, 2$ ) such that  $x \in W^s(\Lambda_1(f), f) \cap W^u(\Lambda_2(f), f)$ . If  $\text{Ind } \Lambda_1(f) = \dim M$  or  $\dim M - 1$ , then by the proof of [4, Theorem 2] we have  $T_x M = T_x W^s(x, f) + T_x W^u(x, f)$ . Thus it is enough to prove the above equality for the case when  $1 \leq \text{Ind } \Lambda_1(f) \leq \dim M - 2$ .

Since  $\Omega(f) = \bar{P}(f)$ , there is  $f' \in \mathcal{P}(M)$  arbitrarily near to  $f$  in a  $C^1$  topology satisfying

(a)  $f(y) = f'(y)$  for all  $y$  outside of a small neighborhood of  $x$ ,

(b) there are  $p = f^n(p) \in \Lambda_1(f)$  for some  $n > 0$  and  $q \in \Lambda_2(f)$  such that  $x \in W^s(p, f') \cap W^u(q, f')$ ,  $T_x W^s(p, f') = T_x W^s(x, f)$  and  $T_x W^u(q, f') = T_x W^u(x, f)$ .

By (a) there are basic sets  $\Lambda_i(f')$  ( $i = 1, 2$ ) for  $f'$  such that  $\Lambda_i(f') = \Lambda_i(f)$  ( $i = 1, 2$ ) since  $f$  is  $\Omega$ -stable. We shall prove that  $T_x M = T_x W^s(p, f') + T_x W^u(q, f')$  for the case when  $1 \leq \text{Ind } \Lambda_1(f) \leq \dim M - 2$ . For simplicity we identify  $f'$  with  $f$ .


Let  $\mathcal{U}(f)$  be a small neighborhood of  $f$  such that  $\mathcal{U}(f) \subset \mathcal{P}(M)$ . Then, by lemma 3 there are  $g \in \mathcal{U}(f)$  and basic sets  $\Lambda_i(g)$  ( $i = 1, 2$ ) satisfying lemma 3 (I), (II) and (III). Thus  $T_x W^s(p, g) = T_x W^s(x, f)$  and  $T_x W^u(q, g) = T_x W^u(x, f)$ . Let  $\varepsilon_3 > 0$  be as in lemma 4 and define

$$V_k(p) = \bigcup_{y \in g^{-nk}(F^u(p, g))} \tilde{W}_{\varepsilon_3}^s(y, g) \text{ for } k \geq 0$$

where  $F^u(p, g)$  is the fundamental domain of  $W^u_{\varepsilon_2}(p, g)$  (see (1)). Notice that  $V_k(p) \subset \Gamma(p)$  for  $k \geq 0$  and that  $V_k(p) \rightarrow \tilde{W}^s_{\varepsilon_3}(p, g) = W^s_{\varepsilon_3}(p, g)$  as  $k \rightarrow \infty$  since  $g^{-nk}(F^u(p, g)) \rightarrow \{p\}$  as  $k \rightarrow \infty$ . Thus there is  $k_0 > 0$  such that

$$V_{k_0}(p) \subset \bigcup_{y \in W^u_{\varepsilon_3}(p, g)} \tilde{W}^s_{\varepsilon_3}(y, g).$$

Obviously  $\bigcup_{k \geq k_0} V_k(p)$  is a neighborhood of  $p$  in  $M$ .



Pick  $l > 0$  such that  $g^l(x) \in \text{int} \left( \bigcup_{k \geq k_0} V_k(p) \right)$  and  $g^{-l}(x) \in W^u_{\varepsilon_0/2}(g^{-l}(q), g)$ , and denote by  $C^u(g^l(x))$  the connected component of  $g^l(x)$  in  $W^u(g^l(q), g) \cap \left( \bigcup_{k \geq k_0} V_k(p) \right)$ . Clearly,  $\exp_p^{-1}(C^u(g^l(x))) \subset T_p M$ .

For a linear subspace  $E$  of  $T_p M$  and  $v > 0$  we write

$$E_v(g^l(x)) = \{v + \exp_p^{-1}(g^l(x)) \mid v \in E \text{ with } \|v\| \leq v\}.$$

Then there are a linear subspace  $E'$  of  $T_p M$  and a number  $0 < v_0 \leq \varepsilon_3$  such that

$$(4) \quad T_{g^l(x)} \exp_p(E'_{v_0}(g^l(x))) = T_{g^l(x)} C^u(g^l(x))$$

and  $\exp_p(E'_{v_1}(g^l(x))) \subset \bigcup_{k \geq k_0} V_k(p)$  for  $0 < v \leq v_0$ .

Since  $g^l(x) \notin \Omega(g)$ , there exists  $0 < v_1 \leq v_0$  such that  $B_{v_1}(g^l(x)) \cap g^i(B_{v_1}(g^l(x))) = \emptyset$  for  $i \in \mathbb{Z} \setminus \{0\}$ . Let  $\mathcal{U}(g)$  be a neighborhood of  $g$  such that  $\mathcal{U}(g) \subset \mathcal{U}(f)$ . By (4) there are  $0 < v_2 < v_1$  and  $\varphi \in \text{Diff}(M)$  such that

$$\left\{ \begin{array}{l} \varphi|_{(B_{v_2}(g^l(x)))^c} = \text{id}, \\ \varphi(g^l(x)) = g^l(x), \\ \varphi(\exp_p(E'_{v_2}(g^l(x)))) \subset C^u(g^l(x)), \\ \dim \varphi(\exp_p(E'_{v_2}(g^l(x)))) = \dim C^u(g^l(x)), \\ g' \in \mathcal{U}(g) \text{ where } g' = \varphi^{-1} \circ g. \end{array} \right.$$

We denote  $\exp_p(E'_{v_2}(g^l(x)))$  by  $\exp_p(E'_{v_2}(g'^l(x)))$  because of  $g^l(x) = g'^l(x)$ .

It is clear that there are two distinct basic sets  $\Lambda_i(g')$  ( $i = 1, 2$ ) such that  $\Lambda_i(g') = \Lambda_i(g)$  ( $i = 1, 2$ ) since  $g$  is  $\Omega$ -stable, and that

$$\begin{aligned} W_{\varepsilon_0}^\sigma(p, g') &= W_{\varepsilon_0}^\sigma(p, g), \\ W_{\varepsilon_0}^\sigma(q, g') &= W_{\varepsilon_0}^\sigma(q, g), \\ T_x W^\sigma(x, g') &= T_x W^\sigma(x, g) \quad (\sigma = s, u), \\ \exp_p(E'_{v_2}(g'^l(x))) &\subset W^u(g'^l(q), g') \cap \Gamma(p), \\ \dim \exp_p(E'_{v_2}(g'^l(x))) &= \dim W^u(q, g') = \dim C^u(g'^l(x)). \end{aligned}$$

**Lemma 5.** *Under the above notations,  $\exp_p(E'_{v_2}(g'^l(x)))$  meets transversely  $W_{\varepsilon_3}^s(p, g')$  at  $g'^l(x)$ .*

*Proof.* Let  $\varepsilon_2 > 0$  be as in lemma 2. Since  $W_{\varepsilon_3}^s(p, g') \subset \exp_p(E_p^s(\varepsilon_2))$  and  $W_{\varepsilon_3}^s(p, g') \subset \exp_p(E_p^u(\varepsilon_2))$ , to get the conclusion it is enough to prove

$$\dim \pi(\exp_p(E'_{v_2}(g'^l(x)))) \geq \dim W_{\varepsilon_3}^s(p, g').$$

Here  $\pi: \Gamma(p) \rightarrow \exp_p(E_p^u(\varepsilon_2))$  is the map defined as in (3).

Assume that  $\dim \pi(\exp_p(E'_{v_2}(g'^l(x)))) < \dim W_{\varepsilon_3}^u(p, g')$  and put  $C_\varepsilon^u(g'^l(x)) = B_\varepsilon(g'^l(x)) \cap g'^{2l}(W_{\varepsilon_0}^u(g'^{-l}(q), g'))$  for  $\varepsilon > 0$ . Take  $0 < \varepsilon < v_2$  such that  $C_\varepsilon^u(g'^l(x))$  is the connected component of  $g'^l(x)$  in  $B_{\tilde{\varepsilon}}(g'^l(x)) \cap g'^{2l}(W_{\varepsilon_0}^u(g'^{-l}(q), g'))$  for  $0 < \varepsilon \leq \tilde{\varepsilon}$ , and such that  $B_{\tilde{\varepsilon}}(g'^l(x)) \cap g'^{2l}(W_{\varepsilon_0}^u(g'^{-l}(q), g')) \subset \exp(E'_{v_2}(g'^l(x)))$ .

**Claim 1.** *Let  $0 < \varepsilon \leq \tilde{\varepsilon}$ . If  $d(g'^{-i}(g^l(x)), g'^{-i}(w)) < \varepsilon$  for  $i \geq 0$ , then  $w \in C_\varepsilon^u(g^l(x))$ .*

It is clear that  $d(g'^{-l-i}(x), g'^{-2l-i}(w)) < \varepsilon \leq \varepsilon_0/2$  for all  $i \geq 0$ . On the other hand, since  $d(g'^{-l-i}(x), g'^{-l-i}(q)) < \varepsilon_0/2$  ( $i \geq 0$ ),

$$d(g'^{-2l-i}(w), g'^{-l-i}(q)) \leq d(g'^{-2l-i}(w), g'^{-l-i}(x)) + d(g'^{-l-i}(x), g'^{-l-i}(q)) < \varepsilon_0$$

for all  $i \geq 0$ , and so  $g'^{-2l}(w) \in W_{\varepsilon_0}^u(g'^{-l}(q), g')$ . Thus  $w \in C_\varepsilon^u(g^l(x)) = B_\varepsilon(g^l(x)) \cap g'^{2l}(W_{\varepsilon_0}^u(g'^{-l}(q), g'))$  since  $d(g^l(x), w) < \varepsilon$ .

We divide the proof of this lemma into two cases:

**Case 1.**  $C_\varepsilon^u(g^l(x)) \subset W_{\varepsilon_3}^s(p, g')$ ,

**Case 2.**  $C_\varepsilon^u(g^l(x)) \not\subset W_{\varepsilon_3}^s(p, g')$ ,

For case 1, put  $\varepsilon = \tilde{\varepsilon}/2$  and let  $0 < \delta = \delta(\varepsilon, g') < \varepsilon$  be the number in the definition of **POTP** of  $g'$ . Recall that  $F^u(p, g') = F^u(p, g)$  and fix  $y \in \bigcup_{k \geq k_0} g'^{-nk}(F^u(p, g)) \setminus \{p\}$  such that  $\tilde{W}_{\varepsilon_3}^s(y, g') \cap B_\delta(g^l(x)) \neq \emptyset$ . For  $z \in \tilde{W}_{\varepsilon_3}^s(y, g') \cap B_\delta(g^l(x))$ ,

$$\{\dots, g'^{-1}(x), x, g'(x), \dots, g'^{l-1}(x), z, g'(z), g'^2(z), \dots\}$$

is a  $\delta$ -pseudo-orbit of  $g'$ . Thus there exists  $w \in M$  such that  $d(g'^i(w), g'^i(z)) < \varepsilon$  ( $i \geq 0$ ) and  $d(g'^{-i}(w), g'^{-i}(g^l(x))) < \varepsilon$  ( $i \geq 1$ ). Since  $d(w, z) < \varepsilon$  and  $d(z, g^l(x)) < \delta < \tilde{\varepsilon}/2$ , we have  $d(g^l(x), w) < \tilde{\varepsilon}$ . Therefore  $d(g'^{-i}(w), g'^{-i}(g^l(x))) < \tilde{\varepsilon}$  for all  $i \geq 0$ , and so  $w \in C_{\tilde{\varepsilon}}^u(g^l(x))$  by claim 1.

Obviously, there is  $\tilde{k} = \tilde{k}(z) > 0$  such that  $g'^{n\tilde{k}}(z) \in V_0(p) = \bigcup_{y \in F^u(p, g')} \tilde{W}_{\varepsilon_3}^s(y, g')$ . By the choice of  $\varepsilon$  and by the definition of  $F^u(p, g)$  we have  $B_\varepsilon(g'^{n\tilde{k}}(z)) \cap W_{\varepsilon_3}^s(p, g') = \emptyset$ . However,  $w \in C_\varepsilon^u(g^l(x)) \subset W_{\varepsilon_3}^s(p, g')$  implies  $(g'^{n\tilde{k}}(w) \in W_{\varepsilon_3}^s(p, g'))$ . Thus  $g'^{n\tilde{k}}(w) \in B_\varepsilon(g'^{n\tilde{k}}(z)) \cap W_{\varepsilon_3}^s(p, g') \neq \emptyset$  (since  $d(g'^{n\tilde{k}}(z), g'^{n\tilde{k}}(w)) < \varepsilon$ ). This is a contradiction and so the lemma is proved for case 1.

For case 2, take  $k_1, k_0$  such that  $k \geq k_1$  implies  $C_\varepsilon^u(g^l(x)) \cap V_k(p) \neq \emptyset$ . By the choice of  $\tilde{\varepsilon}$ ,

$$\pi\left(B_{\tilde{\varepsilon}}(g'^{nk}(C_{\tilde{\varepsilon}}^u(g'^l(x)) \cap V_k(p)))\right) \not\subset F^u(p, g')$$

for all  $k \geq k_1$  since  $\dim \pi(C_{\tilde{\varepsilon}}^u(g'^l(x))) < \dim W_{\varepsilon_3}^u(p, g')$  (see (2)). To simplify we write

$$W_k(p) = \bigcup_{y \in g'^{-nk}(\pi(B_{\tilde{\varepsilon}}(g'^{nk}(C_{\tilde{\varepsilon}}^u(g'^l(x)) \cap V_k(p))))} \tilde{W}_{\varepsilon_3}^s(y, g'),$$


$$W(p) = \left( \bigcup_{k \geq k_1} W_k(p) \right) \cup W_{\varepsilon_3}^s(p, g').$$

Then  $W(p) \subset \Gamma(p)$  and

$$\pi(W(p)) = \left( \bigcup_{k \geq k_1} \pi(W_k(p)) \right) \cup \{p\}$$

$$= \left( \bigcup_{k \geq k_1} g'^{-nk}(\pi(B_{\tilde{\varepsilon}}(g'^{nk}(C_{\tilde{\varepsilon}}^u(g'^l(x)) \cap V_k(p)))) \right) \cup \{p\}$$

is not a neighborhood of  $p$  in  $W_{\varepsilon_3}^u(p, g')$ .



**Claim 2.** Put  $\varepsilon = \tilde{\varepsilon}/2$  and let  $\delta = \delta(\varepsilon, g') < \varepsilon$  be the number in the definition of POTP of  $g'$ . Then we have  $B_\delta(g'^l(x)) \subset W(p)$ .

For every  $z \in B_\delta(g^l(x)) \setminus W_{\varepsilon_0}^s(p, g)$ , there exists  $w \in M$  such that  $d(g^i(w), g^i(z)) < \varepsilon$  and  $d(g'^{-i-1}(w), g'^{-i-1}(g^l(x))) < \varepsilon$  for all  $i \geq 0$  since

$$\{\dots, g'^{-1}(x), x, g'(x), \dots, \\ g'^{l-1}(x), z, g'(z), g'^2(z), \dots\}$$

is a  $\delta$ -pseudo-orbit of  $g'$ . Thus  $d(g'^{-i}(w), g'^{-i}(g^l(x))) < \tilde{\varepsilon}$  for all  $i \geq 0$  (since  $d(g^l(x), w) \leq d(g^l(x), z) + d(z, w) < \varepsilon + \delta < \tilde{\varepsilon}$ , and so  $w \in C_{\tilde{\varepsilon}}^u(g^l(x))$  by claim 1. Fix  $\tilde{k} = \tilde{k}(w) \geq k_1$  such that  $w \in V_{\tilde{k}}(p) \cap C_{\tilde{\varepsilon}}^u(g^l(x))$ . Then  $g'^{n\tilde{k}}(z) \in B_\varepsilon(V_0(p) \cap g'^{n\tilde{k}}(C_{\tilde{\varepsilon}}^u(g^l(x))))$  since  $d(g'^{n\tilde{k}}(w), g'^{n\tilde{k}}(z)) < \varepsilon$ . Thus we have  $z \in W_{\tilde{k}}(p) \subset W(p)$ .

By claim 2 we have  $\pi(B_\delta(g^l(x))) \subset \pi(W(p))$ . If we establish that  $\pi(B_\delta(g^l(x)))$  is a neighborhood of  $p$  in  $W_{\varepsilon_3}^u(p, g')$ , then we get a contradiction and therefore the proof of this lemma is completed.

If  $\pi(B_\delta(g^l(x)))$  is not a neighborhood of  $p$  in  $W_{\varepsilon_3}^u(p, g')$ , then for every  $i > 0$  there is  $y_i \in W_{\varepsilon_3}^u(p, g')$  such that  $y_i \notin \pi(B_\delta(g^l(x)))$  and  $d(y_i, p) < \frac{1}{i}$ . Since  $\tilde{W}_{\varepsilon_3}^s(y_i, g') \rightarrow W_{\varepsilon_3}^s(p, g')$  as  $i \rightarrow \infty$ ,

$$\tilde{W}_{\varepsilon_3}^s(y_i, g') \cap B_\delta(g^l(x)) \neq \emptyset$$

for sufficiently large  $i$  and thus  $y_i \in \pi(B_\delta(g^l(x)))$ . This is a contradiction and so  $\pi(W(p))$  is a neighborhood of  $p$  in  $W_{\varepsilon_3}^u(p, g')$ . For any case lemma 5 was proved.

The proof of the transversality at  $x$  for case  $1 \leq \text{Ind } \Lambda_1(f) \leq \dim M - 2$  follows from lemma 5. Indeed, since  $\exp_p(E'_{v_2}(g^l(x)))$  meets transversely  $W_{\varepsilon_3}^s(p, g')$  at  $g^l(x)$ , we have

$$T_{g^l(x)}M = T_{g^l(x)}\exp_p(E'_{v_2}(g^l(x))) + T_{g^l(x)}W_{\varepsilon_3}^s(p, g') \\ = T_{g^l(x)}W^u(g^l(x), g') + T_{g^l(x)}W_{\varepsilon_3}^s(p, g')$$

by (4). Thus

$$T_xM = T_xW^s(p, g') + T_xW^u(q, g') \\ = T_xW^s(x, g) + T_xW^u(x, g) \\ = T_xW^s(x, f) + T_xW^u(x, f).$$

Therefore the proof of the proposition is completed.

## References

- [1] N. Aoki: *The set of Axiom A diffeomorphisms with no cycle*, Bol. Soc. Bras. Mat. 23-New Series (1992), 21–65.
- [2] J. Franks: *Necessary conditions for stability of diffeomorphisms*, Trans. A.M.S. **158** (1971), 301–308.
- [3] M. Hirsch and C. Pugh: *Stable manifold and hyperbolic sets*, in Global Analysis, Proc. Sympos. Pure Math., A.M.S. **14** (1970), 133–163.
- [4] K. Moriyasu: *The topological stability of diffeomorphisms*, Nagoya Math. J. **123** (1991), 91–102.
- [5] C. Robinson: *Stability theorems and hyperbolicity in dynamical systems*, Rocky Mountain J. Math. **7** (1977), 425–437.
- [6] K. Sakai: *The  $C^1$  uniform pseudo-orbit tracing property*, Tokyo J. Math. **15** (1992), 99–109.
- [7] K. Sakai: *Diffeomorphisms with pseudo orbit tracing property*, Nagoya Math. J. **126** (1992), 125–140.
- [8] S. Smale: *The  $\Omega$ -stability theorem*, in Global Analysis Proc. A.M.S. **14** (1970), 289–297.

Department of Mathematics  
Kanagawa University  
Rokkakubashi, Kanagawa-ku  
Yokohama 221, Japan