OUKA
The University of Osaka
Institutional Knowledge Archive

Title	Hybrid mean value results for a generalization on a problem of D.H.Lehmer and hyper-Kloosterman sums
Author(s)	Liu, Huaning; Zhang, Wenpeng
Citation	Osaka Journal of Mathematics. 2007, 44(3), p. 615-637
Version Type	VoR
URL	https://doi.org/10.18910/9619
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Liu, H. and Zhang, W.
Osaka J. Math.
44 (2007), 615–637

HYBRID MEAN VALUE RESULTS FOR A GENERALIZATION ON A PROBLEM OF D.H. LEHMER AND HYPER-KLOOSTERMAN SUMS

HUANING LIU and WENPENG ZHANG

(Received February 23, 2006, revised October 23, 2006)

Abstract

The main purpose of this paper is by using the Fourier expansion for character sums and the mean value theorems of Dirichlet L-functions to give some hybrid mean value results for a generalization on a problem of D.H. Lehmer and hyper-Kloosterman sums.

1. Introduction

Let $q > 2$ and c be two integers with $(c, q) = 1$. For each integer a with $1 \le a \le q$ and $(a, q) = 1$, we know that there exists one and only one b with $1 \le b \le q$ such that $ab \equiv c \bmod q$. Let

$$L(q, k, c) = \sideset{}{'}\sum_{\substack{a=1 \\ ab \equiv c \bmod q}}^{q} \sideset{}{'}\sum_{b=1}^{q} (a - b)^{2k},$$

where $\sum_a'$ denotes the summation over all a such that $(a, q) = 1$. In reference [1], the second author used the estimates for Kloosterman sums and trigonometric sums to obtain a sharp asymptotic formula for $L(q, k, c)$, and prove the following:

Proposition 1. *Let $q > 2$ and c be two integers with $(c, q) = 1$. Then for any positive integer k, we have the asymptotic formula*

$$L(q, k, c) = \frac{1}{(2k + 1)(k + 1)} \phi(q) q^{2k} + O(4^k q^{(4k+1)/2} d^2(q) \ln^2 q),$$

where $\phi(q)$ is the Euler function, and $d(q)$ is the divisor function.

2000 Mathematics Subject Classification. 11L05.

Supported by the National Natural Science Foundation of China under Grant No. 60472068 and No. 10671155; Natural Science Foundation of Shaanxi province of China under Grant No. 2006A04; and the Natural Science Foundation of the Education Department of Shaanxi Province of China under Grant No. 06JK168.

The error terms in Proposition 1 is best possible. In fact for $k = 1$, let

$$L(q, 1, c) = \frac{1}{6}\phi(q)q^2 + \frac{1}{3}q\prod_{p|q}(1 - p) + G(q, c),$$

where $\prod_{p|q}$ denotes the product over all distinct prime divisors of q, the second author [2] used the properties of Dedekind sums and Cochrane sums to give a sharp mean value formula for $G(q, c)$. That is the following:

Proposition 2. *For any integer $q > 2$, we have the asymptotic formula*

$$\sum_{c=1}^{q}{}' G^2(q, c) = \frac{5}{36}q^3\phi^3(q)\prod_{p^\alpha \| q}\frac{(p+1)^3/(p(p^2+1)) - 1/(p^{3\alpha-1})}{1 + 1/p + 1/p^2} + O\left(q^5\exp\left(\frac{4\ln q}{\ln\ln q}\right)\right),$$

where $\exp(y) = e^y$, $\prod_{p^\alpha \| q}$ *denotes the product over all prime divisors p of q with $p^\alpha | q$ and $p^{\alpha+1} \nmid q$.*

Let $M(q, c)$ be the number of cases in which a and b are of opposite parity. That is,

$$M(q, c) = \sum_{\substack{a=1 \\ ab\equiv c \bmod q \\ 2\nmid a+b}}^{q}{}' \sum_{b=1}^{q}{}' 1.$$

For $q = p$ an odd prime and $c = 1$, D.H. Lehmer [3] asked us to find $M(p, 1)$ or at least to say something nontrivial about it. For the sake of simplicity, we call such a number as a D.H. Lehmer Number. In references [4] and [5], the second author proved that

$$M(q, 1) = \frac{1}{2}\phi(q) + O(q^{1/2}d^2(q)\ln^2 q). \tag{1}$$

For any nonnegative integer n, let

$$M(q, 1, n) = \sum_{\substack{a=1 \\ ab\equiv 1 \bmod q \\ 2\nmid a+b}}^{q}{}' \sum_{b=1}^{q}{}' (a - b)^{2n},$$

the second author [6] also proved the following asymptotic formula:

$$M(q, 1, n) = \frac{1}{(2n+1)(2n+2)}\phi(q)q^{2n} + O(4^n q^{2n+1/2}d^2(q)\ln^2 q).$$

For any fixed positive integer c with $(c, q) = 1$, define

$$F(q, c) = M(q, c) - \frac{1}{2}\phi(q).$$

Then the second author showed in [7] and [8] that for any odd number $q > 2$,

$$\sum_{c=1}^{q}{}' |F(q, c)|^2 = \frac{3}{4}\phi^2(q) \prod_{p^\alpha \| q} \frac{(p+1)^3/(p(p^2+1)) - 1/p^{3\alpha-1}}{1 + 1/p + 1/p^2} + O\left(q \exp\left(\frac{4 \ln q}{\ln \ln q}\right)\right).$$

This proved that the error terms in (1) is also best possible.

In [9], the second author found that there exists some close relation between the error terms $F(q, c)$ and the classical Kloosterman sums:

$$K(m, n; q) = \sum_{b=1}^{q}{}' \mathrm{e}\left(\frac{mb + n\overline{b}}{q}\right),$$

where $\mathrm{e}(y) = \mathrm{e}^{2\pi i y}$, $\overline{b}$ is defined by the equation $b\overline{b} \equiv 1 \mod q$, and obtained the following hybrid mean value formula:

$$\sum_{c=1}^{q}{}' F(q, c)K(\overline{4}c, 1; q) = \frac{4}{\pi^2} q\phi(q) \prod_{p \| q} \left(1 - \frac{1}{p(p-1)}\right) + O(q^{3/2+\epsilon}),$$

where ϵ is any fixed positive number.

In [10], Mordell introduced the hyper-Kloosterman sums as follows:

$$K(h, k, q) = \sum_{\substack{a_1, \cdots, a_k \bmod q \\ (a_1, q) = \cdots = (a_k, q) = 1}} \mathrm{e}\left(\frac{a_1 + \cdots + a_k + h\overline{a}_1 \cdots \overline{a}_k}{q}\right),$$

which is the high-dimensional generalization of the Kloosterman sums. Some applications of the hyper-Kloosterman sums were found in the estimation of Fourier coefficients of Maass forms [11] and the work on Selberg's eigenvalue conjecture [12]. Moreover, there exists some interesting connections between the hyper-Kloosterman sums and the Heibronn sums (see reference [13]).

Now we consider a generalization on this problem of D.H. Lehmer. For any integer $k \geq 1$, let

$$N(q, k, c) = \sum_{a_1=1}^{q}{}' \cdots \sum_{a_k=1}^{q}{}' \sum_{b=1}^{q}{}' (a_1 + \cdots + a_k - b)^2$$
$$a_1 \cdots a_k b \equiv c \bmod q$$
$$2 \nmid a_1 + \cdots + a_k + b$$

and

$$E(q, k, c) = N(q, k, c) - \frac{(3k^2 - 5k + 4)}{24}\phi^k(q)q^2 - \frac{(k+1)}{12}\phi^{k-1}(q)q \prod_{p|q}(1 - p).$$

In this paper, we use the Fourier expansion for character sums and the mean value theorems of Dirichlet L-functions to study the hybrid mean value of $E(q, k, c)$ and the hyper-Kloosterman sums, and give an interesting mean value formula. That is, we shall prove the following:

Theorem. *For any odd number $q \geq 3$ and integer $k \geq 1$, we have the asymptotic formula*

$$\sum_{c=1}^{q}{}' E(q, k, c)K(\overline{2}^{k+1}c, k, q) = \frac{c_k q^{k+2}\phi(q)}{\pi^{k+3}} \prod_{p\|q}\left(1 - \frac{p^k - 1}{p^k(p-1)^2}\right) + O(q^{k+5/2+\epsilon}),$$

where ϵ is any fixed positive number, $\prod_{p\|q}$ denotes the product over all prime divisors p of q with $p|q$ and $p^2 \nmid q$, and

$$c_k = \begin{cases} -6, & \text{if} \quad k = 1, \\ i^{k+3}2^{2k-2}[\pi^2(k^2 - k + 2) - 8(k+1)], & \text{otherwise.} \end{cases}$$

2. Several lemmas

To complete the proof of the theorem, we need the following lemmas.

Lemma 1. *Let $q \geq 3$ be an odd number. Then for any positive integer c with $(c, q) = 1$, we have*

$$\begin{aligned} E(q, 1, c) = {} & \frac{1}{\phi(q)} \sum_{\substack{\chi(-1)=1 \\ \chi \neq \chi_0}} \overline{\chi}(c)\left(\sum_{a=1}^{q}(-1)^a a\chi(a)\right)^2 \\ & + \frac{1}{\phi(q)} \sum_{\chi(-1)=-1} \overline{\chi}(c)(4\chi(4) - 4\chi(2))\left(\sum_{a=1}^{q} a\chi(a)\right)^2 \\ & - \frac{2}{\phi(q)q} \sum_{\chi(-1)=-1} \overline{\chi}(c)(1 - 2\chi(2))\left(\sum_{a=1}^{q} a\chi(a)\right)\left(\sum_{b=1}^{q}(-1)^b b^2\chi(b)\right) + O(q), \end{aligned}$$

where $\sum_{\substack{\chi(-1)=1 \\ \chi \neq \chi_0}}$ denotes the summation over all non-principal even characters modulo q, and $\sum_{\chi(-1)=-1}$ denotes the summation over all odd characters modulo q.

Proof. From the definition of $N(q, 1, c)$ we can get

$$N(q, 1, c) = \sum_{\substack{a=1 \\ ab\equiv c \bmod q \\ 2\nmid a+b}}^{q}{}' \sum_{b=1}^{q}{}' (a - b)^2 = \frac{1}{2} \sum_{\substack{a=1 \\ ab\equiv c \bmod q}}^{q}{}' \sum_{b=1}^{q}{}' [1 - (-1)^{a+b}](a - b)^2$$

$$= \frac{1}{2}\sideset{}{'}\sum_{\substack{a=1\\ ab\equiv c \bmod q}}^{q}\sideset{}{'}\sum_{b=1}^{q}(a-b)^2 - \frac{1}{2}\sideset{}{'}\sum_{\substack{a=1\\ ab\equiv c \bmod q}}^{q}\sideset{}{'}\sum_{b=1}^{q}(-1)^{a+b}(a-b)^2.$$

By the orthogonality relation for character sums mod q we easily deduce

$$\begin{aligned} &N(q, 1, c)\\ &= \frac{1}{12}\phi(q)q^2 + \frac{1}{6}q\prod_{p|q}(1-p) - \frac{1}{\phi(q)}\sum_{\chi(-1)=-1}\overline{\chi}(c)\left(\sum_{a=1}^{q}a\chi(a)\right)^2\\ &\quad - \frac{1}{\phi(q)}\sum_{\chi\neq\chi_0}\overline{\chi}(c)\left[\left(\sum_{a=1}^{q}(-1)^a a^2\chi(a)\right)\left(\sum_{b=1}^{q}(-1)^b\chi(b)\right) - \left(\sum_{a=1}^{q}(-1)^a a\chi(a)\right)^2\right]\\ &\quad + O(q), \end{aligned}$$

where we have used the identities (see reference [14])

$$\sideset{}{'}\sum_{a=1}^{q}a^2 = \frac{1}{3}\phi(q)q^2 + \frac{1}{6}q\prod_{p|q}(1-p), \qquad \sideset{}{'}\sum_{a=1}^{q}a = \frac{\phi(q)q}{2} \tag{2}$$

and

$$\sideset{}{'}\sum_{a=1}^{q}(-1)^a = 0, \qquad \sideset{}{'}\sum_{a=1}^{q}(-1)^a a = -\frac{1}{2}\prod_{p|q}(1-p), \qquad \sideset{}{'}\sum_{a=1}^{q}(-1)^a a^2 = -\frac{1}{2}q\prod_{p|q}(1-p)$$

for any odd number $q \geq 3$.

Note that if $\chi(-1) = 1$ then

$$\sum_{b=1}^{q}(-1)^b\chi(b) = 0, \tag{3}$$

and if $\chi(-1) = -1$, then

$$\sum_{a=1}^{q}(-1)^a a\chi(a) = \frac{q}{2}\sum_{a=1}^{q}(-1)^a\chi(a), \qquad \sum_{b=1}^{q}(-1)^b\chi(b) = 2\chi(2)\sum_{b=1}^{(q-1)/2}\chi(b). \tag{4}$$

From [15] we also know that for any odd character $\chi \bmod q$, we have

$$(1-2\chi(2))\sum_{c=1}^{q}c\chi(c) = \chi(2)q\sum_{c=1}^{(q-1)/2}\chi(c). \tag{5}$$

So from the above formulae we can have

$$E(q,1,c)=\frac{1}{\phi(q)}\sum_{\substack{\chi(-1)=1\\ \chi\neq\chi_0}}\overline{\chi}(c)\left(\sum_{a=1}^{q}(-1)^a a\chi(a)\right)^2$$
$$+\frac{1}{\phi(q)}\sum_{\chi(-1)=-1}\overline{\chi}(c)(4\chi(4)-4\chi(2))\left(\sum_{a=1}^{q}a\chi(a)\right)^2$$
$$-\frac{2}{\phi(q)q}\sum_{\chi(-1)=-1}\overline{\chi}(c)(1-2\chi(2))\left(\sum_{a=1}^{q}a\chi(a)\right)\left(\sum_{b=1}^{q}(-1)^b b^2\chi(b)\right)+O(q).$$

This proves Lemma 1. □

Lemma 2. *Let $q\geq 3$ be an odd number and $k\geq 2$ be an integer. Then for any positive integer c with $(c,q)=1$, we have the identity*

$$E(q,k,c)=\frac{-2^{k-2}}{\phi(q)q^{k-1}}\sum_{\chi(-1)=-1}\overline{\chi}(c)(1-2\chi(2))^k\left(\sum_{a=1}^{q}a\chi(a)\right)^k$$
$$\times\left[\frac{2(k+1)}{q}\left(\sum_{b=1}^{q}(-1)^b b^2\chi(b)\right)+k(k-3)(1-2\chi(2))\left(\sum_{b=1}^{q}b\chi(b)\right)\right].$$

Proof. From the definition of $N(q,k,c)$ we can get

$$N(q,k,c)=\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}(a_1+\cdots+a_k-b)^2$$
$$\substack{a_1\cdots a_k b\equiv c \bmod q\\ 2\nmid a_1+\cdots+a_k+b}$$
$$=\frac{1}{2}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}[1-(-1)^{a_1+\cdots+a_k+b}](a_1+\cdots+a_k-b)^2$$
$$\substack{a_1\cdots a_k b\equiv c \bmod q}$$
$$=\frac{1}{2}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}(a_1+\cdots+a_k-b)^2$$
$$\substack{a_1\cdots a_k b\equiv c \bmod q}$$
$$-\frac{1}{2}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}(-1)^{a_1+\cdots+a_k+b}(a_1+\cdots+a_k-b)^2$$
$$\substack{a_1\cdots a_k b\equiv c \bmod q}$$
$$:=P(q,k,c)+E(q,k,c).$$

Then by (2) and the properties of character sums mod q we have

$$P(q,k,c)=\frac{1}{2}\sideset{}{'}\sum_{\substack{a_1=1\\ a_1\cdots a_k b\equiv c \bmod q}}^{q}\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}(a_1+\cdots+a_k-b)^2$$

$$=\frac{1}{2\phi(q)}\sum_{\chi \bmod q}\overline{\chi}(c)\sum_{a_1=1}^{q}\cdots\sum_{a_k=1}^{q}\sum_{b=1}^{q}\chi(a_1)\cdots\chi(a_k)\chi(b)(a_1+\cdots+a_k-b)^2$$

$$=\frac{1}{2\phi(q)}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}(a_1+\cdots+a_k-b)^2$$

$$=\frac{1}{2\phi(q)}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}\left(\sum_{1\le i\le k}a_i^2+b^2+2\sum_{1\le i<j\le k}a_ia_j-2\sum_{1\le i\le k}a_ib\right)$$

$$=\frac{(k+1)}{2\phi(q)}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}b^2$$

$$+\frac{1}{\phi(q)}((k-1)+(k-2)+\cdots+1-k)\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}a_1b$$

$$=\frac{(k+1)}{2\phi(q)}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}b^2+\frac{k(k-3)}{2\phi(q)}\sideset{}{'}\sum_{a_1=1}^{q}\cdots\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}a_1b$$

$$=\frac{(k+1)\phi^{k-1}(q)}{2}\left[\frac{1}{3}\phi(q)q^2+\frac{1}{6}q\prod_{p\mid q}(1-p)\right]+\frac{k(k-3)}{8}\phi^k(q)q^2$$

$$=\frac{(3k^2-5k+4)}{24}\phi^k(q)q^2+\frac{(k+1)}{12}\phi^{k-1}(q)q\prod_{p\mid q}(1-p).$$

On the other hand, from (3), (4) and (5) we also have

$$E(q,k,c)=-\frac{1}{2}\sideset{}{'}\sum_{\substack{a_1=1\\ a_1\cdots a_k b\equiv c \bmod q}}^{q}\sideset{}{'}\sum_{a_k=1}^{q}\sideset{}{'}\sum_{b=1}^{q}(-1)^{a_1+\cdots+a_k+b}(a_1+\cdots+a_k-b)^2$$

$$=-\frac{1}{2\phi(q)}\sum_{\chi \bmod q}\overline{\chi}(c)\sum_{a_1=1}^{q}\cdots\sum_{a_k=1}^{q}\sum_{b=1}^{q}(-1)^{a_1+\cdots+a_k+b}$$

$$\times\chi(a_1)\cdots\chi(a_k)\chi(b)(a_1+\cdots+a_k-b)^2$$

$$= \frac{-1}{2\phi(q)} \sum_{\chi(-1)=-1} \overline{\chi}(c) \Bigg[(k+1) \Bigg(\sum_{a=1}^{q} (-1)^a \chi(a) \Bigg)^k \Bigg(\sum_{b=1}^{q} (-1)^b b^2 \chi(b) \Bigg) + k(k-3) \Bigg(\sum_{a=1}^{q} (-1)^a a \chi(a) \Bigg)^2 \Bigg(\sum_{b=1}^{q} (-1)^b \chi(b) \Bigg)^{k-1} \Bigg]$$

$$= \frac{-2^{k-2}}{\phi(q) q^{k-1}} \sum_{\chi(-1)=-1} \overline{\chi}(c)(1-2\chi(2))^k \Bigg(\sum_{a=1}^{q} a \chi(a) \Bigg)^k \times \Bigg[\frac{2(k+1)}{q} \Bigg(\sum_{b=1}^{q} (-1)^b b^2 \chi(b) \Bigg) + k(k-3)(1-2\chi(2)) \Bigg(\sum_{b=1}^{q} b \chi(b) \Bigg) \Bigg].$$

This completes the proof of Lemma 2. □

Lemma 3. *Let χ be a primitive character modulo m, and let $m \geq 3$ be an odd number. Then we have*

$$\sum_{a=1}^{m} a \chi(a) = \frac{mi}{\pi} \tau(\chi) L(1, \overline{\chi}), \quad \text{if} \quad \chi(-1) = -1;$$

$$\sum_{a=1}^{m} (-1)^a a \chi(a) = \frac{m \tau(\chi)(1-4\chi(2))}{\pi^2} L(2, \overline{\chi}) + O(m), \quad \text{if} \quad \chi(-1) = 1;$$

$$\sum_{a=1}^{m} (-1)^a a^2 \chi(a) = \frac{m^2(1-2\chi(2))i}{\pi} \tau(\chi) L(1, \overline{\chi}) + \frac{m^2(8\chi(2)-1)i}{\pi^3} \tau(\chi) L(3, \overline{\chi}) + O(m^2), \quad \text{if} \quad \chi(-1) = -1,$$

where $L(1, \chi)$ is the Dirichlet L-function corresponding to χ, $\tau(\chi) = \sum_{a=1}^{m} \chi(a) \mathrm{e}(a/m)$ is the Gauss sum, and $|\tau(\chi)| = \sqrt{m}$.

Proof. From Theorem 12.11 and Theorem 12.20 of [14] we know that if χ is a primitive character modulo m with $\chi(-1) = -1$, then

$$\frac{1}{m} \sum_{b=1}^{m} b \chi(b) = \frac{i}{\pi} \tau(\chi) L(1, \overline{\chi}), \tag{6}$$

so we get the first formula.

Now we prove the third formula, since similarly we can deduce the second one. For any odd primitive character χ modulo m, we have

$$\begin{aligned}\sum_{a=1}^{m}(-1)^a a^2\chi(a) &= \sum_{\substack{a=1\\2|a}}^{m} a^2\chi(a) - \sum_{\substack{a=1\\2\nmid a}}^{m} a^2\chi(a) = \sum_{\substack{a=1\\2|a}}^{m} a^2\chi(a) - \sum_{\substack{a=1\\2|a}}^{m}(m-a)^2\chi(m-a)\\ &= 2\sum_{\substack{a=1\\2|a}}^{m} a^2\chi(a) - 2m\sum_{\substack{a=1\\2|a}}^{m} a\chi(a) + m^2\sum_{\substack{a=1\\2|a}}^{m}\chi(a).\end{aligned} \tag{7}$$

Note that

$$\sum_{a=1}^{m} a\chi(a) = \sum_{\substack{a=1\\2|a}}^{m} a\chi(a) + \sum_{\substack{a=1\\2|a}}^{m}(m-a)\chi(m-a) = 2\sum_{\substack{a=1\\2|a}}^{m} a\chi(a) - m\sum_{\substack{a=1\\2|a}}^{m}\chi(a),$$

so from (6) and (7) we can get

$$\begin{aligned}\sum_{a=1}^{m}(-1)^a a^2\chi(a) &= 2\sum_{\substack{a=1\\2|a}}^{m} a^2\chi(a) - m\sum_{a=1}^{m} a\chi(a)\\ &= 8\chi(2)\sum_{1\le a\le m/2} a^2\chi(a) - \frac{m^2 i}{\pi}\tau(\chi)L(1,\overline{\chi}).\end{aligned} \tag{8}$$

Notice the Fourier expansion for character sums which was first given by Pólya [16]:

$$\sum_{0<n\le my}\chi(n) = \begin{cases}\dfrac{\tau(\chi)}{\pi}\displaystyle\sum_{n=1}^{\infty}\frac{\overline{\chi}(n)\sin(2\pi ny)}{n} + O(1), & \text{if}\quad \chi(-1)=1;\\[2ex] \dfrac{\tau(\chi)L(1,\overline{\chi})}{\pi i} - \dfrac{\tau(\chi)}{\pi i}\displaystyle\sum_{n=1}^{\infty}\frac{\overline{\chi}(n)\cos(2\pi ny)}{n} + O(1), & \text{if}\quad \chi(-1)=-1,\end{cases}$$

where χ is a primitive character modulo m, and $y > 0$ is a real number. Then by Abel's identity we have

$$\begin{aligned}\sum_{0<n\le m/2} n^2\chi(n) &= \frac{m^2}{4}\sum_{0<n\le m/2}\chi(n) - 2\int_0^{m/2} u\sum_{0<n\le u}\chi(n)\,\mathrm{d}u\\ &= \frac{m^2(\overline{\chi}(2)-2)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) - 2\int_0^{m/2} u\sum_{0<n\le u}\chi(n)\,\mathrm{d}u + O(m^2)\\ &= \frac{m^2(\overline{\chi}(2)-2)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) - 2m^2\int_0^{1/2} s\sum_{0<n\le ms}\chi(n)\,\mathrm{d}s + O(m^2)\end{aligned}$$

$$
\begin{aligned}
&= \frac{m^2(\overline{\chi}(2)-2)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) \\
&\quad - 2m^2\int_0^{1/2} s\left[\frac{\tau(\chi)L(1,\overline{\chi})}{\pi i} - \frac{\tau(\chi)}{\pi i}\sum_{n=1}^{\infty}\frac{\overline{\chi}(n)\cos(2\pi ns)}{n} + O(1)\right]\mathrm{d}s \\
&\quad + O(m^2) \\
&= \frac{m^2(\overline{\chi}(2)-1)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) + 2m^2\frac{\tau(\chi)}{\pi i}\sum_{n=1}^{\infty}\frac{\overline{\chi}(n)}{n}\int_0^{1/2} s\cos(2\pi ns)\,\mathrm{d}s \\
&\quad + O(m^2) \\
&= \frac{m^2(\overline{\chi}(2)-1)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) + \frac{m^2\tau(\chi)}{2\pi^3 i}\sum_{n=1}^{\infty}\frac{\overline{\chi}(n)(\cos(\pi n)-1)}{n^3} + O(m^2) \\
&= \frac{m^2(\overline{\chi}(2)-1)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) - \frac{m^2\tau(\chi)}{\pi^3 i}\sum_{\substack{n=1\\ 2\nmid n}}^{\infty}\frac{\overline{\chi}(n)}{n^3} + O(m^2) \\
&= \frac{m^2(\overline{\chi}(2)-1)i}{4\pi}\tau(\chi)L(1,\overline{\chi}) + \frac{m^2(\overline{\chi}(2)-8)}{8\pi^3 i}\tau(\chi)L(3,\overline{\chi}) + O(m^2).
\end{aligned}
$$

Hence from (8) we have

$$
\sum_{a=1}^{m}(-1)^a a^2\chi(a) = \frac{m^2(1-2\chi(2))i}{\pi i}\tau(\chi)L(1,\overline{\chi}) + \frac{m^2(8\chi(2)-1)i}{\pi^3}\tau(\chi)L(3,\overline{\chi}) + O(m^2).
$$

This proves Lemma 3. □

Lemma 4. *Suppose that χ is an even character modulo q, generated by the primitive character χ_m modulo m, and $q \geq 3$ is an odd number. Let l be the largest divisor of q with $(l, m) = 1$. Then we have*

$$
\sum_{a=1}^{q}(-1)^a a\chi(a) = \frac{m\tau(\chi_m)(1-4\chi_m(2))}{\pi^2}L(2,\overline{\chi}_m)\left(\sum_{d|l} d\mu(d)\chi_m(d)\right) + O(q),
$$

where $\mu(n)$ is the Möbius function.

Proof. Note that m and l both are odd numbers, then from (3) we have

$$
\begin{aligned}
\sum_{a=1}^{q}(-1)^a a\chi(a) &= \sum_{i=0}^{(q/ml)-1}\sum_{j=1}^{ml}(-1)^{iml+j}(iml+j)\chi(iml+j) \\
&= \sum_{i=0}^{(q/ml)-1}(-1)^i\sum_{j=1}^{ml}(-1)^j(iml+j)\chi(iml+j) = \sum_{i=0}^{(q/ml)-1}(-1)^i\sum_{j=1}^{ml}(-1)^j j\chi(j)
\end{aligned}
$$

$$= \sum_{j=1}^{ml} (-1)^j j\chi(j) = \sum_{\substack{a=1 \\ (a,l)=1}}^{ml} (-1)^a a\chi_m(a) = \sum_{a=1}^{ml} (-1)^a a\chi_m(a) \sum_{\substack{d|a \\ d|l}} \mu(d)$$

$$= \sum_{d|l} \mu(d) \sum_{\substack{a=1 \\ d|a}}^{lm} (-1)^a a\chi_m(a) = \sum_{d|l} \mu(d) d\chi_m(d) \sum_{b=1}^{lm/d} (-1)^{db} b\chi_m(b).$$

Since d is odd, we have $(-1)^{db} = (-1)^b$. Therefore

$$\begin{aligned}
\sum_{a=1}^{q} (-1)^a a\chi(a) &= \sum_{d|l} \mu(d) d\chi_m(d) \sum_{b=1}^{lm/d} (-1)^b b\chi_m(b) \\
&= \sum_{d|l} \mu(d) d\chi_m(d) \sum_{i=0}^{(l/d)-1} \sum_{j=1}^{m} (-1)^{im+j} (im + j)\chi_m(im + j) \\
&= \sum_{d|l} \mu(d) d\chi_m(d) \sum_{i=0}^{(l/d)-1} (-1)^i \sum_{j=1}^{m} (-1)^j (im + j)\chi_m(j) \\
&= \sum_{d|l} \mu(d) d\chi_m(d) \sum_{j=1}^{m} (-1)^j j\chi_m(j).
\end{aligned}$$

Note that

$$\left| \sum_{d|l} \mu(d) d\chi_m(d) \right| = \left| \prod_{p|l} (1 - p\chi_m(p)) \right| < \prod_{p|l} p \le l,$$

so from Lemma 3 we get

$$\sum_{a=1}^{q} (-1)^a a\chi(a) = \frac{m\tau(\chi_m)(1 - 4\chi_m(2))}{\pi^2} L(2, \overline{\chi}_m) \left(\sum_{d|l} d\mu(d)\chi_m(d) \right) + O(q).$$

This completes the proof of Lemma 4. □

Lemma 5. *Suppose that χ is an odd character modulo q, generated by the primitive character χ_m modulo m, and $q \ge 3$ is an odd number. Let l be the largest divisor of q with $(l, m) = 1$. Then we have*

$$\sum_{a=1}^{q} a\chi(a) = \frac{qi}{\pi} \tau(\chi_m) L(1, \overline{\chi}_m) \left(\sum_{d|l} \mu(d)\chi_m(d) \right).$$

Furthermore, for $q = lm$, we also have

$$\sum_{a=1}^{q}(-1)^a a^2\chi(a) = \frac{q^2(1-2\chi_m(2))i}{\pi}\tau(\chi_m)L(1,\overline{\chi}_m)\left(\sum_{d|l}\mu(d)\chi_m(d)\right) + \frac{m^2(8\chi_m(2)-1)i}{\pi^3}\tau(\chi_m)L(3,\overline{\chi}_m)\left(\sum_{d|l}d^2\mu(d)\chi_m(d)\right) + O(q^2).$$

Proof. The first formula can be easily deduced from Lemma 6 of [5] and Lemma 3. Now let $q = lm$, we have

$$\sum_{a=1}^{q}(-1)^a a^2\chi(a) = \sum_{a=1}^{lm}(-1)^a a^2\chi(a) = \sum_{\substack{a=1\\(a,l)=1}}^{lm}(-1)^a a^2\chi_m(a) = \sum_{d|l}\mu(d)\sum_{\substack{a=1\\d|a}}^{lm}(-1)^a a^2\chi_m(a)$$

$$= \sum_{d|l}\mu(d)d^2\chi_m(d)\sum_{b=1}^{lm/d}(-1)^{db}b^2\chi_m(b)$$

$$= \sum_{d|l}d^2\mu(d)\chi_m(d)\sum_{b=1}^{lm/d}(-1)^b b^2\chi_m(b).$$

Since

$$\sum_{b=1}^{lm/d}(-1)^b b^2\chi_m(b) = \sum_{i=0}^{(l/d)-1}\sum_{j=1}^{m}(-1)^{im+j}(im+j)^2\chi_m(im+j)$$

$$= \sum_{i=0}^{(l/d)-1}(-1)^i\sum_{j=1}^{m}(-1)^j(i^2m^2+2imj+j^2)\chi_m(j)$$

$$= m^2\sum_{i=0}^{(l/d)-1}(-1)^i i^2\sum_{j=1}^{m}(-1)^j\chi_m(j) + 2m\sum_{i=0}^{(l/d)-1}(-1)^i i\sum_{j=1}^{m}(-1)^j j\chi_m(j) + \sum_{i=0}^{(l/d)-1}(-1)^i\sum_{j=1}^{m}(-1)^j j^2\chi_m(j)$$

$$= \frac{m^2l(l-d)}{2d^2}\sum_{j=1}^{m}(-1)^j\chi_m(j) + \frac{m(l-d)}{d}\sum_{j=1}^{m}(-1)^j j\chi_m(j) + \sum_{j=1}^{m}(-1)^j j^2\chi_m(j),$$

and note that

$$\left|\sum_{d|l} d^2\mu(d)\chi_m(d)\right| = \left|\prod_{p|l}(1 - p^2\chi_m(p))\right| < \prod_{p|l} p^2 \le l^2,$$

so from (4), (5), (6) and Lemma 3 we have

$$\begin{aligned}\sum_{a=1}^{q}(-1)^a a^2\chi(a) &= \frac{q^2}{2}\left(\sum_{d|l}\mu(d)\chi_m(d)\right)\left(\sum_{j=1}^{m}(-1)^j\chi_m(j)\right)\\ &\quad - \frac{m^2}{2}\left(\sum_{d|l} d^2\mu(d)\chi_m(d)\right)\left(\sum_{j=1}^{m}(-1)^j\chi_m(j)\right)\\ &\quad + \left(\sum_{d|l} d^2\mu(d)\chi_m(d)\right)\left(\sum_{j=1}^{m}(-1)^j j^2\chi_m(j)\right)\\ &= \frac{q^2(1-2\chi_m(2))i}{\pi}\tau(\chi_m)L(1,\overline{\chi}_m)\left(\sum_{d|l}\mu(d)\chi_m(d)\right)\\ &\quad + \frac{m^2(8\chi_m(2)-1)i}{\pi^3}\tau(\chi_m)L(3,\overline{\chi}_m)\left(\sum_{d|l} d^2\mu(d)\chi_m(d)\right) + O(q^2).\end{aligned}$$

This proves Lemma 5. □

Lemma 6. *Let χ be a character modulo q, generated by the primitive character χ_m modulo m. Then we have the identity*

$$\tau(\chi) = \chi_m\left(\frac{q}{m}\right)\mu\left(\frac{q}{m}\right)\tau(\chi_m).$$

Proof. See Lemma 1.3 of reference [17]. □

Lemma 7. *Let m and r be integers with $m \ge 2$ and $(r, m) = 1$, and let χ be a Dirichlet character modulo m. Then we have the identities*

$$\sum_{\chi \bmod m}^{*}\chi(r) = \sum_{d|(m,r-1)}\mu\left(\frac{m}{d}\right)\phi(d)$$

and

$$J(m) = \sum_{d|m}\mu(d)\phi\left(\frac{m}{d}\right),$$

where $\sum^{*}_{\chi \bmod m}$ denotes the summation over all primitive characters modulo m and $J(m)$ denotes the number of primitive characters modulo m.

Proof. This is Lemma 3 of [18]. □

Lemma 8. *Let $q = uv$, where $(u, v) = 1$, u be a square-full number or $u = 1$, v be a square-free number. Then we have the asymptotic formulae*

$$\Psi_1 := \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \mu(d_1)\cdots\mu(d_{k+1}) \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(d_1\cdots d_{k+1}) L^{k+1}(1, \overline{\chi})$$

$$= \frac{q^k \phi^2(q)}{2} \prod_{p\|q} \left(1 - \frac{p^k - 1}{p^k (p-1)^2}\right) + O(q^{k+1+\epsilon});$$

$$\Psi_2 := \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \frac{\mu(d_1)\cdots\mu(d_{k+1})}{d_1\cdots d_{k+1}} \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=1}} \overline{\chi}(d_1\cdots d_{k+1}) L^{k+1}(2, \overline{\chi})$$

$$= \frac{q^k \phi^2(q)}{2} \prod_{p\|q} \left(1 - \frac{p^k - 1}{p^k (p-1)^2}\right) + O(q^{k+1+\epsilon});$$

$$\Psi_3 := \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \frac{\mu(d_1)\cdots\mu(d_{k+1})}{d_{k+1}^2} \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(d_1\cdots d_{k+1}) L^k(1, \overline{\chi}) L(3, \overline{\chi})$$

$$= \frac{q^k \phi^2(q)}{2} \prod_{p\|q} \left(1 - \frac{p^k - 1}{p^k (p-1)^2}\right) + O(q^{k+1+\epsilon}).$$

Proof. We only prove the first formula, since similarly we can get the others. For any non-principal character χ modulo ud, and parameter $N \geq ud$, applying Abel's identity we have

$$L(s, \overline{\chi}) = \sum_{n=1}^{\infty} \frac{\overline{\chi}(n)}{n^s} = \sum_{1\leq n\leq N} \frac{\overline{\chi}(n)}{n^s} + s \int_N^{\infty} \frac{\sum_{N<n\leq y} \overline{\chi}(n)}{y^{s+1}}\, \mathrm{d}y$$

$$= \sum_{1\leq n\leq N} \frac{\overline{\chi}(n)}{n^s} + O\left(\frac{\sqrt{ud}\log(ud)}{N^s}\right).$$

Let $\tau_{k+1}(n)$ be the $(k+1)$-th divisor function (i.e., the number of positive integer solutions of the equation $n_1 n_2 \cdots n_{k+1} = n$). Then we have

$$\Psi_1 = \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \mu(d_1)\cdots\mu(d_{k+1}) \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(d_1\cdots d_{k+1})$$

$$\times \left(\sum_{1\leq n\leq N} \frac{\overline{\chi}(n)}{n} + O\left(\frac{\sqrt{ud}\log(ud)}{N}\right)\right)^{k+1}$$

$$= \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \mu(d_1)\cdots\mu(d_{k+1}) \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(d_1\cdots d_{k+1})$$

$$\times \left(\sum_{1\le n\le N} \frac{\overline{\chi}(n)}{n}\right)^{k+1} + O\left(\frac{q^{k+5/2+\epsilon}\log^k N}{N}\right)$$

$$= \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \mu(d_1)\cdots\mu(d_{k+1}) \sum_{1\le n\le N^{k+1}} \frac{\tau_{k+1}(n)}{n}$$

$$\times \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(d_1\cdots d_{k+1})\overline{\chi}(n) + O\left(\frac{q^{k+5/2+\epsilon}\log^k N}{N}\right)$$

$$= \sum_{d|v} u^{k+1} d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \sum_{1\le n\le N^{k+1}} \frac{\mu(d_1)\cdots\mu(d_{k+1})d_1\cdots d_{k+1}\tau_{k+1}(n)}{d_1\cdots d_{k+1}n}$$

$$\times \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(d_1\cdots d_{k+1})\overline{\chi}(n) + O\left(\frac{q^{k+5/2+\epsilon}\log^k N}{N}\right)$$

$$:= \Omega + O\left(\frac{q^{k+5/2+\epsilon}\log^k N}{N}\right).$$

For $(a, m) = 1$, by Lemma 7 we have

$$\sideset{}{^*}\sum_{\substack{\chi \bmod m \\ \chi(-1)=-1}} \chi(a) = \frac{1}{2}\sideset{}{^*}\sum_{\chi \bmod m} (1-\chi(-1))\chi(a) = \frac{1}{2}\sideset{}{^*}\sum_{\chi \bmod m} \chi(a) - \frac{1}{2}\sideset{}{^*}\sum_{\chi \bmod m} \chi(-a)$$

$$= \frac{1}{2}\sum_{s|(m,a-1)} \mu\left(\frac{m}{s}\right)\phi(s) - \frac{1}{2}\sum_{s|(m,a+1)} \mu\left(\frac{m}{s}\right)\phi(s). \tag{9}$$

Therefore

$$\Omega = \frac{1}{2}\sum_{d|v} u^{k+1} d^{k+1} \sum_{s|ud} \mu\left(\frac{ud}{s}\right)\phi(s)$$

$$\times \sum_{d_1|(v/d)} \cdots \sum_{\substack{d_{k+1}|(v/d) \\ d_1\cdots d_{k+1}n\equiv 1 \bmod s}} \sum_{\substack{1\le n\le N^{k+1} \\ (n,ud)=1}} \frac{\mu(d_1)\cdots\mu(d_{k+1})d_1\cdots d_{k+1}\tau_{k+1}(n)}{d_1\cdots d_{k+1}n}$$

$$- \frac{1}{2}\sum_{d|v} u^{k+1} d^{k+1} \sum_{s|ud} \mu\left(\frac{ud}{s}\right)\phi(s)$$

$$\times \sum_{d_1|(v/d)} \cdots \sum_{\substack{d_{k+1}|(v/d) \\ d_1\cdots d_{k+1}n\equiv -1 \bmod s}} \sum_{\substack{1\le n\le N^{k+1} \\ (n,ud)=1}} \frac{\mu(d_1)\cdots\mu(d_{k+1})d_1\cdots d_{k+1}\tau_{k+1}(n)}{d_1\cdots d_{k+1}n}$$

$$= \frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)$$
$$\times \sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{\substack{1\le n\le N^{k+1}\\(n,ud)=1\\ d_1\cdots d_{k+1}n=ls+1}}\sum_{0\le l\le((Nv/d)^{k+1}-1)/s}\frac{\mu(d_1)\cdots\mu(d_{k+1})d_1\cdots d_{k+1}\tau_{k+1}(n)}{ls+1}$$
$$-\frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)$$
$$\times \sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{\substack{1\le n\le N^{k+1}\\(n,ud)=1\\ d_1\cdots d_{k+1}n=ls-1}}\sum_{2/s\le l\le((Nv/d)^{k+1}+1)/s}\frac{\mu(d_1)\cdots\mu(d_{k+1})d_1\cdots d_{k+1}\tau_{k+1}(n)}{ls-1}$$
$$= \frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)$$
$$+ O\left(\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\frac{\phi(s)}{s}\sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{1\le l\le((Nv/d)^{k+1}-1)/s}\frac{d_1\cdots d_{k+1}N^{\epsilon}}{l+1/s}\right)$$
$$+ O\left(\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\frac{\phi(s)}{s}\sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{2/s\le l\le((Nv/d)^{k+1}+1)/s}\frac{d_1\cdots d_{k+1}N^{\epsilon}}{l-1/s}\right)$$
$$= \frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}J(ud)+O(q^{k+1+\epsilon}N^{\epsilon}) = \frac{u^k\phi^2(u)}{2}\sum_{d|v} d^{k+1}J(d)+O(q^{k+1+\epsilon}N^{\epsilon})$$
$$= \frac{u^k\phi^2(u)}{2}\prod_{p|v}\left[p^k(p-1)^2\left(1-\frac{p^k-1}{p^k(p-1)^2}\right)\right]+O(q^{k+1+\epsilon}N^{\epsilon})$$
$$= \frac{q^k\phi^2(q)}{2}\prod_{p\|q}\left(1-\frac{p^k-1}{p^k(p-1)^2}\right)+O(q^{k+1+\epsilon}N^{\epsilon}),$$

where we have used the estimate $\tau_{k+1}(n)\ll n^{\epsilon}$, the fact that v is a square-free number, u is a square-full number, and the identity $J(u)=\phi^2(u)/u$, if u is a square-full number.

Now taking $N=q^{3/2}$ in the above, we immediately get

$$\Psi_1 = \frac{q^k\phi^2(q)}{2}\prod_{p\|q}\left(1-\frac{p^k-1}{p^k(p-1)^2}\right)+O(q^{k+1+\epsilon}).$$

This completes the proof of Lemma 8. □

Lemma 9. *Let $q = uv$, where $(u, v) = 1$, u be a square-full number or $u = 1$, v be a square-free number. Then we have the following estimates*

$$\begin{aligned}\Psi_4 &:= \sum_{d|v} u^{k+1}d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \mu(d_1)\cdots\mu(d_{k+1}) \\ &\quad \times \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}^j(2)\overline{\chi}(d_1\cdots d_{k+1})L^{k+1}(1, \overline{\chi}) \\ &\ll q^{k+1+\epsilon}, \quad j = 1, 2, \ldots, k+1; \\ \Psi_5 &:= \sum_{d|v} u^{k+1}d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \frac{\mu(d_1)\cdots\mu(d_{k+1})}{d_1\cdots d_{k+1}} \\ &\quad \times \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=1}} \overline{\chi}^j(2)\overline{\chi}(d_1\cdots d_{k+1})L^{k+1}(2, \overline{\chi}) \\ &\ll q^{k+1+\epsilon}, \quad j = 1, 2, \ldots, k+1; \\ \Psi_6 &:= \sum_{d|v} u^{k+1}d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \frac{\mu(d_1)\cdots\mu(d_{k+1})}{d_{k+1}^2} \\ &\quad \times \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}^j(2)\overline{\chi}(d_1\cdots d_{k+1})L^k(1, \overline{\chi})L(3, \overline{\chi}) \\ &\ll q^{k+1+\epsilon}, \quad j = 1, 2, \ldots, k+1.\end{aligned}$$

Proof. We only prove the first formula, since similarly we can get the others. For parameter $N \geq ud$, using the method of Lemma 8 we have

$$\begin{aligned}\Psi_4 &= \sum_{d|v} u^{k+1}d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \sum_{1\leq n\leq N^{k+1}} \frac{\mu(d_1)\cdots\mu(d_{k+1})2^j d_1\cdots d_{k+1}\tau_{k+1}(n)}{2^j d_1\cdots d_{k+1}n} \\ &\quad \times \sideset{}{^*}\sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}} \overline{\chi}(2^j)\overline{\chi}(d_1\cdots d_{k+1})\overline{\chi}(n) + O\left(\frac{q^{k+5/2+\epsilon}\log^k N}{N}\right) \\ &:= \Upsilon + O\left(\frac{q^{k+5/2+\epsilon}\log^k N}{N}\right).\end{aligned}$$

Then from (9) we get

$$\begin{aligned}
\Upsilon &= \frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)\\
&\quad\times \sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{\substack{1\le n\le N^{k+1}\\(n,ud)=1\\ 2^j d_1\cdots d_{k+1}n\equiv 1 \bmod s}}\frac{\mu(d_1)\cdots\mu(d_{k+1})2^j d_1\cdots d_{k+1}\tau_{k+1}(n)}{2^j d_1\cdots d_{k+1}n}\\
&\quad-\frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)\\
&\qquad\times \sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{\substack{1\le n\le N^{k+1}\\(n,ud)=1\\ 2^j d_1\cdots d_{k+1}n\equiv -1 \bmod s}}\frac{\mu(d_1)\cdots\mu(d_{k+1})2^j d_1\cdots d_{k+1}\tau_{k+1}(n)}{2^j d_1\cdots d_{k+1}n}\\
&= \frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)\\
&\quad\times \sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{\substack{1\le n\le N^{k+1}\\(n,ud)=1\\ 2^j d_1\cdots d_{k+1}n=ls+1}}\sum_{(2^j-1)/s\le l\le(2^j(Nv/d)^{k+1}-1)/s}\frac{\mu(d_1)\cdots\mu(d_{k+1})2^j d_1\cdots d_{k+1}\tau_{k+1}(n)}{ls+1}\\
&\quad-\frac{1}{2}\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\mu\left(\frac{ud}{s}\right)\phi(s)\\
&\qquad\times \sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{\substack{1\le n\le N^{k+1}\\(n,ud)=1\\ 2^j d_1\cdots d_{k+1}n=ls-1}}\sum_{(2^j+1)/s\le l\le(2^j(Nv/d)^{k+1}+1)/s}\frac{\mu(d_1)\cdots\mu(d_{k+1})2^j d_1\cdots d_{k+1}\tau_{k+1}(n)}{ls-1}\\
&\ll \sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\frac{\phi(s)}{s}\sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{(2^j-1)/s\le l\le(2^j(Nv/d)^{k+1}-1)/s}\frac{d_1\cdots d_{k+1}N^{\epsilon}}{l+1/s}\\
&\quad+\sum_{d|v} u^{k+1}d^{k+1}\sum_{s|ud}\frac{\phi(s)}{s}\sum_{d_1|(v/d)}\cdots\sum_{d_{k+1}|(v/d)}\sum_{(2^j+1)/s\le l\le(2^j(Nv/d)^{k+1}+1)/s}\frac{d_1\cdots d_{k+1}N^{\epsilon}}{l-1/s}\\
&\ll q^{k+1+\epsilon}N^{\epsilon}.
\end{aligned}$$

Now taking $N = q^{3/2}$ in the above, we immediately get

$$\Psi_4 \ll q^{k+1+\epsilon}.$$

This proves Lemma 9. □

3. Proof of the theorem

In this section, we complete the proof of the theorem. Let $q \geq 3$ be an odd number and $k \geq 1$ be an integer, for any character $\chi \bmod q$, we have

(10)

$$\sum_{c=1}^{q} \overline{\chi}(c) K\left(\overline{2}^{k+1} c, k, q\right) = \sum_{a_1=1}^{q}{}' \cdots \sum_{a_k=1}^{q}{}' \sum_{c=1}^{q} \overline{\chi}(c) \mathrm{e}\left(\frac{a_1 + \cdots + a_k + \overline{2}^{k+1} c \cdot \overline{a}_1 \cdots \overline{a}_k}{q}\right)$$

$$= \overline{\chi}^{k+1}(2)\tau(\overline{\chi})\left(\sum_{a=1}^{q} \overline{\chi}(a)\mathrm{e}\left(\frac{a}{q}\right)\right)^k = \overline{\chi}^{k+1}(2)\tau^{k+1}(\overline{\chi}).$$

We first treat the case $k = 1$ of the theorem. From Lemma 1 we get

$$\sum_{c=1}^{q}{}' E(q, 1, c) K(\overline{4}c, 1, q)$$

$$= \frac{1}{\phi(q)} \sum_{\substack{\chi(-1)=1 \\ \chi \neq \chi_0}} \overline{\chi}(4)\tau^2(\overline{\chi})\left(\sum_{a=1}^{q}(-1)^a a\chi(a)\right)^2 + \frac{1}{\phi(q)} \sum_{\chi(-1)=-1} (4 - 4\overline{\chi}(2))\tau^2(\overline{\chi})\left(\sum_{a=1}^{q} a\chi(a)\right)^2$$

$$- \frac{2}{\phi(q)q} \sum_{\chi(-1)=-1} (\overline{\chi}(4) - 2\overline{\chi}(2))\tau^2(\overline{\chi})\left(\sum_{a=1}^{q} a\chi(a)\right)\left(\sum_{b=1}^{q}(-1)^b b^2 \chi(b)\right) + O(q^{5/2+\epsilon}).$$

Let $q = uv$, where $(u, v) = 1$, u be a square-full number or $u = 1$, v be a square-free number. Suppose that χ is a character modulo q, generated by the primitive character χ_m modulo m. Note that $\tau(\overline{\chi}) = \overline{\chi}_m(q/m)\mu(q/m)\tau(\overline{\chi}_m) \neq 0$ if and only if $m = ud$, where $d|v$. Then from Lemmas 4, 5, 6 we have

$$\sum_{c=1}^{q}{}' E(q, 1, c) K(\overline{4}c, 1, q)$$

$$= \frac{1}{\pi^4 \phi(q)} \sum_{d|v} u^4 d^4 \sum_{\substack{\chi \bmod ud \\ \chi(-1)=1}}{}^{*} [16 - 8\overline{\chi}(2) + \overline{\chi}(4)]$$

$$\times \overline{\chi}^2\left(\frac{v}{d}\right)\mu^2\left(\frac{v}{d}\right)\left[\sum_{d_1|(v/d)} \frac{v}{dd_1}\mu\left(\frac{v}{dd_1}\right)\chi\left(\frac{v}{dd_1}\right)\right]^2 L^2(2, \overline{\chi})$$

$$+ \frac{q^2}{\pi^2 \phi(q)} \sum_{d|v} u^2 d^2 \sum_{\substack{\chi \bmod ud \\ \chi(-1)=-1}}{}^{*} [4 - 4\overline{\chi}(2) + 2\overline{\chi}(4)]\overline{\chi}^2\left(\frac{v}{d}\right)\mu^2\left(\frac{v}{d}\right)$$

$$\times \left[\sum_{d_1|(v/d)} \mu\left(\frac{v}{dd_1}\right)\chi\left(\frac{v}{dd_1}\right)\right]^2 L^2(1, \overline{\chi})$$

$$+\frac{2}{\pi^4\phi(q)}\sum_{d|v}u^4d^4\sideset{}{^*}\sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}[-16+10\overline{\chi}(2)-\overline{\chi}(4)]\overline{\chi}^2\left(\frac{v}{d}\right)\mu^2\left(\frac{v}{d}\right)$$
$$\times\left[\sum_{d_1|(v/d)}\mu\left(\frac{v}{dd_1}\right)\chi\left(\frac{v}{dd_1}\right)\right]\left[\sum_{d_2|(v/d)}\frac{v^2}{d^2d_2^2}\mu\left(\frac{v}{dd_2}\right)\chi\left(\frac{v}{dd_2}\right)\right]L(1,\overline{\chi})L(3,\overline{\chi})$$
$$+O(q^{7/2+\epsilon}).$$

Note that

$$\overline{\chi}\left(\frac{v}{d}\right)=\overline{\chi}\left(\frac{v}{dd_1}\right)\overline{\chi}(d_1),\quad \mu\left(\frac{v}{d}\right)=\mu\left(\frac{v}{dd_1}\right)\mu(d_1), \tag{11}$$

so by Lemma 8 and Lemma 9 we have

$$\begin{aligned}
&\sum_{c=1}^{q}{}' E(q,1,c)K(\overline{4}c,1,q)\\
&=\frac{q^2}{\pi^4\phi(q)}\sum_{d|v}u^2d^2\sum_{d_1|(v/d)}\sum_{d_2|(v/d)}\frac{\mu(d_1)\mu(d_2)}{d_1d_2}\\
&\quad\times\sideset{}{^*}\sum_{\substack{\chi \bmod ud\\ \chi(-1)=1}}[16-8\overline{\chi}(2)+\overline{\chi}(4)]\overline{\chi}(d_1d_2)L^2(2,\overline{\chi})\\
&\quad+\frac{q^2}{\pi^2\phi(q)}\sum_{d|v}u^2d^2\sum_{d_1|(v/d)}\sum_{d_2|(v/d)}\mu(d_1)\mu(d_2)\\
&\qquad\times\sideset{}{^*}\sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}[4-4\overline{\chi}(2)+2\overline{\chi}(4)]\overline{\chi}(d_1d_2)L^2(1,\overline{\chi})\\
&\quad+\frac{2q^2}{\pi^4\phi(q)}\sum_{d|v}u^2d^2\sum_{d_1|(v/d)}\sum_{d_2|(v/d)}\frac{\mu(d_1)\mu(d_2)}{d_2^2}\\
&\qquad\times\sideset{}{^*}\sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}[-16+10\overline{\chi}(2)-\overline{\chi}(4)]\overline{\chi}(d_1d_2)L(1,\overline{\chi})L(3,\overline{\chi})+O(q^{7/2+\epsilon})\\
&=-\frac{6q^3\phi(q)}{\pi^4}\prod_{p\|q}\left(1-\frac{p-1}{p(p-1)^2}\right)+O(q^{7/2+\epsilon}).
\end{aligned}$$

This proves the theorem with $k=1$.

Now let $k \geq 2$ be an integer. Then from formulae (10), (11), and Lemmas 2, 4, 5, 6, 8, 9 we can have

$$\begin{aligned}
&\sum_{c=1}^{q}{}' E(q,k,c)K\left(\overline{2}^{k+1}c,k,q\right)\\
&= \frac{-2^{k-2}}{\phi(q)q^{k-1}} \sum_{\chi(-1)=-1} \tau^{k+1}(\overline{\chi})\overline{\chi}^{k+1}(2)(1-2\chi(2))^k\\
&\quad \times \left(\sum_{a=1}^{q} a\chi(a)\right)^k \left[\frac{2(k+1)}{q}\left(\sum_{b=1}^{q}(-1)^b b^2\chi(b)\right) + k(k-3)(1-2\chi(2))\left(\sum_{b=1}^{q} b\chi(b)\right)\right]\\
&= \frac{i^{k+1}(-1)^{k+2}2^{k-2}(k^2-k+2)q^2}{\pi^{k+1}\phi(q)} \sum_{d|v} u^{k+1}d^{k+1} \sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}{}^{*} \overline{\chi}^{k+1}\left(\frac{v}{d}\right)\mu^{k+1}\left(\frac{v}{d}\right)\\
&\quad \times [\overline{\chi}(2)-2]^{k+1}\left[\sum_{d_1|(v/d)} \mu\left(\frac{v}{dd_1}\right)\chi\left(\frac{v}{dd_1}\right)\right]^{k+1} L^{k+1}(1,\overline{\chi})\\
&\quad + \frac{i^{k+1}(-1)^{k+2}2^{k-1}(k+1)}{\pi^{k+3}\phi(q)} \sum_{d|v} u^{k+3}d^{k+3} \sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}{}^{*} \overline{\chi}^{k+1}\left(\frac{v}{d}\right)\mu^{k+1}\left(\frac{v}{d}\right)[\overline{\chi}(2)-2]^k\\
&\qquad \times [8-\overline{\chi}(2)]\left[\sum_{d_1|(v/d)} \mu\left(\frac{v}{dd_1}\right)\chi\left(\frac{v}{dd_1}\right)\right]^k \left[\sum_{d_2|(v/d)} \frac{v^2}{d^2d_2^2}\mu\left(\frac{v}{dd_2}\right)\chi\left(\frac{v}{dd_2}\right)\right]\\
&\qquad \times L^k(1,\overline{\chi})L(3,\overline{\chi}) + O(q^{k+5/2+\epsilon})\\
&= \frac{i^{k+1}(-1)^{k+2}2^{k-2}(k^2-k+2)q^2}{\pi^{k+1}\phi(q)} \sum_{d|v} u^{k+1}d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \mu(d_1)\cdots\mu(d_{k+1})\\
&\quad \times \sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}{}^{*} [\overline{\chi}(2)-2]^{k+1}\overline{\chi}(d_1\cdots d_{k+1})L^{k+1}(1,\overline{\chi})\\
&\quad + \frac{i^{k+1}(-1)^{k+2}2^{k-1}(k+1)q^2}{\pi^{k+3}\phi(q)} \sum_{d|v} u^{k+1}d^{k+1} \sum_{d_1|(v/d)} \cdots \sum_{d_{k+1}|(v/d)} \frac{\mu(d_1)\cdots\mu(d_{k+1})}{d_{k+1}^2}\\
&\qquad \times \sum_{\substack{\chi \bmod ud\\ \chi(-1)=-1}}{}^{*} [\overline{\chi}(2)-2]^k[8-\overline{\chi}(2)]\overline{\chi}(d_1\cdots d_{k+1})L^k(1,\overline{\chi})L(3,\overline{\chi}) + O(q^{k+5/2+\epsilon})\\
&= \frac{i^{k+1}(-1)2^{2k-2}(k^2-k+2)q^{k+2}\phi(q)}{\pi^{k+1}} \prod_{p\|q}\left(1-\frac{p^k-1}{p^k(p-1)^2}\right)\\
&\quad + \frac{i^{k+1}2^{2k+1}(k+1)q^{k+2}\phi(q)}{\pi^{k+3}} \prod_{p\|q}\left(1-\frac{p^k-1}{p^k(p-1)^2}\right) + O(q^{k+5/2+\epsilon})
\end{aligned}$$

$$= \frac{i^{k+3}2^{2k-2}q^{k+2}\phi(q)}{\pi^{k+3}}[\pi^2(k^2-k+2)-8(k+1)]\prod_{p\|q}\left(1-\frac{p^k-1}{p^k(p-1)^2}\right)+O(q^{k+5/2+\epsilon}).$$

So from the above we have

$$\sum_{c=1}^{q}{}' E(q,k,c)K\left(\overline{2}^{k+1}c,k,q\right)$$
$$= \frac{c_k q^{k+2}\phi(q)}{\pi^{k+3}}\prod_{p\|q}\left(1-\frac{p^k-1}{p^k(p-1)^2}\right)+O(q^{k+5/2+\epsilon}), \quad \text{for} \quad k\geq 1.$$

This completes the proof of the theorem.

ACKNOWLEDGMENTS. The authors express their gratitude to the referee for his very helpful and detailed comments in improving this paper.

References

[1] W. Zhang: *On the difference between an integer and its inverse modulo n*, J. Number Theory **52** (1995), 1–6.
[2] W. Zhang: *On the difference between an integer and its inverse modulo n* (II), Sci. China Ser. A **46** (2003), 229–238.
[3] R.K. Guy Unsolved Problems in Number Theory, Second edition, Springer, New York, 1994.
[4] W. Zhang: *A problem of D.H. Lehmer and its Generalization* (I), Compositio Math. **86** (1993), 307–316.
[5] W. Zhang: *A problem of D.H. Lehmer and its Generalization* (II), Compositio Math. **91** (1994), 47–56.
[6] W. Zhang: *On the difference between a D.H. Lehmer number and its inverse modulo q*, Acta Arith. **68** (1994), 255–263.
[7] W. Zhang: *A problem of D.H. Lehmer and its mean square value formula*, Japan. J. Math. (N.S.) **29** (2003), 109–116.
[8] W. Zhang, X. Zongben and Y. Yuan: *A problem of D.H. Lehmer and its mean square value formula*, J. Number Theory **103** (2003), 197–213.
[9] W. Zhang: *On a problem of D.H. Lehmer and Kloosterman sums*, Monatsh. Math. **139** (2003), 247–257.
[10] L.J. Mordell: *On a special polynomial congruence and exponential sums*; in Calcutta Math. Soc. Golden Jubilee Commemoration Vol., Part 1, Calcutta Math. Soc., Calcutta., 1963, 29–32.
[11] D. Bump, W. Duke, J. Hoffstein and H. Iwaniec: *An estimate for the Hecke eigenvalues of Maass forms*, Internat. Math. Res. Notices (1992), 75–81.
[12] W. Luo, Z. Rudnick and P. Sarnak: *On Selberg's eigenvalue conjecture*, Geom. Funct. Anal. **5** (1995), 387–401.
[13] R.A. Smith: *On n-dimensional Kloosterman sums*, J. Number Theory **11** (1979), 324–343.
[14] T.M. Apostol: Introduction to Analytic Number Theory, Springer, New York, 1976.
[15] T. Funakura: *On Kronecker's limit formula for Dirichlet series with periodic coefficients*, Acta Arith. **55** (1990), 59–73.
[16] G. Pólya: *Über die Verteilung der quadratishen Reste und Nichtreste*, Gottinger Nachrichten (1918), 21–29.

[17] C. Pan and C. Pan: Goldbach Conjecture, Science Press, Beijing, 1992.
[18] W. Zhang: *On a Cochrane sum and its hybrid mean value formula*, J. Math. Anal. Appl. **267** (2002), 89–96.
[19] W. Zhang, Y. Yi and X. He: *On the* $2k$*-th power mean of Dirichlet* L*-functions with the weight of general Kloosterman sums*, J. Number Theory **84** (2000), 199–213.

Huaning Liu
Department of Mathematics
Northwest University
Xi'an, Shaanxi
P.R. China
e-mail: hnliumath@hotmail.com

Wenpeng Zhang
Department of Mathematics
Northwest University
Xi'an, Shaanxi
P.R. China
e-mail: wpzhang@nwu.edu.cn