



Title	Supplementary remarks on categories of indecomposable modules
Author(s)	Harada, Manabu
Citation	Osaka Journal of Mathematics. 1972, 9(1), p. 49-55
Version Type	VoR
URL	https://doi.org/10.18910/9628
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

SUPPLEMENTARY REMARKS ON CATEGORIES OF INDECOMPOSABLE MODULES

MANABU HARADA

(Received May 11, 1971)

In the previous papers [3], [4], we have defined a full sub-category \mathfrak{A} in the category \mathfrak{M}_R of modules over a ring R , whose objects consist of injective modules or directsums of completely indecomposable modules.

Making use of those ideas, in this short note, we shall give a proof of Z. Papp's theorem in [9] as an application of [3], Theorem 1 and generalize Theorems 4 and 7 in [4] to cases of semi- T -nilpotent system and quasi-projective module, respectively. Especially, we shall show that if R is a right perfect ring, then every quasi-projective module is a directsum of completely indecomposable modules and the Krull-Remak-Schmidt's theorem is valid for those direct decompositions.

In this note, we always assume that the ring R has the identity and every module is an unitary R -module. We shall use the same notations and definitions in [3], [4] and [5] for categories, those in [1] and [8] for semi-perfect modules and those in [5] for quasi-projective modules.

1. Papp's theorem

We shall give an application of [3], Theorem 1.

Theorem 1 ([9], Z. Papp). *Let R be a ring. If every (right) R -injective module is a directsum of indecomposable modules, then R is (right) noetherian.*

Proof. It is known by [2], Proposition 4.1 that R is noetherian if and only if any directsum of injective modules is also injective. Let \mathfrak{A} be the full sub-category of all injective R -modules in the category of right R -modules and \mathfrak{J} the Jacobson radical of \mathfrak{A} . Then $\mathfrak{A}/\mathfrak{J}$ is a completely reducible C_3 -abelian category by the assumption and [3], Theorem 1. Let $\{Q_i\}_1^\infty$ be a family of injective modules, and E an injective hull of $\sum \oplus Q_i$. From the assumption $E = \sum \oplus E_i$, where E_i 's are (completely) indecomposable. Hence, $\sum \oplus E_j = \sum \oplus Q_i$ in $\mathfrak{A}/\mathfrak{J}$ by [3], Theorem 1. Therefore, $E \approx \sum \oplus Q_i$, which means that $\sum \oplus Q_i$ is injective. Hence, R is noetherian.

2. Exchange property

Let M be a directsum of completely indecomposable modules M_α ; $M = \sum \oplus M_\alpha$. We have defined the (\aleph_0-) exchange property in M for a direct summand N of M in [4]. Namely, we have $M = N \oplus \sum_I \oplus T'_\alpha$ for any decomposition $M = \sum_I \oplus T_\alpha$ (with $\text{Card } I \leq \aleph_0$), where $T'_\alpha \subseteq T_\alpha$ for all $\alpha \in I$.

Let $M = N \oplus N'$. If N has the exchange property in M , then N and N' are directsums of indecomposable modules.

Now we assume $M = \sum_I \oplus M_\alpha$. A family $\{M_\beta\}_J$ ($J \subseteq I$) is called a *semi- T -nilpotent system* with respect to the radical of $[M_\beta, M_\beta]_R$ if the following condition is satisfied. J is a finite or empty set or if J is otherwise, for any subfamily $\{M_{\beta_i}\}$ with $\beta_i \in J$ and $\beta_i \neq \beta_j$ if $i \neq j$ and any set of non isomorphisms $f_i: M_{\beta_i} \rightarrow M_{\beta_{i+1}}$, there exists a natural number n such that $f_n f_{n-1} \dots f_1(m) = 0$ for $m \in M_{\beta_1}$, where n may depend on m , (cf. [5]). Then we have a generalization of [4], Theorem 4 as follows;

Theorem 2. *Let $M = \sum_I \oplus M_\alpha$ with M_α completely indecomposable and $M = N_1 \oplus N_2$. If the dense submodule of $N_1^{(1)}$ is a directsum of indecomposable modules which are a semi- T -nilpotent system with respect to the radical, then N_i has the exchange property in M for $i=1, 2$.*

Proof. We first note that $N_1 = \sum \oplus M'_\beta$ by the assumption and [7], Corollary to Theorem 1. Furthermore, since the ideal \mathfrak{J} of $S_{N_1} = [N_1, N_1]_R$ defined in [3], §3 is equal to the Jacobson radical of S_{N_1} by [7], Theorem 1. Hence, we have from the first part of the proof of [4], Theorem 4 that N_2 has the exchange property. Let $M = \sum_K \oplus T_\beta$ with any $\text{Card } K$. We shall use the same notation in [4]. If we consider the category $\mathfrak{A}/\mathfrak{J}$ in [4], then $\bar{M} = \sum \oplus \bar{T}_\beta = \bar{N}_1 \oplus \bar{N}_2$ in $\mathfrak{A}/\mathfrak{J}$ by [4], Theorem 1. Since $\mathfrak{A}/\mathfrak{J}$ is a completely reducible C_3 -abelian category, $\bar{M} = \bar{N}_1 \oplus (\sum \oplus \bar{T}'_\beta)$, where $\bar{T}_\beta = \bar{T}'_\beta \oplus \bar{T}''_\beta$ and we may assume that T'_β and T''_β are in \mathfrak{A} and submodules in T_β by [4], Proposition 2. Since $\sum_K \oplus \bar{T}''_\beta \approx \bar{N}_1$, $N_1 \approx \sum \oplus T''_\beta$ by the assumption and [7], Theorem 1. Let p be a homomorphism of M to $\sum_K \oplus T''_\beta$ such that \bar{p} is a projection of \bar{M} to $\sum_K \oplus \bar{T}''_\beta$ with $\text{Ker } \bar{p} = \sum_K \oplus \bar{T}'_\beta$. Then p splits. Put $L = \text{Ker } p$, then $T_\beta = T''_\beta \oplus T^*_\beta$, where $T^*_\beta = T_\beta \cap L$. Since $L = \sum \oplus T^*_\beta$ and $\bar{L} = \text{Ker } \bar{p}$, $\sum_K \oplus \bar{T}^*_\beta = \sum_K \oplus \bar{T}'_\beta$. Hence, $\bar{M} = \bar{N}_1 \oplus (\sum \oplus \bar{T}'_\beta) = \sum_K \oplus \bar{T}''_\beta \oplus \sum_K \oplus \bar{T}'_\beta$. Therefore, $\text{Ker } \bar{p} \cap \bar{N}_1 = \bar{0}$ and $\bar{p}(\bar{N}_1) = \bar{p}(\bar{M}) = \sum_K \oplus \bar{T}''_\beta$, which implies $p|_{N_1}$ is isomorphic. Hence, $M = N_1 \oplus \text{Ker } p = N_1 \oplus$

1) See [4], §1 for the definition.

$$\sum_K \oplus T_\beta^*.$$

Corollary. *Let M be as above and N a direct summand of M . Then the following statements are equivalent.*

- 1) *Every direct summand of N has the \aleph_0 -exchange property in M .*
- 2) *Every direct summand of N has the exchange property in M .*
- 3) *$N = \sum_j \oplus N_\beta$; $N_\beta \approx M_{\pi(\beta)}$ and $\{N_\beta\}$ is a semi- T -nilpotent system with*

respect to the radical of $[N_\gamma, N_\delta]_R$.

Proof. 1) \rightarrow 3). Let N be a direct summand of M and $N = \sum_j \oplus N'_\beta$, $N'_\beta \approx M_\beta$ with $\text{Card } J \leq \text{Card } I$. We first note that every direct summand P of N has the \aleph_0 -exchange property in N . Let $M = N \oplus Q$ and $N = P_1 \oplus P_2 = \sum_{i=1}^\infty \oplus T_i$. Then $M = P_1 \oplus (P_2 \oplus Q) = \sum \oplus T_i \oplus Q$. Since P_1 has the \aleph_0 -exchange property in M , $M = P_1 \oplus \sum \oplus T'_i \oplus Q'$, where $T'_i \subseteq T_i$ and $Q' \subseteq Q$. Hence, $N = P_1 \oplus \sum \oplus T'_i \oplus P_1 \cap Q'$ and $P_1 \cap Q' \subseteq P_1 \cap Q = (0)$. Now put $P_1 = \sum_{J_0} \oplus N_\gamma$ for any $J_0 \subseteq I$ with $\text{Card } J_0 \leq \aleph_0$. Then $\{N_\gamma\}_{J_0}$ is a semi- T -nilpotent system by [7], Theorem 1. Hence, $\{N_\gamma\}_J$ is a semi- T -nilpotent system. 3) \rightarrow 2). Since the ideal \mathfrak{J} of $[N, N]_R$ defined in [3] is the Jacobson radical by [7], Theorem 1, every direct summand of N is a directsum of indecomposable modules and has the exchange property in M by Theorem 2. 2) \rightarrow 1). It is clear.

Lemma 1. *Let M be as above. We assume that $M = N_1 \oplus N_2 = N'_1 \oplus N'_2$. If N_1 has the exchange property in M and there exists an automorphism f of M such that $f(N_i) = N'_i$ for $i = 1, 2$ then N'_1 has the exchange property in M .*

Proof. It is clear.

Lemma 2. *Let M, N_1 and N_2 be as above. We assume $N_i = \sum_{\alpha \in J_i} \oplus M_{i\alpha}$, $\text{Card } J_i$ are infinite and $M_{i\alpha}$'s are indecomposable modules for $i = 1, 2$. Let $\{f_i\}_1^\infty$, $\{g_i\}_1^\infty$, be sets of non-isomorphic homomorphisms of $M_{1\alpha_i}$ to $M_{2\alpha_i}$ and $M_{2\alpha_i}$ to $M_{1\alpha_{i+1}}$, respectively. Furthermore, we assume that N_1 has the \aleph_0 -exchange property. Then for any m in $M_{1\alpha_1}$ there exists n such that $g_n f_n g_{n-1} f_{n-1} \cdots g_1 f_1(m) = 0$.*

Proof. We shall make use of the same argument in [3], Lemma 9. Put $M'_{1\alpha_i} = \{m_i + f_i(m_i) \mid m_i \in M_{1\alpha_i}\}$ and $M'_{2\alpha_i} = \{m_i + g_i(m_i) \mid m_i \in M_{2\alpha_i}\}$. Then $M = M'_{1\alpha_1} \oplus M_{2\alpha_1} \oplus M'_{1\alpha_2} \oplus M_{2\alpha_2} \oplus \cdots \oplus M_{10} \oplus M_{20} = M_{1\alpha_1} \oplus M'_{2\alpha_1} \oplus M_{1\alpha_1} \oplus M'_{2\alpha_2} \oplus \cdots \oplus M_{10} \oplus M_{20}$, where $M_{i0} = \sum_{J_i - \{1, 2, \dots\}} \oplus M_{i\alpha}$. Since $T = M'_{1\alpha_1} \oplus M'_{1\alpha_2} \oplus \cdots \oplus M_{10} \approx N_1$, T has the \aleph_0 -exchange property in M by Lemma 1. Hence, $M = T \oplus M_{1\alpha_1}^* \oplus M_{2\alpha_1}^* \oplus M_{1\alpha_2}^* \oplus M_{2\alpha_2}^* \oplus \cdots \oplus M_{20}^*$, where $M_{1\alpha_j}^* = 0$ or $M_{1\alpha_j} (M_{2\alpha_j}^* = 0$ or $M'_{2\alpha_j})$. In this case we can use the same argument in [3], Lemma 9.

From Lemma 2 we have

Proposition 1. *Let $M = \sum \oplus M_\alpha$ with M_α completely indecomposable. We assume that $M = N_1 \oplus N_2$ and $N_i = \sum_\gamma \sum_{\beta \in J_\gamma} \oplus M_{\gamma\beta}^{(\iota)}$, where $M_{\gamma\beta}^{(\iota)} \approx M_{\gamma'\beta'}^{(\iota)}$ and $M_{\gamma\beta}^{(\iota)} \not\approx M_{\gamma'\beta'}^{(\iota)}$ if $\gamma \neq \gamma'$, where $M_{\alpha\beta}^{(\iota)}$'s are indecomposable. We further assume $\text{Card } J_\gamma^{(2)} \geq \text{Card } J_\gamma^{(1)}$ for all $\text{Card } J_\gamma^{(2)}$ which is smaller than or equal to \aleph_0 . Then N_1 has the (\aleph_0-) exchange property if and only if $\{M_{\gamma\beta}^{(1)}\}$ is a semi- T -nilpotent system with respect to the radical.*

Now we take the category \mathfrak{A} of all R -modules which is a directsum of some completely indecomposable modules. Let M be an object in \mathfrak{A} . We call M having the exchange property in \mathfrak{A} if M has the exchange property in P for any object P in \mathfrak{A} which contains M as a direct summand.

Corollary 2. *Let \mathfrak{A} be the above. Then we have the following equivalent statements for $M = \sum_I \oplus M_\alpha$ in \mathfrak{A} .*

- 1) M has the exchange property in \mathfrak{A} .
- 2) $\{M_\alpha\}_I$ is a semi- T -nilpotent system with respect to the radical, where M_α 's are completely indecomposable.

Proof. 2) \rightarrow 1). It is clear from Corollary to Theorem 2. 1) \rightarrow 2). Let $M = \sum_\alpha \sum_{I_\alpha \in \beta} \oplus M_{\alpha\beta}$, $M_{\alpha\beta} \approx M_{\alpha'\beta'}$ and $M_{\alpha\beta} \not\approx M_{\alpha'\beta'}$ if $\alpha \neq \alpha'$. Put $P = \sum_1^\infty \oplus P_n$, $P_n = M$. Since M has the exchange property in P by the assumption, $\{M_\alpha\}$ is a semi- T -nilpotent system by Proposition 1.

Finally, we shall consider a special case. Let Z be the ring of integers (or Z may be a Dedekind domain) and $\{P_i\}_I$ a family of primes. Let M be a directsum of any copies of $Z/P_i^{n_i}$, where i runs over a sub-set of I and n_i 's are integers. Then $M = \sum_{i \in I} \oplus M_{P_i}$, where $M_{P_i} = \sum \oplus Z/P_i^{n_i}$. In this case, every submodule N of M is a directsum of N_P , where $N_P = N \cap M_P$. Hence, a direct summand N of M has the exchange property in M if and only if N_P has the exchange property in M_P for each P .

Corollary 3. *Let Z , M_P and M be as above. We assume $M = N_1 \oplus N_2$ and $N_i = \sum \oplus M_{P_j}^{(\iota)}$; $M_{P_j}^{(\iota)} \approx Z/P_j^{n_j}$. Then N_1 has the exchange property in M if and only if either $\{M_{P_j}^{(1)}\}$ or $\{M_{P_j}^{(2)}\}_j$ is a semi- T -nilpotent system with respect to the radical for every P_j .*

Proof. It is clear from Lemma 2 and [3], Lemma 12.

REMARK. If $P_1 \neq P_2$, then $\sum_{n=1}^\infty \oplus Z/P_1^n$ has the exchange property in $P = \sum_1^\infty \oplus Z/P_1^n \oplus \sum_1^\infty \oplus Z/P_2^n$ from the above remark. However, $\{Z/P_i^n\}_n$ are not semi- T -nilpotent systems for $i=1, 2$. Hence, M does not have the exchange

property in \mathfrak{A} .

3. Quasi-projective modules

First, we consider projective modules of a special type.

Lemma 3. *Let P and Q be projective R -modules such that $J(P)$ and $J(Q)$ are small in P and Q , respectively. Then $[P/J(P), Q/J(Q)]_{R/J(R)}=0$ if and only if $[Q/J(Q), P/J(P)]_{R/J(R)}=0$, where $J(*)$ is the Jacobson radical of $(*)$.*

Proof. Put $T=P\oplus Q$. Then $J(T)$ is a unique maximal one among small submodules in T . We assume $[P, Q]_R=[P, J(Q)]_R$ and f an element in $[Q, P]_R$. We put $f_T=(\begin{smallmatrix} 0 & 0 \\ 0 & 0 \end{smallmatrix})$ in $S_T=[T, T]_R$. Since $[P, Q]_R f=[P, J(Q)]_R f\subseteq [Q, J(Q)]_R\subseteq J(S_Q)$ by [4], Proposition 1. Hence, $S_T f_T$ is in $J(S_T)$. Therefore, $f_T(T)\subseteq J(P)\oplus J(Q)$. Hence, $f(Q)\subseteq J(P)$ and $[Q, P]_R=[Q, J(P)]_R$. It is clear that $[P, J(Q)]_R=[P, Q]_R$ if and only if $[P/J(P), Q/J(Q)]_R=0$, since P is projective.

Proposition 2. *Let P and Q be as above. We further assume that P is completely indecomposable, then the following are equivalent.*

- 1) P is isomorphic to a direct summand of Q .
- 2) $P/J(P)$ is isomorphic to a sub-module of $Q/J(Q)$.

Proof. It is clear, since $J(P)$ is a unique maximal sub-module in P by [4], Theorem 5.

Changing slightly the proofs in [10], Lemma 1 and [5], Proposition 1, we have

Lemma 4. *Let M be a quasi-projective, then $J(S_M)=\{f \mid f \in S_M, f(M) \text{ is small in } M\}$. Furthermore, $J(M)$ is small if and only if $[M, J(M)]_R=J(S_M)$, where, $S_M=[M, M]_R$.*

We note that a quasi-projective module with projective cover is nothing but a factor module of projective module P with respect to a small R -sub-module K in P which is a S_P -module by [6], Propositions 2.1 and 2.2. Furthermore, if we take the ring of column summable matrices, we know Proposition 2.4 in [6] is valid for a directsum of infinite components, (cf. [3], § 3).

Proposition 3. *Let M be a quasi-projective. We assume that M has projective cover P . Then $S_M \approx S_P/A$ and $P/J(P) \approx M/J(M)$, where A is an ideal contained in $J(S_P)$. Furthermore, $J(P)$ is small in P if and only if $J(M)$ is small in M .*

Proof. We have the exact sequence $0 \rightarrow [P, K]_R \rightarrow S_P \rightarrow [P, M]_R \rightarrow 0$ from an exact sequence $0 \rightarrow K \rightarrow P \xrightarrow{\nu} M \rightarrow 0$. $A=[P, K]_R$ is a two-sided ideal by [6],

Proposition 2.2. Let f be in $[P, M]_R$. Since P is projective, we have g in S_P such that $\nu g = f$. Hence, $f(K) = \nu g(K) \subseteq \nu(K) = 0$. Therefore, $[P, M]_R = S_M$. Since $K \subseteq J(P)$, $J(M) \approx J(P)/K$ and $P/J(P) \approx M/J(M)$. Furthermore, $A \subseteq J(S_P)$ by Lemma 4. The last part is clear.

Lemma 5. Let $\{M_\alpha\}_I$ be a family of quasi-projective modules and I an infinite set. We assume $M = \sum_I \oplus M_\alpha$ is quasi-projective. Then $J(M)$ is small in M if and only if $J(M_\alpha)$ is small in M_α for all $\alpha \in I$ and $\{M_\alpha\}_I$ is a semi- T -nilpotent system with respect to the radical of $[M_\alpha, M_\beta]_R$.

Proof. We can make use of the same argument in [5], Theorem 3 from Lemma 4.

Theorem 3. Let M be a quasi-projective module with projective cover P . Then P is semi-perfect if and only if 1) $M = \sum_I \oplus M_\alpha$; M_α 's are completely indecomposable R -modules, 2) $\{M_\alpha\}_I$ is a semi- T -nilpotent system with respect to the Jacobson radical of $[M_\alpha, M_\beta]_R$ and 3) M_α has a projective cover for all $\alpha \in I$. In this case any direct decomposition of $M/J(M)$ is lifted to M .

Proof. We assume P is semi-perfect. Then 1) is clear from [6], Proposition 2.4 and the above remark. 2) is clear from Proposition 3 and Lemma 4. 3) is clear from [6], Proposition 2.4. Conversely, we assume 1), 2) and 3). Let P_α be a projective cover of M_α via ν_α and $Q = \sum \oplus P_\alpha$. We have an exact sequence $0 \rightarrow K \rightarrow P \xrightarrow{\nu} M \rightarrow 0$ with K small. Hence, we have $f \in [Q, P]_R$ and $g \in [P, Q]_R$ such that $fg = I_P$ and $\nu' = \nu g$, where $\nu = \sum \oplus \nu_\alpha$. Since ν and ν' induce natural isomorphisms $P/J(P) \approx M/J(M) \approx Q/J(Q)$, g is isomorphic. Furthermore, P_α is semi-perfect from Proposition 3 and [4], Theorem 5. We know from 2) and Lemma 4 that $J(M)$ is small in M . Hence, $J(P)$ is small in P by Proposition 3. Therefore, P is semi-perfect by [8], Theorem 5.2. The last part is clear from Proposition 3, [6], Proposition 2.4 and [8], Theorem 4.3.

Corollary. If R is a right perfect (resp, semi-perfect) ring, then every (resp. finitely generated) quasi-projective module is a directsum of completely indecomposable modules and the Krull-Remak-Schmidt's theorem is valid for those decompositions.

OSAKA CITY UNIVERSITY

References

- [1] H. Bass: *Finitistic dimension and a homological generalization of semi-primary rings*, Trans. Amer. Math. Soc. **95** (1960), 466-488.
- [2] S.U. Chase: *Direct products of modules*, Trans. Amer. Math. Soc. **97** (1960), 457-

- 473.
- [3] M. Harada and Y. Sai: *On categories of indecomposable modules I*, Osaka J. Math. **7** (1970), 323–344.
 - [4] M. Harada: *On categories of indecomposable modules II*, Osaka J. Math. **8** (1971), 309–321.
 - [5] M. Harada and H. Kanbara: *On categories of projective modules*, Osaka J. Math. **8** (1971), 471–483.
 - [6] J.P. Jans and L.E.Wu: *On quasi projectives*, Illinois J. Math. **11** (1967), 439–448.
 - [7] H. Kanbara: *On the Krull-Remak-Schmidt-Azumaya's theorem*, to appear.
 - [8] E. Mares: *Semi-perfect modules*, Math. Z. **83** (1963), 347–360.
 - [9] Z. Papp: *On algebraically closed modules*, Publ. Math. Debrecen **6** (1959), 311–323.
 - [10] R. Ware and J. Zelmanowitz: *The Jacobson radical of the endomorphism ring of a projective module*, Proc. Amer. Math. Soc. **26** (1970), 15–20.

