



Title	Cohomotopy of Lie groups
Author(s)	Ōshima, Hideaki; Takahara, Kōji
Citation	Osaka Journal of Mathematics. 1991, 28(2), p. 213-221
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9680">https://doi.org/10.18910/9680</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## COHOMOTOPY OF LIE GROUPS

HIDEAKI ŌSHIMA AND KŌJI TAKAHARA

(Received May 11, 1990)

### 1. Introduction

The purpose of this note is to study the set  $\text{Cdg}(X, n) = \text{Cdg}([X, S^n])$  when  $X = G$  is a compact simply connected simple Lie group, where

$$\text{Cdg}: [X, S^n] \rightarrow \text{Hom}(\pi_n(X), \pi_n(S^n))$$

assigns the induced homotopy homomorphism  $f_*$  to the homotopy class of a map  $f: X \rightarrow S^n$ . To estimate  $\text{Cdg}(X, n)$  we introduce an invariant  $\text{cdg}(X, n)$  and its stable version  ${}^s\text{cdg}(X, n)$ , which are non-negative integers or infinity, such that  ${}^s\text{cdg}(G, 3)$  was denoted by  $\text{cd}(G)$  in [9]. We denote by  $\text{cdg}_p(X, n)$  the exponent of a prime number  $p$  in the prime power decomposition of  $\text{cdg}(X, n)$  when  $0 < \text{cdg}(X, n) < \infty$ . For convenience's sake we set  $\text{cdg}_p(X, n) = 0$  when  $\text{cdg}(X, n) = 0$ . We define  ${}^s\text{cdg}_p(X, n)$  similarly. We prove the following two theorems.

**Theorem 1.** *If  $G$  is a compact simply connected Lie group, that is,  $G = G_1 \times \cdots \times G_r$  with  $G_i$  a compact simply connected simple Lie group, then  $\text{cdg}(G, n)$  and  ${}^s\text{cdg}(G, n)$  are finite and the following seven statements are equivalent for any prime number  $p$ .*

- (1)  $\text{cdg}_p(G, 3) = 0$ .
- (2)  ${}^s\text{cdg}_p(G, 3) = 0$ .
- (3)  $\text{cdg}_p(G_i, 3) = 0$  for all  $i$ .
- (4)  ${}^s\text{cdg}_p(G_i, 3) = 0$  for all  $i$ .
- (5)  $G_i$  is  $p$ -regular for every  $i$ .
- (6)  $G$  is  $p$ -regular.
- (7)  $\text{cdg}_p(G, n) = 0$  for all  $n$ .

**Theorem 2.** *If  $G$  is a compact simply connected simple Lie group, then  $\text{Cdg}(G, n)$  is a subgroup of  $\text{Hom}(\pi_n(G), \pi_n(S^n))$  of maximal rank. Indeed  $\text{Cdg}(G, n)$  is  $\text{cdg}(G, n)\mathbb{Z}\{s'_1\} \oplus c\mathbb{Z}\{s'_2\}$  if  $(G, n) = (\text{Spin}(4m), 4m-1)$  and  $\text{cdg}(G, n) \cdot \text{Hom}(\pi_n(G), \pi_n(S^n))$  otherwise. Here  $\pi_{4m-1}(\text{Spin}(4m)) = \mathbb{Z}\{s_1\} \oplus \mathbb{Z}\{s_2\}$  and  $s'_i$  is the dual element to  $s_i$ ;  $c$  is 1 if  $m \leq 2$  and 2 if  $m \geq 3$ ;  $\text{cdg}(G, n)$  is non-zero if and only if  $n \in \{n_1, \dots, n_r\}$ , where  $H^*(G; \mathbb{Q}) \cong H^*(\prod_{i=1}^r S^{n_i}; \mathbb{Q})$ .*

In this note all spaces are path-connected with base point and all maps

preserve base point. Base point of any H-space is the unit of it. To simplify notation, we denote a map and its homotopy class by the same letter.

We define invariants  $\text{cdg}(X, n)$  and  ${}^s\text{cdg}(X, n)$  in §2, prove Theorems in §3, and give three results without proofs in §4.

### 2. Homotopy invariants

We will use the following notation and convention: We denote by  $a|b$  that  $b=ca$  for some integer  $c$ . For any subset  $A$  of  $\mathbf{Z}$  which contains a non-zero, we denote by  $\text{GCD}(A)$  the greatest common divisor of the non-zero integers in  $A$ . For convenience' sake we set  $\text{GCD}(0)=0$ ,  $k|\infty$  for any non-zero integer  $k$ , and  $0\cdot\infty=0$ , hence  $\infty|0$ . For any subset  $A$  of  $\{k\in\mathbf{Z}; k\geq 0\} \cup \{\infty\}$ , we denote by  $\text{LCM}(A)$  the least common multiple of  $A$  (it may be  $\infty$ ) if  $A$  is non-empty and contains neither 0 nor  $\infty$ , and 0 if  $A$  is empty or contains 0, and  $\infty$  if  $A$  does not contain 0 but  $\infty$ . For any group  $C$ , we denote by  ${}^{ab}C$  the abelianization of  $C$ , that, is  ${}^{ab}C$  is the quotient group of  $C$  by its commutator subgroup. Note that the canonical surjection  $C\rightarrow{}^{ab}C/\text{Tor}$  induces an isomorphism  $\text{Hom}(C, \mathbf{Z})\cong\text{Hom}({}^{ab}C/\text{Tor}, \mathbf{Z})$ , where  $\text{Tor}$  denotes the torsion subgroup. The group  $C$  has the rank  $r$ ,  $\text{rank}C=r$ , if  ${}^{ab}C/\text{Tor}$  is a free abelian group of rank  $r$ . We denote by  $\mathfrak{B}(C)$  the set of  $x\in{}^{ab}C/\text{Tor}$  which is not divisible by any integer  $\geq 2$ .

Put  $\{X, Y\}=\lim_{k\rightarrow\infty}[\Sigma^k X, \Sigma^k Y]$  and  ${}^s\pi_n(X)=\{S^n, X\}$ . Let  ${}^s\text{Cdg}: \{X, S^n\}\rightarrow\text{Hom}({}^s\pi_n(X), {}^s\pi_n(S^n))$  be the stable version of  $\text{Cdg}$ . For any  $\alpha\in\pi_n(X)$ , we denote by  $\text{cdg}(X, n; \alpha)$  or  $\text{cdg}(\alpha)$  the non-negative generator of the subgroup of  $\mathbf{Z}$  generated by the image of  $\alpha^*: [X, S^n]\rightarrow\pi_n(S^n)=\mathbf{Z}$ . We define  ${}^s\text{cdg}(\alpha)$  similarly for any  $\alpha\in{}^s\pi_n(X)$ . If  $\alpha, \beta\in\pi_n(X)$  represent the same element in  ${}^{ab}\pi_n(X)/\text{Tor}$ , then  $\text{cdg}(\alpha)=\text{cdg}(\beta)$ . Thus  $\text{cdg}$  can be defined on  ${}^{ab}\pi_n(X)/\text{Tor}$ . Similarly  ${}^s\text{cdg}$  is defined on  ${}^s\pi_n(X)/\text{Tor}$ .

DEFINITION 2.1.

$$\begin{aligned} \text{cdg}(X, n) &= \text{LCM}\{\text{cdg}(\alpha); \alpha\in\mathfrak{B}(\pi_n(X))\}, \\ {}^s\text{cdg}(X, n) &= \text{LCM}\{{}^s\text{cdg}(\alpha); \alpha\in\mathfrak{B}({}^s\pi_n(X))\}. \end{aligned}$$

The invariant  ${}^s\text{cdg}(X, n)$  has been studied by several people when  $X$  is the Thom space of an  $n$ -dimensional vector bundle [8]. Note from [10] that  $\text{cdg}(Sp(n)/Sp(k), 4n-1)$  and  $\text{cdg}(U(n)/U(k), 2n-1)$  are James numbers [2] for  $0\leq k<n$ , though  $\text{cdg}(O(8)/O(1), 7)=6$  and the James number of  $SO(8)=O(8)/O(1)$  is 1.

The invariant  $\text{cdg}(X, n)$  may be  $\infty$ , though it is finite if  $X$  is a finite CW-complex. Indeed we have

EXAMPLE 2.2. For each prime  $p$ , let  $\alpha_1(3; p)$  and  $\alpha_1(2; p)$  be generators of the  $p$ -components of  $\pi_{2p}(S^3)$  and  $\pi_{2p}(S^2)$ , respectively [12]. Set  $\alpha_1(n; p)=\sum^{n-3}\alpha_1(3; p)\in\pi_{n+2p-3}(S^n)$  and  $X(n; p)=S^n\cup_{\alpha_1(n; p)}e^{n+2p-2}$  for  $n\geq 3$ , and set  $X(2; p)=$

$S^2 \cup_{\sigma_1(2; p)} e^{2p+1}$ . Then  $\text{cdg}(X(n; p), n) = p$  and  $\text{cdg}(\prod_p X(n; p), n) \neq 0$ , hence  $\text{cdg}(\prod_p X(n; p), n) = \infty$  by Proposition 2.6 below.

**Proposition 2.3.** *If  $\pi_n(X)$  is of finite rank, then the following three assertions hold.*

- (1)  $\text{cdg}(X, n) < \infty$ , and  $\text{cdg}(X, n) = \text{cdg}(\alpha)$  for some  $\alpha \in \mathcal{B}(\pi_n(X))$  if  $\mathcal{B}(\pi_n(X))$  is non-empty.
- (2)  $\text{cdg}(X, n) \neq 0$  if and only if  $\text{rank } \pi_n(X) = \text{rank } \langle \text{Cdg}(X, n) \rangle \geq 1$ , where  $\langle \text{Cdg}(X, n) \rangle$  is the subgroup of  $\text{Hom}(\pi_n(X), \pi_n(S^n))$  generated by  $\text{Cdg}(X, n)$ .
- (3)  $\text{cdg}(X, n) \neq 0$  if and only if  $\text{rank } \pi_n(X) \geq 1$  and there exists an integer  $r \geq 1$  such that  $r \cdot \text{Hom}(\pi_n(X), \pi_n(S^n)) \subset \langle \text{Cdg}(X, n) \rangle$ . In the latter case  $\text{cdg}(X, n)$  is equal to the least of such  $r$ .

*Stable version also holds.*

*Proof.* Put  $t = \text{rank } \pi_n(X)$  and  $s = \text{rank } \langle \text{Cdg}(X, n) \rangle$ . We denote by  $\{a_1, \dots, a_t\}$  and  $\{\alpha_1, \dots, \alpha_t\}$  a free basis of  $\text{Hom}(\pi_n(X), \pi_n(S^n))$  and its dual basis of  ${}^{ab}\pi_n(X)/\text{Tor}$ , respectively.

First we prove (2). Suppose  $\text{cdg}(X, n) \neq 0$ . Then trivially  $t \geq 1$ . To induce a contradiction, suppose  $t > s$ . Then we can take  $\{a_1, \dots, a_t\}$  satisfying  $\langle \text{Cdg}(X, n) \rangle \subset \langle a_1, \dots, a_s \rangle$ . It follows that  $f_*(\alpha_i) = 0$  for all  $f: X \rightarrow S^n$ , hence  $\text{cdg}(\alpha_i) = 0$  and  $\text{cdg}(X, n) = 0$ . This is a contradiction. Hence  $t = s$ . Conversely suppose that  $t = s \geq 1$  and  $\text{cdg}(X, n) = 0$ . Then we can take  $\{\alpha_1, \dots, \alpha_t\}$  satisfying  $\text{cdg}(\alpha_i) = 0$ . It follows that  $\text{Cdg}(X, n) \subset \langle a_1, \dots, a_{t-1} \rangle$  so that  $s \leq t - 1$ . This is a contradiction. Hence  $\text{cdg}(X, n) \neq 0$  if  $t = s \geq 1$ . This proves (2).

Next we prove (1). If  $\text{cdg}(X, n) = 0$ , then there is no problem. So suppose that  $\text{cdg}(X, n) \neq 0$ . Then  $t = s \geq 1$  as shown above. Choose  $\{a_i; 1 \leq i \leq t\}$  such that  $\{k_i a_i; 1 \leq i \leq t\}$  is a basis of  $\langle \text{Cdg}(X, n) \rangle$ , where  $k_i \geq 1$ . Put  $k = \text{LCM}\{k_i\}$ . Then

$$k = \text{Min}\{r > 0; r \cdot \text{Hom}(\pi_n(X), \pi_n(S^n)) \subset \langle \text{Cdg}(X, n) \rangle\}$$

and hence  $k \cdot \text{Hom}(\pi_n(X), \pi_n(S^n)) \subset \langle \text{Cdg}(X, n) \rangle$ , where Min denotes the minimum. Evaluating at any  $\beta = \sum c_i \alpha_i \in {}^{ab}\pi_n(X)/\text{Tor}$ , we have  $k \cdot \text{GCD}\{c_i\} \mathbf{Z} \subset \text{cdg}(\beta) \mathbf{Z} = \text{GCD}\{k_i c_i\} \mathbf{Z}$  so that  $\text{cdg}(\beta) = \text{GCD}\{k_i c_i\} | k \cdot \text{GCD}\{c_i\}$ . If  $\beta \in \mathcal{B}(\pi_n(X))$ , then  $\text{GCD}\{c_i\} = 1$  and  $\text{cdg}(\beta) | k$ , hence  $\text{cdg}(X, n) | k$ . Set  $d_i = k/k_i$  and  $\alpha = \sum d_i \alpha_i$ . Then  $\text{GCD}\{d_i\} = 1$  and  $\alpha \in \mathcal{B}(\pi_n(X))$ . We then have  $\text{cdg}(\alpha) = k$ , so  $\text{cdg}(X, n) = \text{cdg}(\alpha) = k$ . This proves (1) and a part of (3).

Other part of (3) follows immediately from (2). The same proof is valid for stable case. This completes the proof of Proposition 2.3.

**Proposition 2.4.** (1) *If all of the following five conditions are satisfied, then  $\text{cdg}(X, n)$  is non-zero.*

- (i)  $X$  is a finite CW-complex.
- (ii)  $\text{rank } \pi_n(X) \geq 1$ .

- (iii)  $X$  is simply connected if  $n \geq 2$ .  
 (iv) Image  $\{\pi_n(X^{(n-1)}) \rightarrow \pi_n(X)\}$  is a torsion, where  $X^{(k)}$  is the  $k$ -skeleton of  $X$ .  
 (v) All attaching maps of  $2n$ -cells in  $X/X^{(n-1)}$  are null homotopic if  $n$  is even.  
 (2) If  $X$  is a finite CW-complex with  $\text{rank}^s \pi_n(X) \geq 1$ , then  ${}^s\text{cdg}(X, n)$  is non-zero.

Proof. The assertions for  $n=1$  can be proved by using the facts that the composite of  $[X, S^1] \cong H^1(X) \cong \text{Hom}(H_1(X), \mathbf{Z}) \cong \text{Hom}(\pi_1(X), \mathbf{Z})$  is  $\text{Cdg}$  and that  ${}^{ab}\pi_1(X) \cong {}^s\pi_1(X)$ .

Suppose  $n \geq 2$  and five conditions in (1).

First we shall show that  $\text{Cdg}$  is a surjection on  $[X^{(n+1)}/X^{(n-1)}, S^n]$ . This is trivial if  $X$  has no  $(n+1)$ -cell, so we assume that  $X$  has  $(n+1)$ -cells. We then have a cofibre sequence  $\vee S^n \xrightarrow{p} \vee S^n \xrightarrow{i} X^{(n+1)}/X^{(n-1)}$  and the commutative diagram:

$$\begin{array}{ccccc} [\vee S^n, S^n] & \xleftarrow{p^*} & [\vee S^n, S^n] & \xleftarrow{i^*} & [X^{(n+1)}/X^{(n-1)}, S^n] \\ \text{Cdg} \downarrow \cong & & \text{Cdg} \downarrow \cong & & \downarrow \text{Cdg} \\ \text{Hom}(\pi_n(\vee S^n), \mathbf{Z}) & \xleftarrow{p_*^*} & \text{Hom}(\pi_n(\vee S^n), \mathbf{Z}) & \xleftarrow{i_*^*} & \text{Hom}(\pi_n(X^{(n+1)}/X^{(n-1)}), \mathbf{Z}) \end{array}$$

In this diagram, the upper horizontal sequence is the same as the stable one and hence exact,  $i_*^*$  is a monomorphism, and  $p_*^* \circ i_*^* = 0$ . By chasing the diagram, it follows that the third  $\text{Cdg}$  is a surjection.

Given any  $a \in \text{Hom}(\pi_n(X^{(n+1)}/X^{(n-1)}), \mathbf{Z})$ , choose  $b: X^{(n+1)}/X^{(n-1)} \rightarrow S^n$  such that  $\text{Cdg}(b) = a$ . By (v) and [1, 3.1], we can construct skeleton-wise a map  $f: X/X^{(n-1)} \rightarrow S^n$  such that  $f \circ i = k \circ b$  for some  $k \neq 0$ , where  $i: X^{(n+1)}/X^{(n-1)} \subset X/X^{(n-1)}$ . This implies that  $\langle \text{Cdg}(X/X^{(n-1)}), n \rangle$  is of maximal rank, since

$$i_*^*: \text{Hom}(\pi_n(X/X^{(n-1)}), \mathbf{Z}) \cong \text{Hom}(\pi_n(X^{(n+1)}/X^{(n-1)}), \mathbf{Z}).$$

By (iii) and a theorem of Blakers-Massey,  $\pi_n(X, X^{(n-1)}) \cong \pi_n(X/X^{(n-1)})$ . Then by (iv) the homomorphism

$$q_*^*: \text{Hom}(\pi_n(X/X^{(n-1)}), \mathbf{Z}) \rightarrow \text{Hom}(\pi_n(X), \mathbf{Z})$$

induced by the quotient map  $q$  has a finite cokernel. Therefore  $\langle \text{Cdg}(X, n) \rangle$  is of maximal rank, since  $q_*^* \langle \text{Cdg}(X/X^{(n-1)}), n \rangle \subset \langle \text{Cdg}(X, n) \rangle$ . Hence  $\text{cdg}(X, n) \neq 0$  by Proposition 2.3. This proves (1). By almost the same proof as the above, we have (2).

The following two results can be proved easily. So we omit their proofs.

**Proposition 2.5.** (1) If  $X$  is  $k$ -connected with  $n \leq 2k+1$  and  $\mathcal{B}({}^s\pi_n(X))$  is non-empty, then  ${}^s\text{cdg}(X, n) \mid \text{cdg}(X, n)$ .

(2) If  $\text{rank} \pi_n(X) = \text{rank} {}^s\pi_n(X) = 1$ , then  $m \cdot {}^s\text{cdg}(X, n) \mid \text{cdg}(X, n)$ , where

the suspension  $\Sigma^\infty: {}^{ab}\pi_n(X)/\text{Tor}=\mathbf{Z}\rightarrow {}^s\pi_n(X)/\text{Tor}=\mathbf{Z}$  is multiplication by  $m$ .

(3) If  $G$  is a connected simple Lie group, then  $\text{rank } \pi_3(G)=\text{rank } {}^s\pi_3(G)=1$  and

$${}^s\text{cdg}(\tilde{G}, 3) | m \cdot {}^s\text{cdg}(G, 3) | \text{cdg}(G, 3)$$

where  $\tilde{G}$  is a universal covering group of  $G$  and  $m$  is a non-zero integer defined as in (2) for  $X=G$ .

We denoted  $m \cdot {}^s\text{cdg}(G, 3)$  in 2.5(3) by  $\text{cd}(G)$  in [9]. Hence  ${}^s\text{cdg}(G, 3)=\text{cd}(G)$  if  $G$  is simple and simply connected.

**Proposition 2.6.** (1) If  $\mathcal{B}(\pi_n(X_i))$  is non-empty for  $i=1, 2$ , then  $\text{LCM}\{\text{cdg}(X_1, n), \text{cdg}(X_2, n)\} | \text{cdg}(X_1 \times X_2, n)$ . Stable version also holds.

(2) If  ${}^{ab}\pi_n(X_1)$  is a torsion, then  $\text{cdg}(X_1 \times X_2, n)=\text{cdg}(X_2, n)$ .

(3) If  $\pi_n(X_i)$  is of finite rank and  $\text{cdg}(X_i, n) \neq 0$  for  $i=1, 2$ , then  $\text{cdg}(X_1 \times X_2, n)=\text{LCM}\{\text{cdg}(X_1, n), \text{cdg}(X_2, n)\}$ .

(4) If  $X_i$  is  $(n-1)$ -connected and  ${}^s\pi_n(X_i)$  is of finite rank for  $i=1, 2$ , then  ${}^s\text{cdg}(X_1 \times X_2, n) | {}^s\text{cdg}(X_1, n) \cdot {}^s\text{cdg}(X_2, n)$ .

### 3. Proof of Theorem

In this section  $G$  denotes a compact connected Lie group of type  $\{n_1, \dots, n_r\}$ , that is,  $H^*(G; \mathbf{Q}) \cong H^*(\prod_{i=1}^r S^{n_i}; \mathbf{Q})$ . As is well-known,  $n_i$  is odd and there are maps  $f: \prod_i S^{n_i} \rightarrow G$  and  $g: G \rightarrow \prod S^{n_i}$  which induce isomorphisms  $\pi_*(\prod S^{n_i}) \otimes \mathbf{Q} \cong \pi_*(G) \otimes \mathbf{Q}$  (see [7]). From this and Proposition 2.3 we have

**Proposition 3.1.** The following five statements are equivalent.

- (1)  $\text{Cdg}(G, n)$  is non-trivial.
- (2)  $\text{cdg}(G, n)$  is non-zero.
- (3)  $\text{rank } \pi_n(G)=\text{rank } \langle \text{Cdg}(G, n) \rangle \geq 1$ .
- (4)  $\text{rank } \pi_n(G) \geq 1$ .
- (5)  $n \in \{n_1, \dots, n_r\}$ .

*Proof of Theorem 1.* Numbers  $\text{cdg}(G, n)$  and  ${}^s\text{cdg}(G, n)$  are finite by Proposition 2.3. Put  $A(n)=\{i; \text{rank } \pi_n(G_i) \geq 1\}$  and define  ${}^sA(n)$  similarly. Then  $A(3)={}^sA(3)=\{i; 1 \leq i \leq \ell\}$ . We have  ${}^s\text{cdg}(G, 3) | \text{cdg}(G, 3)$  by 2.5 (1). Thus (1) implies (2). We have  $\text{LCM}\{{}^s\text{cdg}(G_i, 3)\} | {}^s\text{cdg}(G, 3)$  and  $\text{cdg}(G, n)=\text{LCM}\{\text{cdg}(G_i, n); i \in A(n)\}$  by 2.6. Hence (2) implies (4), and (1) and (3) are equivalent. By Theorem 4.1 (1) of [9], (4) and (5) are equivalent. Trivially (5) implies (6), and (7) implies (1).

To prove that (6) implies (7), suppose (6). By Proposition 3.1, we may suppose that  $n \in \{n_1, \dots, n_r\}$ . Then there is a  $p$ -equivalence  $f: G \rightarrow S = \prod_{i=1}^r S^{n_i}$  so that  $\text{rank } \pi_n(G)=\text{rank } \pi_n(S)=u$ , say, and the image of  $f_*: \pi_n(G) \rightarrow \pi_n(S)$  is of maximal rank. Let  $\{\alpha_1, \dots, \alpha_u\}$  be a free basis of  $\pi_n(G)/\text{Tor}$  and  $\{a_1, \dots, a_u\}$  its

dual basis of  $\text{Hom}(\pi_n(G), \pi_n(S^n))$ . Let  $\{k_1, \dots, k_n\}$  be positive integers and  $\{\beta_1, \dots, \beta_n\}$  a free basis of  $\pi_n(S)$  such that  $f_*(\alpha_i) = k_i\beta_i$ . Then  $k_i$  is prime to  $p$ . Since  $f_*^* \circ \text{Cdg} = \text{Cdg} \circ f^* : [S, S^n] \rightarrow \text{Hom}(\pi_n(G), \pi_n(S^n))$  and since  $\text{Cdg}$  is surjective on  $[S, S^n]$ , we have  $\text{Cdg}(G, n) \supset \text{Image}(f_*^*) = \bigoplus_{i=1}^n k_i \mathbf{Z}\{a_i\}$ . Hence  $\text{Cdg}(G, n)$  contains  $\text{LCM}\{k_i\} \cdot \text{Hom}(\pi_n(G), \pi_n(S^n))$  so that  $\text{cdg}(G, n) \mid \text{LCM}\{k_i\}$  by Proposition 2.3 (3), therefore  $\text{cdg}_p(G, n) = 0$ . This implies (7) and completes the proof of Theorem 1.

EXAMPLE 3.2. For  $G$  non-simply connected, Theorem 1 does not hold in general:  $\text{cdg}(SO(3), 3) = 2$  and  ${}^s\text{cdg}(SO(3), 3) = 1$  (see [10]).

Recall that if  $G$  is simple then  $n \in \{n_1, \dots, n_r\}$  if and only if  $\text{rank } \pi_n(G)$  is 1 or 2 and  $\text{rank } \pi_n(G) = 2$  if and only if  $(\tilde{G}, n) = (\text{Spin}(4m), 4m - 1)$  for  $m \geq 2$ . Then the following and Proposition 3.1 prove Theorem 2 except for the case  $(G, n) = (\text{Spin}(4m), 4m - 1)$ .

**Proposition 3.3** (James). *If  $n$  is odd, then the image of  $\alpha^* : [X, S^n] \rightarrow \pi_n(S^n)$  is a subgroup for every  $\alpha \in \pi_n(X)$ . In particular if  $n$  is odd and  $\text{rank } \pi_n(X) = 1$ , then  $\text{Cdg}(X, n) = \text{cdg}(X, n) \cdot \text{Hom}(\pi_n(X), \pi_n(S^n))$ .*

Proof. The first assertion can be proved by the method in [3, p.88]. The second assertion then follows, since  $\alpha^* = \text{ev}_\alpha \circ \text{Cdg}$  and  $\text{ev}_\alpha$  is an isomorphism if  $\text{rank } \pi_n(X) = 1$  and  $\alpha$  represents a generator of  ${}^{ab}\pi_n(X)/\text{Tor} = \mathbf{Z}$ , where  $\text{ev}_\alpha : \text{Hom}(\pi_n(X), \pi_n(S^n)) \rightarrow \pi_n(S^n)$  is the evaluation at  $\alpha$ , that is,  $\text{ev}_\alpha(\theta) = \theta(\alpha)$ .

Let

$$\text{Spin}(4m - 1) \xrightarrow{i} \text{Spin}(4m) \xrightarrow{p} S^{4m - 1}$$

be the canonical bundle for  $m \geq 1$ . Then we have

$$\begin{aligned} \pi_{4m - 1}(\text{Spin}(4m)) &= \mathbf{Z}\{s_1\} \oplus \mathbf{Z}\{s_2\}, \\ \text{Hom}(\pi_{4m - 1}(\text{Spin}(4m)), \pi_{4m - 1}(S^{4m - 1})) &= \mathbf{Z}\{s'_1\} \oplus \mathbf{Z}\{s'_2\} \end{aligned}$$

where  $s_1$  is the image under  $i_*$  of a generator of  $\pi_{4m - 1}(\text{Spin}(4m - 1)) = \mathbf{Z}$  and  $s_2$  is an element such that  $p_*(s_2)$  is 2 if  $m \geq 3$  and 1 if  $m \leq 2$  (cf., [5]);  $s'_j$  is the dual element to  $s_j$ . Then the following completes the proof of Theorem 2.

**Proposition 3.4.** *The number  $\text{cdg}(\text{Spin}(4m), 4m - 1)$  is non-zero and*

$$\text{Cdg}(\text{Spin}(4m), 4m - 1) = \text{cdg}(\text{Spin}(4m), 4m - 1) \mathbf{Z}\{s'_1\} \oplus c \mathbf{Z}\{s'_2\}$$

where  $c$  is 2 if  $m \geq 3$  and 1 if  $m \leq 2$ .

Proof. If  $m \leq 2$ , then  $\text{Spin}(4m) \approx \text{Spin}(4m - 1) \times S^{4m - 1}$  and the assertion can be obtained easily.

Suppose that  $m \geq 3$ . Then  $s_2^*(p) = 2$ , hence  $\text{cdg}(s_2) = 2$  by the following

lemma.

**Lemma 3.5.** *If  $X$  is an  $H$ -space and  $n$  is odd with  $n \neq 1, 3, 7$ , then  $\text{cdg}(\alpha)$  is even for every  $\alpha \in \pi_n(X)$ .*

To simplify notations, we set  $(G, n) = (\text{Spin}(4m), 4m-1)$ . By definition, we have

$$\text{Cdg}(G, n) \subset \text{cdg}(s_1)\mathbf{Z}\{s'_1\} \oplus 2\mathbf{Z}\{s'_2\}.$$

Take any integers  $k_1$  and  $k_2$ . Then there exists a map  $f: G \rightarrow S^n$  such that  $\text{Cdg}(f) = \text{cdg}(s_1)k_1s'_1 + 2js'_2$  for some integer  $j$ . Let  $I: G \rightarrow G$  be the inversion, that is,  $I(A) = A^{-1}$ . Then  $\text{Cdg}$  of the composition of

$$G \xrightarrow{d} G \times G \xrightarrow{1 \times f} G \times S^n \xrightarrow{g_{\pm}} S^n$$

is  $\text{cdg}(s_1)k_1s'_1 + (2j \pm 2)s'_2$ , where  $d$  is the diagonal map,  $g_+$  the canonical action and  $g_- = g_+ \circ (I \times 1)$ . Inductively we then have  $\text{cdg}(s_1)k_1s'_1 + 2k_2s'_2 \in \text{Cdg}(G, n)$ . Hence  $\text{Cdg}(G, n) = \text{cdg}(s_1)\mathbf{Z}\{s'_1\} \oplus 2\mathbf{Z}\{s'_2\}$ . Also  $\text{cdg}(s_1)$  is even from Lemma 3.5, hence  $\text{cdg}(G, n) = \text{cdg}(s_1) \neq 0$  from Proposition 2.3(3) and the following lemma.

**Lemma 3.6.**  $\text{cdg}(s_1) \neq 0$ .

*Proof of 3.5.* Let  $g: X \rightarrow S^n$  be a map such that  $g \circ \alpha = \text{cdg}(\alpha) \in \pi_n(S^n) = \mathbf{Z}$ . Then the degree of the composition of

$$S^n \xrightarrow{i_j} S^n \times S^n \xrightarrow{\alpha \times \alpha} X \times X \xrightarrow{\mu} X \xrightarrow{g} S^n$$

is  $\text{cdg}(\alpha)$  for  $j=1, 2$ , where  $i_j$  is the inclusion to the  $j$ -th factor and  $\mu$  is the multiplication. Hence  $\text{cdg}(\alpha)^2[\iota_n, \iota_n] = [\text{cdg}(\alpha)\iota_n, \text{cdg}(\alpha)\iota_n] = 0$ , so  $\text{cdg}(\alpha)$  is even, because the Whitehead square  $[\iota_n, \iota_n]$  of the identity map  $\iota_n$  of  $S^n$  is of order 2.

*Proof of 3.6.* Set  $n = 4m - 1$ . Then the homomorphism  $\pi_n(\text{Spin}(n)) = \mathbf{Z} \rightarrow \pi_n(\text{Spin}(n+2)) = \mathbf{Z}$  induced by the inclusion is multiplication by  $e$ , where  $e$  is 1 if  $m \geq 3$  and 2 if  $m \leq 2$ . Thus we have  $\text{cdg}(\text{Spin}(n+1), n; s_1) | e \cdot \text{cdg}(\text{Spin}(n+2), n)$ . Since the latter number is non-zero by Proposition 3.1, so is the former.

This completes the proofs of Proposition 3.4 and Theorem 2.

**REMARK 3.7** ([10]). By almost the same proof as the above, we can prove that  $\text{Cdg}(SO(m), n)$  is a subgroup of maximal rank. By using Proposition 4.1 below, we can prove that if  $G$  is simple but not necessarily simply connected, then  $\text{Cdg}(G, n)$  contains a subgroup of maximal rank.

#### 4. Other results

We give three results. See [6] and [10] for their proofs. When we study  $\text{Cdg}(G, n)$  for non-simply connected  $G$ , the following is useful.

**Proposition 4.1.** *Let  $q: H \rightarrow G$  be a finite covering homomorphism and  $m$  the least positive integer such that  $x^m = 1$  for all  $x$  in the kernel of  $q$ . Then we have*

- (1)  $m \cdot \text{Cdg}(H, n) \subset q_*^* \text{Cdg}(G, n) \subset \text{Cdg}(H, n)$ ,
- (2)  $\text{cdg}(\beta) | \text{cdg}(q_*\beta) | m \cdot \text{cdg}(\beta)$  for every  $\beta \in \pi_n(H)$ ,
- (3)  $\text{cdg}(H, n) | \text{cdg}(G, n) | m \cdot \text{cdg}(H, n)$  for  $n \geq 2$ ,
- (4)  $\text{cdg}(H, 1) | m$ .

Let  $\Xi: \pi_n(X) \rightarrow H_n(X)$  be the Hurewicz homomorphism. Put  $PH_n(X) = \{x \in H_n(X); d_*(x) = x \otimes 1 + 1 \otimes x\}$ , where  $d: X \rightarrow X \times X$  is the diagonal map. As is easily seen,  $\Xi(\pi_n(X)) \subset PH_n(X)$ . It is known as a theorem of Cartan-Serre that  $\Xi \otimes \mathbb{Q}: \pi_*(G) \otimes \mathbb{Q} \cong PH_*(G) \otimes \mathbb{Q}$ . L. Smith[11] studied the problem: What is the smallest positive integer  $N(G, n)$  such that  $N(G, n)x$  is contained in the image of the modulo torsion Hurewicz homomorphism

$$\Xi: \pi_n(G)/\text{Tor} \rightarrow PH_n(G)/\text{Tor}$$

for every  $x \in PH_n(G)/\text{Tor}$ ?

**Proposition 4.2.** *If  $G$  is simple or simply connected, then  $\text{cdg}(G, n)$  is a multiple of  $N(G, n)$ .*

**EXAMPLE 4.3.** The number  $N(G, n)$  has been determined for classical groups,  $G_2$  and  $F_4$  (see e.g., [4]). The first few values of the Smith's upper bound  $N(n)$  of  $N(G, n)$  are  $N(3)=1$ ,  $N(5)=2^2$ ,  $N(7)=2^4 \cdot 3$ ,  $N(9)=2^6 \cdot 3$ ,  $N(11)=2^8 \cdot 3^2 \cdot 5$  (see[11]). If  $G$  is simple and simply connected, then  $N(G, 3)=1$  and  $\text{cdg}(G, 3)$  is even except for  $G=S^3$ . We have  $N(SU(3), 5)=\text{cdg}(SU(3), 5)=2$ ;  $N(Sp(2), 7)=\text{cdg}(Sp(2), 7)=2^2 \cdot 3=N(Sp(3), 7)$ ;  $2^5 \cdot 3 | \text{cdg}(Sp(3), 7) | 2^8 \cdot 3$ ;  $N(SU(5), 9)=\text{cdg}(SU(5), 9)=2^3 \cdot 3$ ;  $N(G_2, 11)=\text{cdg}(G_2, 11)=2^3 \cdot 3 \cdot 5$ .

---

#### References

- [1] M. Arkowitz and C.R. Curjel: *The Hurewicz homomorphism and finite homotopy invariants*, Trans. Amer. Math. Soc. **101** (1964), 538–551.
- [2] M.C. Crabb and K. Knapp: *James numbers*, Math. Ann. **282** (1988), 395–422.
- [3] I.M. James: *Symmetric functions of several variables, whose range and domain is a sphere*, Bol. Mat. Mex. **1** (1956), 85–88.
- [4] R. Kane and G. Moreno: *Spherical homology classes in the bordism of Lie groups*, Can. J. Math. **40** (1988), 1331–1374.
- [5] M.A. Kervaire: *Some nonstable homotopy groups of Lie groups*, Ill.J. Math. **4** (1960), 161–169.

- [6] A. Kono and H.  $\bar{O}$ shima: *Codegree of simple Lie groups-II*, Osaka J. Math. **28** (1991), 129–139.
- [7] M. Mimura and H. Toda: *On  $p$ -equivalences and  $p$ -universal spaces*, Comment. Math. Helv. **46** (1971), 87–97.
- [8] H.  $\bar{O}$ shima: *Remarks on the  $j$ -codegree of vector bundles*, Japanese J. Math. **16** (1990), 97–117.
- [9] H.  $\bar{O}$ shima: *Codegree of simple Lie groups*, Osaka J. Math. **26** (1989), 759–773.
- [10] H.  $\bar{O}$ shima and K. Takahara: *Codegrees and Lie groups*, Mimeographed note, 1990.
- [11] L. Smith: *On the relation between spherical and primitive homology classes in topological groups*, Topology **8** (1969), 69–80.
- [12] H. Toda: *Composition methods in homotopy groups of spheres*, Ann. Math. Stud. **49**, Princeton Univ. Press, Princeton, 1962.

Hideaki  $\bar{O}$ shima  
Department of Mathematics  
Osaka City University  
Sugimoto, Sumiyoshi-ku  
Osaka 558, Japan

Koji Takahara  
Department of Mathematics  
Osaka City University  
Sugimoto, Sumiyoshi-ku  
Osaka 558, Japan

and  
Fujitsu  
Miyamoto, Numazu  
Shizuoka 410-03, Japan

