

Title	Everolimus Rescue Treatment for Chronic Rejection After Pediatric Living Donor Liver Transplantation: 2 Case Reports
Author(s)	Ueno, T.; Hiwatashi, S.; Saka, R. et al.
Citation	Transplantation Proceedings. 2018, 50(9), p. 2872-2876
Version Type	AM
URL	https://hdl.handle.net/11094/97129
rights	© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Figure captions:

Fig. 1 Case 1 course summary: Liver biopsies were performed pre and post treatment with everolimus. ALT: alanine aminotransferase; LBx: liver biopsy; rATG: rabbit antithymoglobulin; T-Bil : total bilirubin

Fig. 2 Case 1 liver biopsy findings: LBx1 showed pre everolimus treatment liver biopsy. H&E staining showed lymphocyte infiltration in the portal tract. Azan staining showed possible bridging fibrosis. CK7 staining showed damaged bile duct. LBx2 showed post everolimus treatment liver biopsy. H&E stain showed improved lymphocyte infiltration. Azan and CK7 staining showed better fibrosis and improved bile ducts. H&E: hematoxylin and eosin; CK7: cytokeratin 7

Fig. 3 Case 2 course summary : Liver biopsies were performed pre and post treatment with everolimus. ALT: alanine aminotransferase; LBx: liver biopsy; rATG: rabbit antithymoglobulin; T-Bil : total bilirubin

Fig. 4 Case 2 liver biopsy findings: LBx3 showed pre everolimus treatment liver biopsy. H&E and azan staining showed fibrosis of portal area. CK7 staining showed proliferation of pseudocholeangiolar, almost ductpenia. LBx4 showed post everolimus treatment liver biopsy. H&E and azan staining showed improved fibrosis. CK7 staining showed normal bile ducts. H&E: hematoxylin and eosin; CK7: cytokeratin 7

