



|              |                                                                             |
|--------------|-----------------------------------------------------------------------------|
| Title        | Structural properties of functional-differential equations in Banach spaces |
| Author(s)    | Nakagiri, Shin-ichi                                                         |
| Citation     | Osaka Journal of Mathematics. 1988, 25(2), p. 353-398                       |
| Version Type | VoR                                                                         |
| URL          | <a href="https://doi.org/10.18910/9745">https://doi.org/10.18910/9745</a>   |
| rights       |                                                                             |
| Note         |                                                                             |

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## STRUCTURAL PROPERTIES OF FUNCTIONAL DIFFERENTIAL EQUATIONS IN BANACH SPACES

SHIN-ICHI NAKAGIRI

(Received November 13, 1986)

### Contents

1. Introduction and summary of the results
2. Linear functional differential equations in Banach spaces
  - 2.1. Notation and terminology.
  - 2.2. Fundamental solution, mild solution and retarded resolvent
3. Semigroups associated with functional differential equations
4. Structural operators  $F$ ,  $G$  and their adjoint operators
5. Characterizations of  $\text{Ker } F$  and  $\text{Im } F$
6. Representations of resolvent operators
7. Spectral decomposition
  - 7.1. Classification of spectrum
  - 7.2. Generalized eigenspaces and spectral decomposition
8. Adjoint spectral decomposition
  - 8.1. Generalized eigenspaces and structural operators
  - 8.2. Representations of spectral projections
9. Completeness of generalized eigenfunctions
10. Illustrative examples

### 1. Introduction and summary of the results

In the present work we study the structural properties of linear autonomous functional differential equations in Banach spaces within the framework of linear operator theory. We shall explain our motivation of this study.

In a series of papers Bernier and Manitius [4], Manitius [29] and Delfour and Manitius [15] they have developed an excellent state space theory for linear retarded functional differential equations (FDE's) in the product space  $\mathbf{R}^n \times L_p([-h, 0]; \mathbf{R}^n)$ ,  $h > 0$ . The theory is based on certain relations between semigroups associated with the FDE's and the so-called structural operators  $F$  and  $G$ . The structural operators have enriched the qualitative theory of linear FDE's and have provided various new and efficient techniques for the study of control

theory involving retarded FDE's. The power of the theory has been shown and increased by a number of contributions (refer to Delfour [12], Delfour, Lee and Manitius [14], Manitius [30,31], Salamon [39] and Vinter and Kwong [47]). Recently Salamon has been extended the state space theory to the controlled neutral FDE's and has used the theory to expand a system theory for neutral systems in his book [40].

We now pose a question. Is it possible to construct an analogous theory for partial FDE's? For the question we shall give an affirmative answer for certain class of partial FDE's. We study partial FDE's in the class consists of abstract delay evolutionary equations in Banach spaces similarly as in Travis and Webb [44,45], Webb [48], Datko [9, 10] and Kunisch and Schappacher [26, 27]. Partial FDE's in this class are very general and are appropriate for system theoretical study as shown in Curtain and Pritchard [8] and Fuhrmann [16], so we take this class. Let  $X$  be a reflexive Banach space and consider the evolution equation with delay

$$(E) \quad \frac{dx(t)}{dt} = A_0 x(t) + \int_{-h}^0 d\eta(s)x(t+s), \quad t > 0$$

on  $X$ , where  $A_0$  generates a  $C_0$ -semigroup and  $\eta$  is a bounded Stieltjes measure on  $I_h = [-h, 0]$ . We study the equation (E) on the state space  $M_p = X \times L_p(I_h; X)$ . The structural operator  $F$  is concerned with the retarded part of (E) and is defined through the measure  $\eta$  quite similarly as in [4]. In [33, 34] the author has constructed the fundamental solution of (E) under the natural condition on  $\eta$  and has shown its prominent role in the optimal control theory involving (E). The introduction of fundamental solution permits us to define the structural operator  $G$ , and these  $G$  and  $F$  have made it possible to develop the state space for (E).

The objective of this paper is to extend and give certain new contributions to the state space theory for the equation (E) on a reflexive Banach Space. Many results obtained here, which are useful in applications, are considered to be possible generalizations of the results in [4, 15, 29] to infinite dimensions. However it is also the objective of this paper to propose an approach for simplifying the state space theory. Due to our approach heavily depending on functional analysis method, many of the proofs can be improved. The author believes that the results presented here will provide a useful tool in studying the control theory for partial FDE's.

We enumerate the contents of this paper. Section 2 gives some preliminary results on the equation (E). The notations and terminology to be used for (E) are given in Subsection 2.1. In Subsection 2.2 various fundamental concepts relating to (E) are introduced; e.g., the fundamental solution  $W(t)$ ,

the retarded resolvent  $R(\lambda; A_0, \eta)$  which is a bounded inverse of  $\Delta(\lambda) = \lambda I - A_0 - \int_{-h}^0 e^{\lambda s} d\eta(s)$ , the three kinds of retarded, point spectrum  $\sigma_p(A_0, \eta)$ , continuous spectrum  $\sigma_c(A_0, \eta)$ , and the residual spectrum  $\sigma_R(A_0, \eta)$ , the mild solution; and the basic fact that  $R(\lambda; A_0, \eta)$  is given by the Laplace transform of  $W(t)$  for  $\text{Re } \lambda$  large is stated. Also, a variation of constants formula for the mild solution in terms of  $W(t)$ , which is essential in our treatment, is given. In the remainder part of this subsection we introduce the transposed equation  $(E^T)$  on the adjoint space  $M_p^*$  of  $M_p$  and give an elementary adjoint theory. In Section 3 we define semigroups  $S(t)$  and  $S_T(t)$  associated with  $(E)$  and  $(E^T)$  respectively, by the translation segments of mild solutions. The basic properties of the semigroups like infinitesimal generator or compactness for  $t > h$  are investigated as well as those for their adjoint semigroups  $S^*(t)$  and  $S_T^*(t)$ . Section 4 is devoted to study the properties of structural operators. As in Bernier and Manitius [4] we define the structural operators  $F$  and  $G_t$ ,  $t > 0$ , then a key relation  $S(t) = G_t F$ ,  $t \geq h$  in our theory follows from the variation of constants formula. The representations of the adjoints  $F^*$  and  $G_t^*$  are shown to be of same type as  $F$  and  $G_t$ , so analogous decomposition for  $S_T(t)$  holds. As a consequence of such decompositions we can show that the adjoint semigroup  $S^*(t)$  is realized via a modified transposed equation with non-zero forcing term by regarding the forcing term as the initial state of the transposed equation. Other fundamental properties of  $F$  and  $G = G_h$  shown in [4, 15, 29] are the intertwined property  $S(t)G = GS_T^*(t)$  and  $FS(t) = S_T^*(t)F$ . These relations are extended to our Banach space case and their simple proofs based on the new formula  $S(t)G = G_{t+h}$  are presented. It is also proved in Section 4 that the null space  $\text{Ker } G$  of  $G$  is  $\{0\}$  and the image  $\text{Im } G$  of  $G$  is dense in  $M_p$ . We know that similar conditions for  $F$  are hopeful in establishing good qualitative properties of  $(E)$ , but these are not true in general. Thus, in Section 5 we examine conditions for  $F$  such that  $\text{Ker } F = \{0\}$ ,  $\text{Im } F = M_p$  or  $\text{Cl}(\text{Im } F) = M_p$ , where  $\text{Cl}$  denotes the closure operation. A number of necessary and/or sufficient conditions for these criterion expressed by  $\eta$  are established by solving a Volterra integral equation with delays induced by the operator  $F$ . Among those it is shown that for differential difference equations with the retarded term  $\int_{-h}^0 d\eta(s)x(t+s) = \sum_{r=1}^m A_r x(t-h_r)$ ,  $0 < h_1 < \dots < h_m = h$ , an equivalent condition to  $\text{Ker } F = \{0\}$  is  $\text{Ker } A_m = \{0\}$ . Section 6 is devoted to studying the resolvent operators of infinitesimal generators which generates the semigroups given above. According to [7, 15, 40] various spectral operators containing exponential function terms are introduced and the relations each other and connections between  $F$  and/or  $\Delta(\lambda)$  are investigated. Using such relations we show, via the characterizations of generators given in Section 3, that each resolvent is described as a composition of  $F$  (or  $F^*$ ),

retarded resolvent and other spectral operators. Such representations for the resolvents play an important role in the spectral analysis for  $(E)$ . With the help of such forms a detailed and somewhat complicated spectral theory than [15], [18] is developed in Section 7 and Section 8. Section 7 studies the spectral decomposition theory for  $(E)$ . In Subsection 7.1 the spectrum of the generator  $A$  of  $S(t)$  is determined. The spectrum of  $A$  coincides with the retarded spectrum completely. Strictly speaking, it is shown that  $\sigma_p(A) = \sigma_p(A_0, \eta)$ ,  $\sigma_c(A) = \sigma_c(A_0, \eta)$  and  $\sigma_R(A) = \sigma_R(A_0, \eta)$ . In Subsection 7.2 a rather sophisticated spectral decomposition is presented. A characterization of the null space  $\text{Ker}(\lambda I - A)^l, l=1, 2, \dots$  in terms of  $\Delta(\lambda)$  and its derivatives is established for  $\lambda \in \sigma_p(A)$ . If  $\lambda$  is a pole of  $R(\mu; A_0, \eta)$  of order  $k_\lambda$ , then  $M_p$  can be decomposed as the direct sum of the generalized eigenspace  $\mathcal{M}_\lambda = \text{Ker}(\lambda I - A)^{k_\lambda}$  and its complementary space  $\mathcal{M}_\lambda^\perp = \text{Im}(\lambda I - A)^{k_\lambda}$ . In view of the representation of the resolvent  $R(\lambda; A)$  of  $A$  given in Section 6, the canonical spectral projection  $P_\lambda$  on  $\mathcal{M}_\lambda$  is expressed as a composition of  $F$  and other operators containing the retarded resolvent. Finally in this section we restrict a set  $\Lambda \subset \sigma(A)$  to a subset of discrete spectrum and establish the group property of  $S(t)$  on the decomposed space  $\mathcal{M}_\Lambda = \bigoplus_{\lambda \in \Lambda} \mathcal{M}_\lambda$  (direct sum) with a clear picture of the asymptotic behaviour of the mild solution of  $(E)$ . In Section 8 we develop the adjoint spectral decomposition theory by emphasizing the role of structural operators  $F$  and  $G$ . The main concern in Subsection 8.1 is to clarify the relation between the spectrums of the adjoint  $A^*$  of  $A$  and the generator  $A_T$  of  $S_T(t)$ . Thus it is shown that three kinds of spectrums of  $A^*$  and  $A_T$  coincide entirely and the generalized eigenspace  $\mathcal{M}_\lambda^*$  of  $A^*, \lambda \in \sigma_p(A^*)$  is given by  $\mathcal{M}_\lambda^* = F^* \mathcal{M}_\lambda^T$ , where  $\mathcal{M}_\lambda^T$  denotes the generalized eigenspace of  $A_T$  corresponding to  $\lambda$ . We now denote by  $\sigma_d(A^*)$  the discrete spectrum of  $A^*$ , i.e.,  $\dim \mathcal{M}_\lambda^* < \infty$  if  $\lambda \in \sigma_d(A^*) \subset \sigma_p(A^*)$ . Then it is also established that  $\sigma_d(A^*) = \sigma_d(A_T)$  and  $G^* \mathcal{M}_\lambda^* = \mathcal{M}_\lambda^T$ . This implies, by the property of  $G^*$ , that  $\dim \mathcal{M}_\lambda^* = \dim \mathcal{M}_\lambda^T$ . The last result in Subsection 8.1 gives the  $M_p$ -adjoint result for  $A$ , in which a fact that  $\dim \mathcal{M}_\lambda = \dim \mathcal{M}_\lambda^*$  is shown for a pole  $\lambda$  of  $R(\mu; A_0, \eta)$ . In Subsection 8.2 we are concerned with the representations of spectral projections. From the results in Subsection 8.1 we know that  $\dim \mathcal{M}_\lambda^* = \dim \mathcal{M}_\lambda^T < \infty$  for  $\lambda \in \sigma_d(A)$ . Using this fact the spectral projection  $P_\lambda$  for  $\lambda \in \sigma_d(A)$  is expressed in terms of the bases of  $\mathcal{M}_\lambda, \mathcal{M}_\lambda^T$  and the operator  $F$ . In Section 9 we study the problem of completeness of generalized eigenfunctions, which means  $\text{Cl}(\bigcup_{\lambda \in \sigma_p(A)} \mathcal{M}_\lambda) = M_p$ . First a characterization of the null space  $\text{Ker} P_\lambda$  for a pole  $\lambda$  of  $R(\mu; A_0, \eta)$  is given. Then a number of necessary and sufficient conditions for the completeness are established by the use of the representation of  $\text{Ker} P_\lambda$ . In the final Section 10 we give some examples of practical partial FDE's which illustrate the contents of this paper.

## 2. Linear functional differential equations in Banach spaces

### 2.1. Notation

The sets of real and complex numbers are denoted by  $\mathbf{R}^1$  and  $\mathbf{C}^1$ , respectively.  $\mathbf{R}^+$  denotes the set of non-negative numbers and  $\mathbf{R}^n$  denotes the  $n$ -dimensional Euclidean space. Let  $X$  and  $Y$  be complex (separable) Banach spaces with norms  $|\cdot|$  and  $\|\cdot\|_Y$ , respectively. For  $E \subset Y$  the closure of  $E$  is denoted by  $C1(E)$ . The adjoint spaces of  $X, Y$  are denoted by  $X^*, Y^*$  and their norms are denoted by  $|\cdot|_*, \|\cdot\|_{Y^*}$ , respectively. For a closed linear operator  $A$  on a dense domain  $D(A) \subset X$  into  $Y$ , its adjoint operator is denoted by  $A^*$ . The symbols  $\text{Im } A$  and  $\text{Ker } A$  will denote the image and the null space of  $A$ , respectively. The duality pairing between  $X$  and  $X^*$  is denoted by  $\langle \cdot, \cdot \rangle$  and the pairing between  $Y$  and  $Y^*$  by  $\langle \cdot, \cdot \rangle_Y$ . For  $E \subset Y$  the orthogonal complement  $\{y^* \in Y^* : \langle y, y^* \rangle_Y = 0 \text{ for all } y \in E\}$  of  $E$  is denoted by  $E^\perp$ .  $B(Y, X)$  denotes the Banach space of bounded linear operators from  $Y$  into  $X$ . When  $X = Y$ ,  $B(Y, X)$  is denoted by  $B(X)$ . Every operator norm simply is denoted by  $\|\cdot\|$ .

Given an interval  $I \subset \mathbf{R}^1$ ,  $L_p(I; X)$  and  $C(I; X)$  will denote the usual Banach space of  $X$ -valued measurable functions which are  $p$ -Bochner integrable ( $1 \leq p < \infty$ ) or essentially bounded ( $p = \infty$ ) on  $I$  and the Banach space of strongly continuous functions on  $I$ , respectively. The norm of  $L_p(I; X)$  is denoted by  $\|\cdot\|_{p,I}$ .  $W_p^{(1)}(I; X)$  denotes the Sobolev space of  $X$ -valued functions  $x(s)$  on  $I$  such that  $x(s)$  and its distributional derivative  $\dot{x}(s) = \frac{dx(s)}{ds}$  belong to  $L_p(I; X)$ .

For each integer  $k \geq 1$ ,  $C^k(I; X)$  denotes the Banach space of all  $k$ -times continuously differentiable functions from  $I$  into  $X$ .  $C(\mathbf{R}^+; X)$  (resp.  $L_p^{loc}(\mathbf{R}^+; X)$ ) will denote the Fréchet space of functions which belong to  $C([0, t]; X)$  (resp.  $L_p([0, t]; X)$ ) for any  $t > 0$ . Let  $M_p(I; X)$  denote the product space  $X \times L_p(I; X)$ . Given an element  $g \in M_p(I; X)$ ,  $g^0 \in X, g^1(\cdot) \in L_p(I; X)$  will denote the two coordinates of  $g$ , i.e.,  $g = (g^0, g^1)$ .  $M_p(I; X)$  is the Banach space with norm

$$\|g\|_{M_p(I; X)} = \begin{cases} (|g^0|^p + \|g^1\|_{p,I}^p)^{1/p} & \text{if } 1 \leq p < \infty \\ |g^0| + \|g^1\|_{\infty,I} & \text{if } p = \infty. \end{cases}$$

The symbol  $\chi_E$  denotes the characteristic function of the set  $E$ .

### 2.2. Fundamental solution, mild solution and retarded resolvent

We shall review some basic results on linear functional differential equations (FDE's) in Banach spaces. Let  $h > 0$  be fixed and  $I_h = [-h, 0]$ . Consider the following autonomous retarded FDE ( $E$ ) on a Banach space  $X$ :

$$(2.1) \quad \frac{dx(t)}{dt} = A_0 x(t) + \int_{-h}^0 d\eta(s)x(t+s) + u(t) \quad \text{a.e. } t \geq 0$$

$$(2.2) \quad x(0) = g^0, \quad x(s) = g^1(s) \quad \text{a.e. } s \in [-h, 0),$$

where  $g = (g^0, g^1) \in M_p \equiv M_p(I_h; X)$ ,  $u \in L_p^{\text{loc}}(\mathbf{R}^+; X)$ ,  $p \in [1, \infty]$ , and  $A_0$  generates a  $C_0$ -semigroup  $T(t)$  on  $X$ . The Stieltjes measure  $\eta$  in (2.1) is given by

$$(2.3) \quad \eta(s) = - \sum_{r=1}^m \mathcal{X}_{(-\infty, -h_r]}(s) A_r - \int_s^0 A_I(\xi) d\xi, \quad s \in I_h,$$

where  $0 < h_1 < \dots < h_m = h$ ,  $A_r \in B(X)$  ( $r = 1, \dots, m$ ) and  $A_I \in L_1(I_h; B(X))$ .

Let  $W(t)$  be the fundamental solution of (E), which is a unique solution of

$$W(t) = \begin{cases} T(t) + \int_0^t T(t-s) \int_{-h}^0 d\eta(\xi) W(\xi+s) ds, & t \geq 0 \\ 0, & t < 0. \end{cases}$$

Then  $W(t)$  is strongly continuous on  $\mathbf{R}^+$  and satisfies, for some  $M, \gamma_0 > 0$ ,

$$(2.4) \quad \|W(t)\| \leq M \exp(\gamma_0 t), \quad t \geq 0.$$

If the condition

$$(2.5) \quad A_I(\cdot) \in L_{p'}(I_h; B(X)), \quad 1/p + 1/p' = 1$$

is satisfied, then for each  $t \in \mathbf{R}^+$  the operator valued function  $U_t(\cdot)$  given by

$$(2.6) \quad \begin{aligned} U_t(s) &= \sum_{r=1}^m W(t-s-h_r) A_r \mathcal{X}_{[-h_r, 0]}(s) + \int_{-h}^s W(t-s+\xi) A_I(\xi) d\xi \\ &= \int_{-h}^s W(t-s+\xi) d\eta(\xi), \quad \text{a.e. } s \in I_h \end{aligned}$$

belongs to  $L_p(I_h; B(X))$ . This follows from the Hausdorff-Young inequality. Hence the function

$$(2.7) \quad x(t; g, u) = \begin{cases} W(t)g^0 + \int_{-h}^0 U_t(s)g^1(s) ds + \int_0^t W(t-s)u(s) ds, & t \geq 0 \\ g^1(t) & \text{a.e. } t \in [-h, 0). \end{cases}$$

is well defined and is an element of  $C(\mathbf{R}^+; X) \cap L_p(I_h; X)$ . From (2.4)–(2.7) we can derive the following estimate

$$(2.8) \quad |x(t; g, u)| \leq (M_0 \|g\|_{M_p} + M_1 \|u(\cdot)\|_{p, [0, t]}) \exp(\gamma_0 t), \quad t \geq 0,$$

where  $M_0$  and  $M_1$  are constants depending only on  $p, \eta$  and  $A_0$ .

**Theorem 2.1.** *Let (2.5) be satisfied. Then the function  $x(t) = x(t; g, u)$  in (2.7) is the unique solution of the following functional integral equation:*

$$(2.9) \quad x(t) = \begin{cases} T(t)g^0 + \int_0^t T(t-s) \int_{-h}^0 d\eta(\xi) x(s+\xi) ds + \int_0^t T(t-s)u(s) ds, & t \geq 0 \\ g^1(t) & \text{a.e. } t \in [-h, 0). \end{cases}$$

In the sense of Theorem 2.1 we shall call this  $x(t)$  the mild solution of  $(E)$ . The formula (2.7) is well known as a variation of constants formula for retarded FDE's in  $\mathbf{R}^n$  (cf. Hale [18, Chap. 6]). Since we use the class of mild solutions (2.7) throughout this paper, the condition (2.5) is always assumed. A sufficient condition for the existence of differentiable solution of  $(E)$  is given by the next corollary (for the proof see [34]).

**Corollary 2.1.** *Let  $X$  be reflexive. If  $g=(g^0, g^1)$  and  $u$  satisfy*

$$\begin{aligned} g^1 &\in W_p^{(1)}(I_h; X), \quad g^1(0) = g^0 \in D(A_0), \\ u &\in W_p^{(1)}([0, t]; X) \quad \text{for each } t > 0, \end{aligned}$$

*then the function  $x(t)=x(t; g, u)$  given in (2.7) is a strong solution of  $(E)$ , i.e.,  $x(t)$  satisfies (i)  $x \in C(\mathbf{R}^+; X) \cap W_p^{(1)}([0, t]; X)$  for all  $t > 0$ ; (ii)  $x(t) \in D(A_0)$  for a.e.  $t \geq 0$ ,  $x(t)$  is strongly differentiable and satisfies the equation (2.1); (iii)  $x(0)=g^0$ ,  $x(s)=g^1(s)$  a.e.  $s \in [-h, 0)$ .*

For each  $\lambda \in \mathbf{C}^1$  we define the densely defined closed linear operator  $\Delta(\lambda) = \Delta(\lambda; A_0, \eta)$  by

$$(2.10) \quad \Delta(\lambda) = \lambda I - A_0 - \int_{-h}^0 e^{\lambda s} d\eta(s),$$

where  $I$  denotes the identity operator on  $X$ . The retarded resolvent set  $\rho(A_0, \eta)$  we understand the set of all values  $\lambda$  in  $\mathbf{C}^1$  for which the operator  $\Delta(\lambda)$  has a bounded inverse with dense domain in  $X$ . In this case  $\Delta(\lambda)^{-1}$  is denoted by  $R(\lambda; A_0, \eta)$  and is called the retarded resolvent. The complement of  $\rho(A_0, \eta)$  in the complex plane is called the retarded spectrum and is denoted by  $\sigma(A_0, \eta)$ . The three different types of retarded spectrum can be defined as in the following manner. The continuous retarded spectrum  $\sigma_c(A_0, \eta)$  is the set of values  $\lambda$  for which  $\Delta(\lambda)$  has an unbounded inverse with dense domain in  $X$ . The residual retarded spectrum  $\sigma_r(A_0, \eta)$  is the set of values  $\lambda$  for which  $\Delta(\lambda)$  has an inverse whose domain is not dense in  $X$ . The point retarded spectrum  $\sigma_p(A_0, \eta)$  is the set of values  $\lambda$  for which no inverse of  $\Delta(\lambda)$  exists (cf. Hille and Phillip [31, p. 54], Tanabe [43, Chap. 8]).

We know that the retarded resolvent set  $\rho(A_0, \eta)$  is open in  $\mathbf{C}^1$  and contains right half plane and the retarded resolvent  $R(\lambda; A_0, \eta)$  is holomorphic on  $\rho(A_0, \eta)$ . In fact, we have the following

**Theorem 2.2.** *Let*

$$(2.11) \quad \omega_0 = \inf \{ \alpha : \|W(t)\| \leq M e^{\alpha t}, \quad t > 0 \text{ for some } M > 0 \}.$$

*If  $\text{Re } \lambda > \omega_0$ , then  $\lambda \in \rho(A_0, \eta)$  and the retarded resolvent  $R(\lambda; A_0, \eta)$  is given by the Laplace transform of  $W(t)$ , i.e.,*



$$(2.12) \quad R(\lambda; A_0, \eta) = \int_0^\infty e^{-\lambda t} W(t) dt.$$

Next we give an elementary adjoint theory for  $(E)$  under the assumption that  $X$  is reflexive and  $p \neq \infty$ . Then the adjoint space  $M_p^*$  of  $M_p$  is identified with the product space  $X^* \times L_{p'}(I_h; X^*)$ , where  $1/p + 1/p' = 1$ . Let  $f = (f^0, f^1) \in M_p^*$  and  $(v \in L_{p'}^{loc}(\mathbf{R}^+; X^*))$ . The transposed equation  $(E^T)$  on  $X^*$  is defined by

$$(2.13) \quad \frac{dz(t)}{dt} = A_0^* z(t) + \int_{-h}^0 d\eta^*(s) z(t+s) + v(t) \quad \text{a.e. } t \geq 0$$

$$(2.14) \quad z(0) = f^0, \quad z(s) = f^1(s) \quad \text{a.e. } s \in [-h, 0).$$

Since  $X$  is reflexive, the adjoint operator  $A_0^*$  generates a  $C_0$ -semigroup  $T^*(t)$  on  $X^*$  which is given by the adjoint of  $T(t)$  (see [37]). Hence we can construct the fundamental solution  $W^T(t)$  of  $(E^T)$  as the unique solution of the equation

$$W^T(t) = \begin{cases} T^*(t) + \int_0^t T^*(t-s) \int_{-h}^0 d\eta^*(\xi) W^T(\xi+s) ds, & t \geq 0 \\ 0, & t < 0. \end{cases}$$

We denote by  $W^*(t)$  the adjoint of  $W(t)$ . Then we can show that  $W^T(t) = W^*(t)$ ,  $t \in \mathbf{R}^1$ . This implies that  $W^*(t)$  is strongly continuous on  $\mathbf{R}^+$ . Throughout this paper the condition

$$A_f^*(\cdot) \in L_p(I_h; B(X^*)), \quad 1/p + 1/p' = 1$$

is assumed whenever the transposed equation  $(E^T)$  is in consideration. Thus, the (unique) mild solution  $z(t)$  of  $(E^T)$  exists and is represented by

$$(2.15) \quad z(t) = z(t; f, v) = W^*(t) f^0 + \int_{-h}^0 V_t(s) f^1(s) ds + \int_0^t W^*(t-s) v(s) ds, \quad t \geq 0,$$

where

$$(2.16) \quad V_t(s) = \sum_{r=1}^m W^*(t-s-h_r) A_r^* \chi_{[-h_r, 0]}(s) + \int_{-h}^s W^*(t-s+\xi) A_f^*(\xi) d\xi, \quad \text{a.e. } s \in I_h,$$

For  $\lambda \in \mathbf{C}^1$  define the operator

$$\Delta_T(\lambda) = \Delta(\lambda; A_0^*, \eta^*) = \lambda I - A_0^* - \int_{-h}^0 e^{\lambda s} d\eta^*(s).$$

The retarded resolvent and three kinds of spectrum corresponding to  $\Delta_T(\lambda)$  are defined similarly as for  $\Delta(\lambda)$ .

**Theorem 2.3.** (i)  $\lambda \in \rho(A_0, \eta)$  if and only if  $\bar{\lambda}$  (complex conjugate)  $\in \rho(A_0^*, \eta^*)$  and

$$(2.17) \quad R(\lambda; A_0, \eta)^* = R(\bar{\lambda}; A_0^*, \eta^*).$$

(ii) Both retarded resolvent sets  $\rho(A_0, \eta)$  and  $\rho(A_0^*, \eta^*)$  contain the half plane

$\{\lambda \in \mathbb{C}^1: \operatorname{Re} \lambda > \omega_0\}$ , where  $\omega_0$  is given in (2.11).

$$(iii) \quad R(\lambda; A_0^*, \eta^*) = \int_0^\infty e^{-\lambda t} W^*(t) dt \quad \text{for } \operatorname{Re} \lambda > \omega_0.$$

Here in Theorem 2.3 (i) we remark that the duality pairing  $\langle, \rangle$  between  $X$  and  $X^*$  satisfies

$$(2.18) \quad \langle x, \alpha x^* \rangle = \langle \alpha x, x^* \rangle \quad \text{for } \alpha \in \mathbb{C}^1, (x, x^*) \in X \times X^*.$$

Complete proofs of these results in this section can be found in [34, 35].

### 3. Semigroups associated with functional differential equations

This section is devoted to studying basic properties of semigroups associated with the equations (E) and (E<sup>T</sup>). In what follows we assume that  $X$  is reflexive and  $1 < p < \infty$ .

Let  $x(t; g)$  be the mild solution of (E) with  $u \equiv 0$  and  $g \in M_p$ . The solution operator  $S(t): M_p \rightarrow M_p, t \geq 0$  is defined by

$$(3.1) \quad S(t)g = (x(t; g), x_t(\cdot; g)) \quad \text{for } g \in M_p,$$

where  $x_t(s; g) = x(t+s; g)$  a.e.  $s \in I_h$ . The operator  $S(t)$  is bounded and linear on  $M_p$  by (2.7) and has the following properties (for similar results, see [2,4,5, 6,44,46,48]).

**Proposition 3.1.** (i) *The family of operators  $\{S(t): t \geq 0\}$  is a  $C_0$ -semigroup on  $M_p$ .*

(ii) *If  $T(t)$  is compact for all  $t > 0$ , then  $S(t)$  is compact for  $t > h$ .*

(iii) *The infinitesimal generator  $A$  of  $S(t)$  is given by*

$$(3.2) \quad D(A) = \{g = (g^0, g^1) \in M_p: g^1 \in W_p^{(1)}(I_h; X), g^1(0) = g^0 \in D(A_0)\},$$

$$(3.3) \quad Ag = (A_0 g^0 + \int_{-h}^0 d\eta(s)g^1(s), \frac{dg^1}{ds}(\cdot)) \quad \text{for } g = (g^0, g^1) \in D(A),$$

and for  $g \in D(A)$ ,

$$(3.4) \quad \frac{dS(t)g}{dt} = AS(t)g = S(t)Ag, \quad t > 0.$$

Proof. (i) The semigroup property  $S(t+s) = S(t)S(s), S(0) = I$  is obvious from the definition (3.1). Strong continuity of  $S(t)$  on  $M_p$  follows from that  $x(t; g) \rightarrow g^0$  in  $X$  as  $t \rightarrow 0+$  by (2.9) and that  $x_t(\cdot; g) \rightarrow g^1$  in  $L_p(I_h; X)$  as  $t \rightarrow 0+$  by the absolute continuity of Bochner integrable functions (cf. Ahmed and Teo [1, p. 16]).

(ii) First we introduce the operator  $Q^t: M_p \rightarrow X, t \geq 0$  defined by

$$(3.5) \quad Q^t g = \int_0^t T(t-s)k(s; g)ds, \quad g \in M_p,$$

where

$$(3.6) \quad k(s; g) = \int_{-h}^0 d\eta(\xi)x(s+\xi; g), \quad s \geq 0.$$

Using Hölder inequality and the estimate (2.8), we have

$$(3.7) \quad \|k(\cdot; g)\|_{p, [0, t]} \leq M_2(t)\|g\|_{M_p},$$

where

$$M_2(t) = \left( \sum_{r=1}^m \|A_r\| + \|A_t(\cdot)\|_{p', I_h} \cdot t^{1/p} \right) (1 + M_0 t^{1/p} \exp(\gamma_0 t)).$$

In order to prove the compactness of  $Q^t$  for  $t > 0$  under the compactness of  $T(t)$ ,  $t > 0$ , we define the  $\varepsilon$ -approximation  $Q_\varepsilon^t: M_p \rightarrow X$  of  $Q^t$  for  $\varepsilon \in (0, t]$  by

$$(3.8) \quad Q_\varepsilon^t g = T(\varepsilon) \int_0^{t-\varepsilon} T(t-\varepsilon-s)k(s; g)ds, \quad g \in M_p$$

Since  $T(\varepsilon)$  is compact,  $Q_\varepsilon^t$  is also compact. The compactness of  $Q^t$  follows from

$$(3.9) \quad |(Q_\varepsilon^t - Q^t)g| = \left| \int_{t-\varepsilon}^t T(t-s)k(s; g)ds \right| \leq M_3(t) \cdot \varepsilon^{1/p'} \cdot \|g\|_{M_p},$$

where  $M_3(t) = \left( \sup_{s \in [0, t]} \|T(s)\| \right) M_2(t)$ .

Now let  $t > h$  be fixed and let the operator  $R^t: M_p \rightarrow C([t-h, t]; X)$  be defined by

$$(R^t g)(s) = x(s; g), \quad s \in [t-h, t].$$

Let  $E$  be a bounded set in  $M_p$ . Since  $T(s)$  and  $Q^s$  are compact for  $s > 0$ , from the equation (2.9) it follows that for each  $s \in [t-h, t]$ , the set  $\{(R^t g)(s) \in X: g \in E\}$  is precompact in  $X$ . Next we shall prove that  $\{R^t g; g \in E\}$  is an equicontinuous family of  $C([t-h, t]; X)$ . Let  $0 < a < t-h$ ,  $g \in E$  and  $t-h \leq s' < s \leq t$ . Then we obtain from (3.5) and (3.9) that

$$(3.10) \quad \begin{aligned} & |(R^t g)(s) - (R^t g)(s')| \\ & \leq \|T(s) - T(s')\| \cdot |g^0| + \int_{s'}^s \|T(s-\cdot) - T(s'-\cdot)\| \cdot |k(\tau; g)| d\tau \\ & \quad + \int_0^{s'-a} \|T(s-\tau) - T(s'-\tau)\| \cdot |k(\tau; g)| d\tau \\ & \quad + \int_{s'-a}^{s'} \|T(s-\tau) - T(s'-\tau)\| \cdot |k(\tau; g)| d\tau \end{aligned}$$

$$\begin{aligned} &\leq \|T(s) - T(s')\| \cdot |g^0| + M_3(t) \|g\|_{M_p} (s - s')^{1/p'} \\ &\quad + (\sup \{ \|T(\tau) - T(\tau')\| : \tau, \tau' \in [a, t], |\tau - \tau'| = |s - s'| \}) \\ &\quad \times t^{1/p'} M_3(t) \|g\|_{M_p} + 2M_3(t) \|g\|_{M_p} \cdot a^{1/p'}. \end{aligned}$$

For each fixed  $a > 0$ , it is verified via Hille and Phillips [21, p. 304] that  $T(s)$  is uniformly continuous on  $[a, t]$  in the operator norm topology of  $B(X)$ . Taking  $a > 0$  sufficiently small and applying the uniform continuity to (3.10), we have the desired equi-continuity. Therefore by Royden [38, p. 155],  $R^t$  is compact. Now we introduce the immersion  $I^t: C([t - h, t]; X) \rightarrow M_p$  by  $I^t x(\cdot) = (x(t), x_t(\cdot))$  for  $x \in C([t - h, t]; X)$ . Clearly  $I^t$  is bounded. Since  $S(t)$  can be decomposed as  $S(t) = I^t R^t$  for  $t > h$ ,  $S(t)$  is compact for  $t > h$ .

(iii) We denote by  $\bar{A}$  and  $D(\bar{A})$  the infinitesimal generator of  $S(t)$  and its domain, respectively. Let  $g \in D(\bar{A})$  and

$$(3.11) \quad \bar{A}g = (y^0, y^1).$$

Since the second coordinate of  $S(t)g$  is the  $t$ -shift  $x(t + \cdot; g)$ , it follows immediately that

$$(3.12) \quad x(\cdot; g) = g^1 \in W_p^{(1)}(I_h; X) \quad \text{and} \quad \frac{d^+}{ds} x(\cdot; g) = \dot{g}^1 = y^1 \quad \text{in} \quad L_p(I_h; X),$$

where  $\frac{d^+}{ds}$  denotes the right hand derivative. By redefining on the set of measure

0 we can suppose that  $x(s; g) = g^1(s)$  is absolutely continuous from  $I_h$  to  $X$  (cf. Barbu [3, p. 19, Theorem 2.2]). Since  $x(0; g) = g^0$ , this implies  $g^1(0) = g^0$  and  $x(\cdot; g) \in C([-h, \infty); X)$ . Then the function  $k(s; g)$  in (3.6) is continuous in  $s \geq 0$  and satisfies  $\lim_{t \rightarrow 0^+} k(t; g) = \int_{-h}^0 d\eta(s)g^1(s)$ . So that

$$(3.13) \quad \lim_{t \rightarrow 0^+} \frac{1}{t} \int_0^t T(t-s)k(s; g)ds = \int_{-h}^0 d\eta(s)g^1(s).$$

Applying (2.9) and (3.13) to the first coordinate of (3.11), we obtain that

$$\begin{aligned} (3.14) \quad y^0 &= \lim_{t \rightarrow 0^+} \frac{1}{t} (x(t; g) - g^0) \\ &= \lim_{t \rightarrow 0^+} \frac{1}{t} (T(t)g^0 + \int_0^t T(t-s)k(s; g)ds - g_0) \\ &= \lim_{t \rightarrow 0^+} \frac{1}{t} (T(t)g^0 - g) + \int_{-h}^0 d\eta(s)g^1(s) \quad \text{exists in } X. \end{aligned}$$

Hence  $\lim_{t \rightarrow 0^+} t^{-1}(T(t)g^0 - g^0)$  exists in  $X$ , i.e.,  $g^0 \in D(A_0)$  and  $y^0 = A_0g^0 + \int_{-h}^0 d\eta(s)g^1(s)$ .

This shows

$$D(\bar{A}) \subset D(A) \quad \text{and} \quad Ag = \bar{A}g \quad \text{for} \quad g \in D(\bar{A}).$$

Next we show the reverse inclusion. Let  $g \in D(A)$ . According to Corollary 2.1 we have  $x(\cdot; g) \in W_p^{(1)}([-h, a]; X)$  for any  $a > 0$ , from which (3.13) follows. Combining this with  $g^0 \in D(A_0)$  we see that

$$\lim_{t \rightarrow +0} \frac{1}{t} (x(t; g) - g^0) = A_0 g^0 + \int_{-h}^0 d\eta(s) g^1(s).$$

Noting

$$\begin{aligned} (3.15) \quad & \frac{1}{t} (x_t(\xi; g) - g^1(\xi)) - \dot{g}^1(\xi) \\ &= \frac{1}{t} (x(t+\xi; g) - x(\xi; g)) - \dot{x}(\xi; g) = \frac{1}{t} \int_0^t (\dot{x}(s+\xi; g) - \dot{x}(\xi; g)) ds, \end{aligned}$$

for  $\xi \in [-h, 0]$  we obtain with the aid of Hölder inequality that

$$(3.16) \quad \left\| \frac{1}{t} (x_t(\cdot; g) - g^1) - \dot{g}^1 \right\|_{p, I_h} \leq \frac{1}{t} \int_0^t \left[ \int_{-h}^0 |\dot{x}(s+\xi; g) - \dot{x}(\xi; g)|^p d\xi \right] ds.$$

This implies that  $\lim_{t \rightarrow 0^+} t^{-1} (x_t(\cdot; g) - g^1)$  exists in  $L_p(I_h; X)$  and equals  $\dot{g}^1$ . Thus, we prove  $D(A) \subset D(\tilde{A})$  and  $Ag = \tilde{A}g$  for  $g \in D(A)$ , and hence (3.2), (3.3) are shown. The remaining equality (3.4) is obvious.

Concerning the transposed equation ( $E^T$ ) we define the semigroup  $S_T(t)$  on  $M_p^*$  in an analogous manner. Thus we have:

**Proposition 3.2.** (i) *The family of operators  $\{S_T(t): t \geq 0\}$  is a  $C_0$ -semigroup on  $M_p^*$ .*

(ii) *If  $T(t)$  is compact for  $t > 0$ , then  $S_T(t)$  is compact for  $t > h$ .*

(iii) *The infinitesimal generator  $A_T$  of  $S_T(t)$  is given by*

$$\begin{aligned} D(A_T) &= \{f = (f^0, f^1) \in M_p^*: f^1 \in W_{p'}^{(1)}(I_h; X^*), f^1(0) = f^0 \in D(A_0^*)\}, \\ A_T f &= (A_0^* f^0 + \int_{-h}^0 d\eta^*(s) f^1(s), \frac{df^1}{ds}(\cdot)) \quad \text{for } f = (f^0, f^1) \in D(A_T), \end{aligned}$$

where  $1/p + 1/p' = 1$ .

Since  $M_p$  is reflexive, we know that the adjoint  $S^*(t)$  of  $S(t)$  generates a  $C_0$ -semigroup on  $M_p^*$ . Probably it was Vinter [46] who first characterized the infinitesimal generator of the semigroup  $S^*(t)$  in the case  $X = \mathbf{R}^n$  and  $p = 2$ . His article seems hardly to be available, however, we shall give a complete proof of the result in our Banach space case.

**Proposition 3.3.** *The infinitesimal generator  $A^*$  of  $S^*(t)$  is given by*

$$(3.17) \quad D(A^*) = \{f = (f^0, f^1) \in M_p^*: w(f) \in W_{p'}^{(1)}(I_h; X^*), w(f)(-h) = 0, f^0 \in D(A_0^*)\}$$

$$(3.18) \quad A^*f = (A_0^*f^0 + f^1(0), w(f)) \quad \text{for } f = (f^0, f^1) \in D(A^*),$$

where

$$(3.19) \quad w(f)(s) = \int_{-h}^s d\eta^*(s)f^0 - f^1(s), \quad s \in I_h.$$

Proof. Note first that the infinitesimal generator of  $S^*(t)$  is given by the adjoint  $A^*$  of  $A$ . Let  $(g^0, g^1) \in D(A)$  and  $(f^0, f^1) \in M_p^*$ . Assume that there exists a  $(k^0, k^1) \in M_p^*$  such that for all  $(g^0, g^1) \in D(A)$ ,

$$\langle A(g^0, g^1), (f^0, f^1) \rangle_{M_p} = \langle (g^0, g^1), (k^0, k^1) \rangle_{M_p}$$

or, equivalently by Proposition 3.1,

$$(3.20) \quad \langle A_0g^0 + \int_{-h}^0 d\eta(s)g^1(s), f^0 \rangle + \int_{-h}^0 \langle \dot{g}^1(s), f^1(s) \rangle ds = \langle g^0, k^0 \rangle + \int_{-h}^0 \langle g^1(s), k^1(s) \rangle ds$$

Set  $M(s) = \int_{-h}^s k^1(\xi)d\xi, s \in I_h$ .  $M(-h) = 0$  is evident. It is easy to see that, by using integration by parts,

$$(3.21) \quad \int_{-h}^0 \langle g^1(s), k^1(s) \rangle ds = \langle g^1(0), M(0) \rangle - \int_{-h}^0 \langle \dot{g}^1(s), M(s) \rangle ds.$$

Next we set  $h_0 = 0$  and

$$N(s) = \int_{-h}^s d\eta^*(s)f^0 = \sum_{r=1}^m A_r^* \chi_{[-h_r, 0]}(s)f^0 + \int_{-h}^s A_1^*(\xi)f^0 d\xi.$$

Again, using integration by parts on each  $[-h_r, -h_{r-1}]$ ,  $r = 1, \dots, m$ , it is not difficult to show that

$$(3.22) \quad \int_{-1}^0 \langle \dot{g}^1(s), N(s) \rangle ds = \langle g^0, N(0) \rangle - \langle \int_{-h}^0 d\eta(s)g^1(s), f^0 \rangle$$

Then by (3.20)–(3.22), we see for  $(g^0, g^1) \in D(A)$ ,

$$(3.23) \quad \int_{-h}^0 \langle \dot{g}^1(s), f^1(s) - N(s) + M(s) \rangle ds + \langle A_0g^0, f^0 \rangle = \langle g^0, k^0 - N(0) + M(0) \rangle.$$

For  $g^0 \in D(A_0)$  and  $g^1(s) \equiv g^0$ , it is obvious that  $(g^0, g^1) \in D(A)$  and  $\dot{g} = 0$ . Hence applying such  $(g^0, g^1)$  to (3.23), we have

$$(3.24) \quad \langle A_0g^0, f^0 \rangle = \langle g^0, k^0 - N(0) + M(0) \rangle \quad \text{for all } g^0 \in D(A_0).$$

This proves that  $f^0 \in D(A_0^*)$  and

$$(3.25) \quad k^0 = A_0^*f^0 + \int_{-h}^0 d\eta^*(s)f^0 - \int_{-h}^0 k^1(s)ds.$$

Since  $\{g^1: (g^0, g^1) \in D(A)\}$  is dense in  $L_p(I_h; X)$ , from (3.23) and (3.24) it follows that

$$(3.26) \quad f^1(s) - N(s) + M(s) = 0 \quad \text{a.e. } s \in I_h.$$

If we put  $w(f)(s) = N(s) - f^1(s)$ ,  $s \in I_h$ , then by (3.26)  $w(f)$  satisfies

$$(3.27) \quad w(f) \in W_p^1(I_h; X^*), \quad \dot{w}(f) = \dot{M} = k^1 \quad \text{in } L_{p'}(I_h; X^*)$$

and

$$(3.28) \quad w(f)(-h) = 0, \quad w(f)(0) = \int_{-h}^0 k^1(s) ds = \int_{-h}^0 d\gamma^*(s) f^0 - f^1(0).$$

Therefore, by (3.25), (3.27) and (3.28) we conclude that  $D(A^*)$  is given by (3.17) and  $A^*f, f \in D(A^*)$  is represented by  $A^*f = (k^0, k^1) = (A_0^*f^0 + f^1(0), \dot{w}(f))$ , which is (3.18). Conversely it is not difficult to show that any element of the right hand side of (3.17) belongs to  $D(A^*)$ . Thus the proof is complete.

#### 4. Structural operators $F, G$ and their adjoint operators

In this section we extend the structural operator  $F$  and  $G_t$  introduced in [4] for the case  $X = \mathbb{R}^n$  to our Banach space case and study their basic properties including the decomposition formula as well as their adjoint operators.

Define the operator  $F_1: L_p(I_h; X) \rightarrow L_p(I_h; X)$  by

$$(4.1) \quad [F_1 g^1](s) = \int_{-h}^s d\gamma(\xi) g^1(\xi - s) \\ = \sum_{r=1}^m A_r \chi_{[-h_r, 0]}(s) g^1(-h_r - s) + \int_{-h}^s A_I(\xi) g^1(\xi - s) d\xi \quad \text{a.e. } s \in I_h.$$

By direct calculations using Hölder inequality it is verified that  $F_1$  is into, linear and bounded.

First we give an equivalent representation formula of the mild solution  $x(t; g)$  to (2.7) in terms of  $W(t)$  and  $F_1$ , which is given by another complicated form in [4, p. 902]. The following one is explicit.

**Lemma 4.1.** *The mild solution  $x(t; g)$  is represented by*

$$(4.2) \quad x(t; g) = W(t)g^0 + \int_{-h}^0 W(t+s) [F_1 g^1](s) ds, \quad t \geq 0.$$

*Proof.* In view of (2.7) we are left to prove the equality

$$(4.3) \quad \int_{-h}^0 U_t(s) g^1(s) ds = \int_{-h}^0 W(t+s) [F_1 g^1](s) ds.$$

With the aid of suitable changes of variables and Fubini's theorem we obtain

$$\begin{aligned} \int_{-h}^0 U_i(s)g^1(s)ds &= \sum_{r=1}^m \int_{-h_r}^0 W(t-s-h_r)A_r g^1(s)ds \\ &\quad + \int_{-h}^0 \left( \int_{-h}^s W(t-s+\xi)A_I(\xi)d\xi \right) g^1(s)ds \\ &= \int_{-h}^0 W(t+s) \left\{ \sum_{r=1}^m A_r \mathcal{X}_{[-h_r, 0]}(s)g^1(-h_r-s) \right\} ds \\ &\quad + \int_{-h}^0 W(t+s) \left\{ \int_{-h}^s A_I(\xi)g^1(\xi-s)d\xi \right\} ds \\ &= \int_{-h}^0 W(t+s)[F_1 g^1](s)ds . \end{aligned}$$

The first structural operator  $F: M_p \rightarrow M_p$  is defined by  $F = \begin{bmatrix} I & O \\ O & F_1 \end{bmatrix}$ , i.e.,

$$(4.4) \quad [Fg]^0 = g^0, \quad [Fg]^1 = F_1 g^1 \quad \text{for } g = (g^0, g^1) \in M_p .$$

By (2.7), (4.4) and Lemma 4.1, we have

$$(4.5) \quad x(t+s; g) = \begin{cases} W(t+s)g^0 + \int_{-h}^0 W(t+s+\xi)[Fg]^1(\xi)d\xi, & t+s \geq 0 \\ g^1(t+s), & t+s < 0 . \end{cases}$$

The equality (4.5) suggests us to introduce the operator  $G_t: M_p \rightarrow M_p, t \geq 0$  defined by

$$(4.6) \quad [G_t g]^1(s) = W(t+s)g^0 + \int_{-h}^0 W(t+s+\xi)g^1(\xi)d\xi, \quad s \in I_h ,$$

$$(4.7) \quad [G_t g]^0 = [G_t g^1](0), \quad g = (g^0, g^1) \in M_p .$$

Clearly  $G_t$  is linear and bounded. Notice that the right hand side of (4.6) vanishes if  $t+s < 0$ . Especially we define the second structural operator  $G: M_p \rightarrow M_p$  by

$$(4.8) \quad G = G_h .$$

We remark here that  $G_t g \in C(I_h; X)$  for  $t \geq h$  and  $g \in M_p$ .

The following proposition is obvious from (4.5) and the definitions of  $F, G_t, G$  and  $\kappa(t)$ .

**Proposition 4.1.** *The semigroup  $S(t)$  is represented by*

$$(4.9) \quad S(t) = G_t F + \kappa(t), \quad t \geq 0 ,$$

where  $\kappa(t): M_p \rightarrow M_p$  is given by

$$(4.10) \quad [\kappa(t)g]^0 = 0, \quad [\kappa(t)g]^1(s) = g^1(t+s)\mathcal{X}_{[-h, -t]}(s) \quad \text{a.e. } s \in I_h .$$

In particular,  $S(h)$  is decomposed as



$$(4.11) \quad S(h) = GF.$$

To obtain a similar representation formula for the transposed semigroup  $S_T(t)$ , we have to compute the adjoints of  $G_t$  and  $F$  (cf. (2.15), (2.16)).

The following proposition can be established by a direct calculation.

**Proposition 4.2.** *The adjoint  $F^*: M_p^* \rightarrow M_p^*$  of  $F$  is given by*

$$F^* = \begin{bmatrix} I & O \\ O & F_1^* \end{bmatrix},$$

where  $F_1^*: L_p(I_h; X^*) \rightarrow L_p(I_h; X^*)$  denotes the adjoint of  $F_1$  and is represented by

$$(4.12) \quad [F_1^* f^1](s) = \int_{-h}^s d\eta^*(\xi) f^1(\xi - s) \\ = \sum_{r=1}^m A_r^* \chi_{[-h_r, 0]}(s) f^1(-h_r - s) + \int_{-h}^s A_1^*(\xi) f^1(\xi - s) d\xi \quad \text{a.e. } s \in I_h.$$

The following proposition is also easily proved.

**Proposition 4.3.** *The adjoint  $G_t^*: M_p^* \rightarrow M_p^*$  of  $G_t$ ,  $t \geq 0$  is represented by*

$$\begin{cases} [G_t^* f]^1(s) = W^*(t+s) f^0 + \int_{-h}^0 W^*(t+s+\xi) f^1(\xi) d\xi, & s \in I_h, \\ [G_t^* f]^0 = [G_t^* f]^1(0), & f = (f^0, f^1) \in M_p^*. \end{cases}$$

Consider the transposed equation  $(E^T)$ . By (2.15) and Proposition 4.2, we see that the mild solution  $z(t; f) = z(t; f, 0)$  of  $(E^T)$  is written as

$$z(t; f) = W^*(t) f^0 + \int_{-h}^0 W^*(t+s) [F^* f]^1(s) ds, \quad t \geq 0.$$

Hence by Proposition 4.3, we obtain the following

**Proposition 4.4.** *The semigroup  $S_T(t)$  is represented by*

$$S_T(t) = G_t^* F^* + \kappa(t), \quad t \geq 0,$$

where  $\kappa(t): M_p^* \rightarrow M_p^*$  is same as given in (4.10). In particular,

$$(4.13) \quad S_T(h) = G^* F^*.$$

We can verify by standard manipulation involving the pairing  $\langle, \rangle_{M_p}$  that the adjoint  $\kappa^*(t): M_p^* \rightarrow M_p^*$  of  $\kappa(t)$  in (4.10) is given by

$$(4.14) \quad [\kappa^*(t) f]^0 = 0, \quad [\kappa^*(t) f]^1(s) = \chi_{[0, s+h]}(t) f^1(s-t), \quad f \in M_p^*.$$

Since the same operator as in (4.14) can be defined on  $M_p$ , we denote this operator

rator by the same symbol  $\kappa^*(t)$ . Then taking adjoints of  $S(t)$  and  $S_T(t)$ , we have the following result.

**Corollary 4.1.** *The adjoint semigroups  $S^*(t)$  and  $S_T^*(t)$  are represented by*

$$(4.15) \quad S^*(t) = F^*G_t^* + \kappa^*(t), \quad S_T^*(t) = FG_t + \kappa^*(t), \quad t \geq 0,$$

respectively. In particular,

$$(4.16) \quad S^*(h) = F^*G^*, \quad S_T^*(h) = FG.$$

It is well known that the adjoint semigroup  $S^*(t)$  plays an important role in the study of linear quadratic optimal control problem associated with FDE's including their numerical computations (cf. [11, 12, 14, 17, 47]). The structure of  $S^*(t)$  is not straightforward compared with  $S_T(t)$ , since a functional differential equation which realizes  $S^*(t)$  has not been unknown. The advantage of the use of transposed semigroup  $S_T(t)$  depends on this fact and that  $S^*(t)$  and  $S_T(t)$  are connected by the operators  $F^*$  and  $G^*$  in an appropriate way (see Theorems 4.1, 4.2 below).

A somewhat simple property of  $G$  and  $G^*$  is the following

**Proposition 4.5.** (i)  $C1(\text{Im } G) = M_p, \text{ Ker } G = \{0\}$ ;

(ii)  $C1(\text{Im } G^*) = M_p^*, \text{ Ker } G^* = \{0\}$ .

Proof. First we shall show  $\text{Ker } G = \{0\}$ . Assume  $Gg = 0$  in  $M_p$ . Then by (4.6) and (4.8),  $0 = [Gg]^1(-h) = W(0)g^0 = g^0$ . Using this and changing variables  $\xi \rightarrow -\xi$  and  $h + s \rightarrow s$  in (4.6), we have

$$(4.17) \quad [Gg]^1(s-h) = \int_0^s W(s-\xi)g^1(-\xi)d\xi = 0 \quad \text{for all } s \in [0, h].$$

Now we can use a convolution type result on the fundamental solution in Nakagiri [34, Lemma 5.1] to obtain from (4.17) that  $g^1(-\xi) = 0$  a.e.  $\xi \in [0, h]$ , i.e.,  $g^1 = 0$  in  $L_p(I_h; X)$ . Hence  $g = (g^0, g^1) = 0$  in  $M_p$ , which proves  $\text{Ker } G = \{0\}$ . Similarly, by Proposition 4.3  $\text{Ker } G^* = \{0\}$  holds. Since  $M_p$  is reflexive, it follows from the duality theorem (cf. Kato [25, p. 243], Tanabe [43, Chapter III]) that  $\text{Ker } G = \{0\}$  (resp.  $\text{Ker } G^* = \{0\}$ ) is equivalent to  $C1(\text{Im } G^*) = M_p^*$  (resp.  $C1(\text{Im } G) = M_p$ ). This proves (i) and (ii).

In the special case where  $A_0$  is bounded, we have the following sharper result for  $G$  than Proposition 4.5.

**Proposition 4.6.** *Let  $A_0$  be bounded. Then*

- (i)  $\text{Im } G = D(A)$  and  $G: M_p \rightarrow D(A)$  is bijective;
- (ii)  $G^{-1}: D(A) \rightarrow M_p$  is given by

$$(4.18) \quad \begin{cases} [G^{-1}g]^1(s) = \dot{g}^1(-s-h) - A_0g^1(-s-h) - \int_s^0 d\eta(\xi)g^1(\xi-s-h) & \text{a.e. } s \in I_h \\ [G^{-1}g]^0 = g^1(-h), & g \in D(A); \end{cases}$$

(iii)

$$(4.19) \quad \text{Im } S(t) \subset D(A) \quad \text{for } t \geq h.$$

Proof. Since  $A_0$  is bounded, we see from Delfour [13, Theorems 1.1, 1.2] that the function

$$(4.20) \quad W(t)k^0 + \int_0^t W(t-s)k^1(s)ds, \quad t \in [0, h],$$

where  $k^0 \in X, k^1 \in L_p([0, h]; X)$ , gives a unique strong solution of (E) with  $g^0 = k^0, g^1 = 0$  and  $u = k^1$ . For  $g = (g^0, g^1) \in M_p$  the function  $y(t) = [Gg]^1(t-h), t \in [0, h]$  is given by (4.20) with  $k^0 = g^0$  and  $k^1(s) = g^1(-s)$ . Then  $[Gg]^1(\cdot) = y(\cdot + h) \in W_p^{(1)}(I_h; X)$ , and hence by  $[Gg]^1(0) = [Gg]^0, Gg \in D(A)$  i.e.,  $\text{Im } G \subset D(A)$ . To prove the reverse inclusion let  $g = (g^1(0), g^1) \in D(A)$  and define  $\psi = (\psi^0, \psi^1)$  by the right hand side of (4.18). It is clear that  $\psi \in M_p$ . Put  $y(t) = g^1(t-h), t \in [0, h]$  and  $y(0) = g^1(-h), y(s) = 0$  a.e.  $s \in I_h$ . Then from (4.18) it follows immediately that  $y(t)$  is a strong solution of (E) with  $k^0 = \psi^0$  and  $k^1(s) = \psi^1(-s)$ . Hence by uniqueness,  $y(t) = [G\psi]^1(t-h) = g^1(t-h), t \in [0, h]$  and especially  $y(h) = [G\psi]^1(0) = [G\psi]^0 = g^1(0)$ . So that  $G\psi = g$ , or  $\psi = G^{-1}g$ . This shows (i) and (ii) simultaneously. Since  $S(t) = S(h)S(t-h) = GFS(t-h)$  by (4.11) and  $\text{Im } G = D(A)$ , we have (iii). In other words,  $S(t)$  is differentiable for  $t \geq h$ .

We note that  $S^*(t)$  also satisfies  $\text{Im } S^*(t) \subset D(A^*)$  for  $t \geq h$ . Next we establish a simple and fundamental relation between  $S(t)$  and  $G_t$  which can not be found in any literatures studying state space theory for (E).

**Proposition 4.7.** (i)  $S(t)G = G_{t+h}, S_T(t)G^* = G_{t+h}^*, t \geq 0;$

(ii)  $G^*S^*(t) = G_{t+h}^*, GS_T^*(t) = G_{t+h}, t \geq 0.$

Proof. (i) With the aid of (2.7), (4.6), (4.8), we have for  $g = (g^0, g^1) \in M_p,$

$$(4.21) \quad \begin{aligned} [S(t)Gg]^1(s) &= (W(t+s)W(h) + \int_{-h}^0 U_{t+s}(\xi)W(h+\xi)d\xi)g^0 \\ &\quad + \int_{-h}^0 W(t+s)W(h+\xi)g^1(\xi)d\xi \\ &\quad + \int_{-h}^0 \int_{-h}^0 U_{t+s}(\xi)W(h+\xi+\alpha)g^1(\alpha)d\alpha d\xi = I_1 + I_2 + I_3. \end{aligned}$$

As is easily seen

$$(4.22) \quad I_2 + I_3 = \int_{-h}^0 (W(t+s)W(h+\xi) + \int_{-h}^0 U_{t+s}(\alpha)W(h+\xi+\alpha)d\alpha)g^1(\xi)d\xi.$$

We now recall the following quasi-semigroup property of  $W(t)$  given in [35,

Eq. (4.9)];

$$(4.23) \quad W(t_1+t_2) = W(t_1)W(t_2) + \int_{-h}^0 U_{t_1}(\xi)W(t_2+\xi)d\xi, \quad t_1, t_2 \geq 0.$$

Applying (4.23) to  $I_1$  in (4.21) and the integrand in (4.22), we obtain that

$$(4.24) \quad [S(t)Gg]^1(s) = W(t+s+h)g^0 + \int_{-h}^0 W(t+s+h+\xi)g^1(\xi)d\xi = [G_{t+h}g]^1(s).$$

Substituting  $s=0$  in (4.24), we have  $[S(t)Gg]^0 = [G_{t+h}g]^0$ . Therefore,  $S(t)G = G_{t+h}$  is proved. Similarly  $S_T(t)G^* = G_{t+h}^*$  is true.

(ii) Take adjoints of the equalities in (i).

We are now ready to give the main theorem which is one of the key results in the state space theory. A similar result for  $X = \mathbf{R}^n$  is already proved by Manitius [29, Theorem 3.3], however his proof is much complicated and can not be carried to our Banach space case ( $W(t)$  is not differentiable!). Here we shall give a very simple proof based on Proposition 4.7.

**Theorem 4.1.** (i)

$$(4.25) \quad S(t)G = GS^*(t), \quad G^*S^*(t) = S_T(t)G^*, \quad t \geq 0;$$

(ii)

$$(4.26) \quad GD(A^*) \subset D(A) \quad \text{and} \quad AG = GA^* \quad \text{on} \quad D(A^*);$$

$$(4.27) \quad G^*D(A^*) \subset D(A_T) \quad \text{and} \quad G^*A^* = A_TG^* \quad \text{on} \quad D(A^*).$$

Proof. The part (i) is a direct consequence from Proposition 4.7 and the part (ii) follows from (i) and the definition of infinitesimal generator.

The next is the second key result related to  $F$ , which is first proved by Bernier and Manitius [4, Theorem 5.4] and later by Delfour and Manitius [15, Theorem 3.1] for more general measure  $\eta$ . Compare their proofs and our simple proof.

**Theorem 4.2.** (i)

$$(4.28) \quad FS(t) = S^*(t)F, \quad S^*(t)F^* = F^*S_T(t), \quad t \geq 0;$$

(ii)

$$(4.29) \quad FD(A) \subset D(A^*) \quad \text{and} \quad FA = A^*F \quad \text{on} \quad D(A);$$

$$(4.30) \quad F^*D(A_T) \subset D(A^*) \quad \text{and} \quad A^*F^* = F^*A_T \quad \text{on} \quad D(A_T).$$

Proof. Since (ii) follows from (i), we prove only (i). By (4.25) and (4.11),

$$(4.31) \quad \begin{aligned} G(S^*(t)F) &= (GS^*(t))F = S(t)GF = S(t)S(h) = S(h)S(t) \\ &= GFS(t) = G(FS(t)), \quad t \geq 0. \end{aligned}$$

Since  $\text{Ker } G = \{0\}$ , it follows from (4.31) that  $S^*(t)F = FS(t)$ ,  $t \geq 0$ . The second equality in (i) is proved analogously.

**Corollary 4.2.** (i)

$$(4.32) \quad \text{Ker } F = \{g \in M_p : x(t; g) = 0 \text{ for } t \in [0, h]\};$$

(ii)

$$(4.33) \quad \text{Ker } F^* = \{f \in M_p^* : z(t; f) = 0 \text{ for } t \in [0, h]\}.$$

Proof. Since  $S(t)$  is a semigroup defined by (3.1) and  $(E)$  is autonomous, we see easily that

$$(4.34) \quad \begin{aligned} S(h)g = 0 & \text{ if and only if } S(t+h)g = 0 \text{ for all } t \geq 0, \\ & \text{if and only if } x(t; g) = 0 \text{ for all } t \geq 0, \\ & \text{if and only if } x(t; g) = 0 \text{ for all } t \in [0, h]. \end{aligned}$$

From (4.11) and  $\text{Ker } G = \{0\}$ , we have  $\text{Ker } F = \text{Ker } S(h)$ . Hence (4.34) implies (4.32). Similarly (4.33) is proved by (4.13) and Proposition 4.5 (ii).

Lastly in this section we introduce a bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  between  $M_p$  and  $M_p^*$  defined by

$$(4.35) \quad \langle\langle g, f \rangle\rangle = \langle Fg, f \rangle_{M_p} = \langle g, F^*f \rangle_{M_p}.$$

The form  $\langle\langle \cdot, \cdot \rangle\rangle$  is considered a time reversing one of the Hale's bilinear form (see [18, p. 173]) and appears in the representation of basis for generalized eigenspaces associated with  $(E)$  (which will be given in Section 8 below). The following corollary is obvious from Theorem 4.2 and the definition (4.35).

**Corollary 4.3.** (i)  $\langle\langle Ag, f \rangle\rangle = \langle\langle g, A_T f \rangle\rangle$ ,  $(g, f) \in D(A) \times D(A_T)$ .

(ii)  $\langle\langle S(t)g, f \rangle\rangle = \langle\langle g, S_T(t)f \rangle\rangle$ ,  $t \geq 0$ ,  $(g, f) \in M_p \times M_p^*$ .

**5. Characterizations of Ker F and Im F**

In this section we shall give a number of necessary and/or sufficient conditions for  $\text{Ker } F = \{0\}$ ,  $\text{Im } F = M_p$  and  $C1(\text{Im } F) = M_p$  in terms of the coefficient operators appearing in the measure  $\eta$ .

By definition it is clear that

$$(5.1) \quad \text{Ker } F = \{0\} \times \text{Ker } F_1, \quad \text{Im } F = X \times \text{Im } F_1.$$

Further, we know by the duality theorem that

$$(\text{Im } F_1)^\perp = (C1(\text{Im } F_1))^\perp = \text{Ker } F_1^*.$$

Since  $F_1^*$  is an operator of the same type as  $F_1$  (Proposition 4.2), we mainly

investigate the structure of  $\text{Ker } F_1$ . From the definition (4.1) of  $F_1$ , we see that the condition  $g^1 \in \text{Ker } F_1$  is equivalent to that  $g^1$  satisfies

$$(5.2) \quad \sum_{r=1}^m A_r \mathcal{X}_{[-h_r, 0]}(s) g^1(-h_r - s) + \int_{-h}^s A_I(\xi) g^1(\xi - s) d\xi = 0 \quad \text{a.e. } s \in I_h.$$

The equation (5.2) can be written by the following homogeneous Volterra integral equation with delays:

$$(5.3) \quad A_m \psi(t) + \sum_{r=1}^{m-1} A_r \mathcal{X}_{[\tau_r, h]}(t) \psi(t - \tau_r) + \int_0^t A_I(\xi - h) \psi(t - \xi) d\xi = 0$$

a.e.  $t \in [0, h]$

where  $\psi(t) = g^1(-t)$ ,  $t \in [0, h]$  and  $\tau_r = h - h_r > 0$ ,  $r = 1, \dots, m - 1$ .

Hence by the first equality in (5.1),  $\text{Ker } F = \{0\}$  is equivalent to that the equation (5.3) admits a unique trivial solution  $\psi(t) = 0$  a.e.  $t \in [0, h]$ . In order to give conditions for  $\text{Ker } F = \{0\}$  we introduce the following null space  $N(A_I; \alpha)$ ,  $\alpha \in I_h$  associated with the kernel  $A_I(\xi)$ :

$$N(A_I; \alpha) = \{x \in X: A_I(\xi)x = 0 \quad \text{for a.e. } \xi \in [-h, \alpha]\}.$$

**Proposition 5.1.** *A necessary condition for  $\text{Ker } F = \{0\}$  is*

$$(5.4) \quad \text{Ker } A_m \cap N(A_I; \alpha) = \{0\} \quad \text{for each } \alpha \in (-h, -h_{m-1}].$$

Proof. Suppose (5.4) does not hold. Then there exist  $\alpha_0 \in (-h, -h_{m-1}]$  and  $x_0 \neq 0$  such that  $x_0 \in \text{Ker } A_m \cap N(A_I; \alpha_0)$ , i.e.,

$$(5.5) \quad A_m x_0 = 0 \quad \text{and} \quad A_I(\xi)x_0 = 0 \quad \text{a.e. } \xi \in [-h, \alpha_0].$$

Define

$$(5.6) \quad \psi(t) = \begin{cases} 0, & t \in [0, -\alpha_0] \\ x_0, & t \in (-\alpha, h_0], \end{cases}$$

where  $\psi(\cdot) \neq 0$  in  $L_p([0, h]; X)$ . Making use of (5.5) we can verify straightforwardly that  $\psi(t)$  in (5.6) satisfies (5.3). Hence  $\psi(-\cdot) \in \text{Ker } F_1$ , so that  $\text{Ker } F \neq \{0\}$ . This proves the proposition.

**Theorem 5.1.** *Assume that  $A_I(s) = 0$  in a neighbourhood of  $-h$ . A necessary and sufficient condition for  $\text{Ker } F = \{0\}$ , or equivalently  $C1(\text{Im } F^*) = M_b^*$ , is*

$$(5.7) \quad \text{Ker } A_m = \{0\}.$$

Proof. Since the condition (5.7) is necessary by Proposition 5.1 and assumption, it suffices to prove that  $\text{Ker } A_m = \{0\}$  implies  $\text{Ker } F_1 = \{0\}$ . Let  $g^1 \in \text{Ker } F_1$  and  $\psi$  be given in (5.3). Suppose  $A_I(s) = 0$  a.e.  $s \in [-h, -h + \tau]$  for some  $\tau \in (0, h - h_{m-1}]$  by assumption. Then by (5.3),

$$(5.8) \quad A_m \psi(t) + \sum_{r=1}^{m-1} A_r \mathcal{X}_{[\tau_r, h]}(t) \psi(t - \tau_r) = 0$$

for a.e.  $t \in [0, \tau]$ ; in particular  $A_m \psi(t) = 0$  a.e.  $t \in [0, \min(\tau, \tau_{m-1})]$ . So that  $\psi(t) = 0$  a.e.  $t \in [0, \min(\tau, \tau_{m-1})]$  by (5.7). Using this, via step by step argument, we obtain from (5.8) that  $\psi(t) = 0$  a.e.  $t \in [0, \tau]$ . Let  $k \geq 1$  and suppose  $\psi(t) = 0$  a.e.  $t \in [0, k\tau]$ . Then for  $t \in [k\tau, (k+1)\tau]$ , we have

$$\int_0^t A_I(\xi - h) \psi(t - \xi) d\xi = \int_0^{t-k\tau} A_I(\xi - h) \psi(t - \xi) d\xi = 0.$$

Thus,  $\psi(t)$  satisfies (5.8) for a.e.  $t \in [0, (k+1)\tau]$ . Consequently we have  $\psi(t) = 0$  a.e.  $t \in [0, (k+1)\tau]$  similarly as above. Then by mathematical induction  $\psi = 0$ , or  $g^1 = 0$  in  $L_p(I_k; X)$  follows. This shows  $\text{Ker } F = \{0\}$ .

**Proposition 5.2.** *If  $0 \in \rho(A_m)$ , then*

$$(5.9) \quad \text{Im } F = M_p, \quad \text{Im } F^* = M_p^*.$$

Proof. Since  $0 \in \rho(A_m)$ , the inverse  $A_m^{-1}$  exists and is bounded. Let  $\phi \in L_p([0, h]; X)$  be given. Consider the following inhomogeneous Volterra integral equation with delays

$$(5.10) \quad \psi(t) + \sum_{r=1}^{m-1} C_r \mathcal{X}_{[\tau_r, h]}(t) \psi(t - \tau_r) + \int_0^t C_I(\xi - h) \psi(t - \xi) d\xi = A_m^{-1} \phi(t),$$

where  $C_r = A_m^{-1} A_r \in B(X)$ ,  $r = 1, \dots, m-1$  and  $C_I(\cdot) = A_m^{-1} A_I(\cdot) \in L_p(I_h; B(X))$ . For  $t \in [0, \tau_{m-1}]$  the equation (5.10) becomes a Volterra integral equation

$$(5.11) \quad \psi(t) + \int_0^t C_I(t - h - \xi) \psi(\xi) d\xi = A_m^{-1} \phi(t) \quad \text{a.e. } t \in [0, \tau_{m-1}].$$

Since the term  $A_m^{-1} \phi(t)$  belongs to  $L_p([0, \tau_{m-1}]; X)$ , the equation (5.11) admits a unique solution  $\psi \in L_p([0, \tau_{m-1}]; X)$ . This can be proved in the usual manner using the contraction mapping principle in  $L_p$ -space (see e.g. Miller [32] or Hönig [22]). Then (5.10) is solvable on  $[0, \tau_{m-1}]$ . Suppose that  $\psi$  solves (5.1) a.e. in  $[0, k\tau_{m-1}]$ ,  $k \geq 1$ . Then for a.e.  $t \in [k\tau_{m-1}, (k+1)\tau_{m-1}]$ , the equation (5.10) is written by the equivalent form as

$$(5.12) \quad \psi(t) + \int_{k\tau_{m-1}}^t C_I(t - h - \xi) \psi(\xi) d\xi \\ = A_m^{-1} \phi(t) - \int_0^{k\tau_{m-1}} C_I(t - h - \xi) \psi(\xi) d\xi - \sum_{r=1}^{m-1} C_r \mathcal{X}_{[\tau_r, h]}(t) \psi(t - \tau_r).$$

Because  $t - \tau_r \leq t - \tau_{m-1} \leq k\tau_{m-1}$ ,  $r = 1, \dots, m-1$ , the last term in the righthand side of (5.12) is a known function, and hence the right hand side denotes a known function in  $L_p([k\tau_{m-1}, (k+1)\tau_{m-1}]; X)$ . We then have that the equation (5.12)

is a Volterra integral equation which can be solved in the space  $L_p([k\tau_{m-1}, (k+1)\tau_{m-1}]; X)$ . This concludes that (5.10) is solvable on  $[0, (k+1)\tau_{m-1}]$ . Hence by induction, (5.10) is solvable on whole  $[0, h]$ . By a change of variables  $t \rightarrow -s$  and an application of  $A_m$  to (5.10), we derive  $\text{Im } F_1 = L_p(I_h; X)$ , and this implies  $\text{Im } F = M_p$ . It is well known (cf. Kato [25, p. 184]) that  $0 \in \rho(A_m)$  is equivalent to  $0 \in \rho(A_m^*)$ . Thus we have the second equality in (5.9) similarly as above.

The following corollaries are obvious from Theorem 5.1 and Proposition 5.2.

**Corollary 5.1.** *For the differential difference equation*

$$\frac{dx(t)}{dt} = A_0x(t) + \sum_{r=1}^m A_r x(t-h_r),$$

*a necessary and sufficient condition for  $\text{Ker } F = \{0\}$  (resp.  $\text{Ker } F^* = \{0\}$ ) is  $\text{Ker } A_m = \{0\}$  (resp.  $\text{Ker } A_m^* = \{0\}$ ).*

**Corollary 5.2.** *Assume that  $A_I(s) = 0$  in a neighbourhood of  $-h$ . If  $0 \in \rho(A_m)$ , then  $0 \in \rho(F)$  and  $0 \in \rho(F^*)$ , in other words,  $F$  and  $F^*$  are boundedly invertible.*

The above results are infinite dimensional analogue of those given in Delfour and Manitius [15, Section 2.1], in which the proofs are more complicated than those given here, because they have intended to include a very general Stieltjes measure  $\eta$  on  $\mathbf{R}^n$  of bounded variation. Our proofs are simple and easy because of the restricted form of  $\eta$  given in (2.3).

### 6. Representations of resolvent operators

This section is devoted to give convenient forms of the resolvents of  $A, A_T, A^*$  and  $A_T^*$ . In order to give such forms we require some definitions. According to Delfour and Manitius [15], Burns and Herdman [7] and Salamon [40] we introduce the following linear operators  $E_\lambda, T_\lambda, K_\lambda$  and  $H_\lambda$ . Let  $\lambda \in \mathbf{C}^1$  and an ordered pair of spaces  $(Y, Z_p)$  be the pair  $(X, M_p)$  or  $(X^*, M_p^*)$ . Define  $E_\lambda: Y \rightarrow Z_p, T_\lambda: Z_p \rightarrow Z_p, K_\lambda: Z_p \rightarrow Z_p$  and  $H_\lambda: Z_p \rightarrow Y$  by

$$(6.1) \quad \begin{cases} [E_\lambda z]^0 = z, \\ [E_\lambda z]^1(s) = e^{\lambda s} z, \quad s \in I_h \end{cases} \quad \text{for } z \in Y,$$

$$(6.2) \quad \begin{cases} [T_\lambda y]^0 = 0, \\ [T_\lambda y]^1(s) = \int_s^0 e^{\lambda(s-\xi)} y^1(\xi) d\xi, \quad s \in I_h \end{cases} \quad \text{for } y = (y^0, y^1) \in Z_p,$$



$$(6.3) \quad \begin{cases} [K_\lambda y]^0 = 0, \\ [K_\lambda y]^1(s) = \int_{-h}^s e^{\lambda(\xi-s)} y^1(\xi) d\xi, \end{cases} \quad \text{for } y = (y^0, y^1) \in Z_p, \quad s \in I_h$$

$$(6.4) \quad H_\lambda y = y^0 + \int_{-h}^0 e^{\lambda s} y^1(s) ds \quad \text{for } y = (y^0, y^1) \in Z_p,$$

respectively. The operator  $E_\lambda$  is often called the exponential map. All above are operator valued entire functions in  $\lambda$ . In what follows we denote the  $k$ -th derivative  $\frac{d^k}{d\lambda^k} f(\lambda)$  of  $f(\lambda)$  by  $f^{(k)}(\lambda)$ , or simply  $f^{(k)}$  for  $k=0, 1, 2, \dots$ .

**Proposition 6.1.** For each  $\lambda \in \mathbf{C}^1$  and integer  $k \geq 0$ ,

(i)

$$(6.5) \quad FT_\lambda^k = K_\lambda^k F, \quad F^* T_\lambda^k = K_\lambda^k F^*;$$

(ii)

$$(6.6) \quad T_\lambda^k E_\lambda = \frac{(-1)^k}{k!} E_\lambda^{(k)};$$

(iii)

$$(6.7) \quad (FE_\lambda)^{(k)} = FE_\lambda^{(k)}, \quad (F^* E_\lambda)^{(k)} = F^* E_\lambda^{(k)};$$

(iv)

$$(6.8) \quad H_\lambda FT_\lambda^k E_\lambda = \frac{(-1)^k}{(k+1)!} \Delta^{(k+1)}(\lambda), \quad H_\lambda F^* T_\lambda^k E_\lambda = \frac{(-1)^k}{(k+1)!} \Delta_T^{(k+1)}(\lambda).$$

Proof. (i). First we shall show  $FT_\lambda^k = K_\lambda^k F$  for  $k=1$ . Let  $g = (g^0, g^1) \in M_p$ . Since  $[FT_\lambda g]^0 = [K_\lambda Fg]^0 = 0$ , in order to prove  $FT_\lambda = K_\lambda F$  we have to prove, by (6.2) and (6.3), that

$$(6.9) \quad \int_{-h}^s e^{\lambda(\xi-s)} \int_{-h}^\xi d\eta(\tau) g^1(\tau - \xi) d\xi = \int_{-h}^s d\eta(\tau) \int_{\tau-s}^0 e^{\lambda(\tau-s-\beta)} g^1(\beta) d\beta, \quad s \in I_h.$$

Since the relation can be shown with the aid of the Fubini theorem, the detailed proof is omitted. The equality  $FT_\lambda^k = K_\lambda^k F$  for  $k \geq 2$  follows easily by induction. Since  $F^*$  has the same form as  $F$  (Proposition 4.2), we can verify  $F^* T_\lambda^k = K_\lambda^k F^*$  similarly as above.

(ii), (iii). These parts are proved easily by straightforward calculations using the definitions (4.1), (6.1), (6.2) of  $F, E_\lambda, T_\lambda$ .

(iv). We prove only the first equality in (6.8). By virtue of (6.4) and (6.6), the element  $H_\lambda FT_\lambda^k E_\lambda x, x \in X$  is written by

$$(6.10) \quad \begin{aligned} H_\lambda FT_\lambda^k E_\lambda x &= \frac{(-1)^k}{k!} H_\lambda FE_\lambda^{(k)} x \\ &= \delta_{k,0} x + (-1)^k \int_{-h}^0 e^{\lambda \xi} \int_{-h}^\xi d\eta(s) \frac{(s-\xi)^k}{k!} e^{\lambda(s-\xi)} x ds. \end{aligned}$$

Thus, the equations (6.10) and (6.9) with  $g^1(s) = \frac{1}{k!} s^k e^{\lambda s} x$  imply that

$$\begin{aligned} H_\lambda F T_\lambda^k E_\lambda x &= \delta_{k,0} x + (-1)^k \int_{-h}^0 d\eta(s) \int_s^0 e^{\lambda s} \cdot \frac{\xi^k}{k!} x d\xi \\ &= \delta_{k,0} x + (-1)^{k+1} \int_{-h}^0 d\eta(s) e^{\lambda s} \cdot \frac{s^{k+1}}{(k+1)!} x \\ &= \delta_{k,0} x - \frac{(-1)^k}{(k+1)!} \cdot \frac{d^{k+1}}{d\lambda^{k+1}} \int_{-h}^0 e^{\lambda s} d\eta(s) x \\ &= \frac{(-1)^k}{(k+1)!} \Delta^{(k+1)}(\lambda) x. \end{aligned}$$

This completes the proof.

Now we can give explicit representations of the resolvents of  $A$  and  $A_T$  in terms of the retarded resolvent, structural operator  $F$  and other operators introduced in this section.

**Theorem 6.1.** (i)  $\rho(A) = \rho(A_0, \eta)$  and the resolvent  $R(\lambda; A)$  of  $A$  is given by

$$(6.11) \quad R(\lambda; A) = E_\lambda R(\lambda; A_0, \eta) H_\lambda F + T_\lambda, \quad \lambda \in \rho(A).$$

(ii)  $\rho(A_T) = \rho(A_0^*, \eta^*)$  and the resolvent  $R(\lambda; A_T)$  of  $A_T$  is given by

$$(6.12) \quad R(\lambda; A_T) = E_\lambda R(\lambda; A_0^*, \eta^*) H_\lambda F^* + T_\lambda, \quad \lambda \in \rho(A_T).$$

Proof. For a given  $\phi = (\phi^0, \phi^1) \in M_p$ , we construct a  $g = (g(0), g(\cdot)) \in D(A)$  such that  $(\lambda I - A)g = \phi$ . This is equivalent, in view of Proposition 3.1 (iii), to that

$$(6.13) \quad \lambda g(0) - A_0 g(0) - \int_{-h}^0 d\eta(s) g(s) = \phi^0, \quad g(0) \in D(A_0)$$

$$(6.14) \quad \lambda g(s) - \frac{d}{ds} g(s) = \phi^1(s), \quad s \in I_h.$$

We solve the differential equation (6.14) to obtain

$$(6.15) \quad g(s) = e^{\lambda s} g(0) + \int_s^0 e^{\lambda(s-\xi)} \phi^1(\xi) d\xi,$$

i.e.,

$$(6.16) \quad (g(0), g) = E_\lambda g(0) + T_\lambda \phi.$$

Substituting (6.15) in (6.13) and using (6.9), we have

$$\begin{aligned} (6.17) \quad \Delta(\lambda)g(0) &= \int_{-h}^0 d\eta(s) \int_s^0 e^{\lambda(s-\xi)} \phi^1(\xi) d\xi + \phi^0 \\ &= \int_{-h}^0 e^{\lambda s} \int_{-h}^s d\eta(\xi) \phi^1(\xi - s) ds + \phi^0 = H_\lambda F \phi. \end{aligned}$$

Assume that  $\lambda \in \rho(A_0, \eta)$ . Then by definition,  $\Delta(\lambda)$  has a bounded inverse  $\Delta^{-1}(\lambda) = R(\lambda; A_0, \eta)$ . So that by (6.16) and (6.17), we derive

$$(6.18) \quad (g(0), g) = E_\lambda R(\lambda; A_0, \eta) H_\lambda F \phi + T_\lambda \phi .$$

Since all operators appearing in (6.18) are bounded,  $\lambda \in \rho(A)$  and the resolvent  $R(\lambda; A)$  is given by (6.11). Next we show the inclusion  $\rho(A) \subset \rho(A_0, \eta)$ . Let  $\lambda \in \rho(A)$ . Then for any  $\phi = (\phi^0, \phi^1) \in M_p$ , there exists a unique  $g = (g(0), g) \in D(A)$  such that  $(\lambda I - A)g = \phi$ , or equivalently, (6.16) and (6.17) hold. We note that  $\Delta(\lambda)$  is one to one. Because if not, there exists a  $g^0 \in D(A_0)$ ,  $g^0 \neq 0$  such that  $\Delta(\lambda)g^0 = 0$ . The element  $g = E_\lambda g^0 \in D(A)$  satisfies  $(\lambda I - A)g = 0$ ,  $g \neq 0$ , which contradicts to  $\lambda \in \rho(A)$ . For special  $\phi = (\phi^0, 0)$ ,  $\phi^0 \in X$ , the equality (6.17) means that there exists a  $g = (g(0), g) \in D(A)$  such that  $\Delta(\lambda)g(0) = \phi^0$ . This concludes that the densely defined closed linear operator  $\Delta(\lambda): D(A_0) \subset X \rightarrow X$  is onto and one to one. Hence by open mapping theorem,  $\Delta(\lambda)^{-1}$  exists and is bounded, i.e.,  $\lambda \in \rho(A_0, \eta)$ . Therefore (i) is proved. The part (ii) is proved in quite analogous manner as in (i).

Next we characterize the resolvents of the adjoint operators  $A^*$  and  $A_0^*$ .

**Lemma 6.1.** *The relation*

$$(6.19) \quad (\lambda I - A^*)f = \psi, \quad f \in D(A^*), \quad \psi \in M_p^*$$

*is equivalent to*

$$(6.20) \quad \Delta_T(\lambda)f^0 = H_\lambda \psi, \quad f^0 \in D(A_0^*) \quad \text{and} \quad f = K_\lambda \psi + F^* E_\lambda f^0 .$$

*Proof.* In view of Proposition 3.3, (6.19) is written by the following equivalent condition

$$(6.21) \quad \lambda f^0 - A_0^* f^0 - f^1(0) = \psi^0, \quad f^0 \in D(A_0^*),$$

$$(6.22) \quad \lambda f^1(s) - \frac{d}{ds} w(f)(s) = \psi^1(s), \quad s \in I_h,$$

where  $w(f)$  is given in (3.19). Put  $\tilde{f}^0 = (f^0, f^0) \in M_p^*$  i.e.,  $[\tilde{f}^0]^0 = f^0$ ,  $[\tilde{f}^0]^1(s) \equiv f^0$ , and  $\tilde{w} = -w(f)$ . Since  $f^1 = [F^* \tilde{f}^0]^1 + \tilde{w}$  by (3.19), we can solve the differential equation (6.22) with the initial condition  $\tilde{w}(-h) = 0$  to obtain

$$\begin{aligned} \tilde{w}(s) &= e^{-\lambda s} \int_{-h}^s e^{\lambda \xi} \{ \psi^1(\xi) - \lambda [F^* \tilde{f}^0]^1(\xi) \} d\xi \\ &= \int_{-h}^s e^{-\lambda(s-\xi)} \psi^1(\xi) d\xi - \int_{-h}^s e^{\lambda(\xi-s)} \lambda \int_{-h}^\xi d\eta^*(\beta) f^0 d\xi, \quad s \in I_h . \end{aligned}$$

By (6.3) and applying the Fubini theorem to the last term of the above equality, we obtain without difficulty that

$$(6.23) \quad \tilde{w}(s) = [K_\lambda \psi]^1(s) - [F^* \tilde{f}^0]^1(s) + [F^* E_\lambda f^0]^1(s), \quad s \in I_h.$$

Thus,

$$f^1 = [F^* \tilde{f}^0]^1 + \tilde{w} = [K_\lambda \psi]^1 + [F^* E_\lambda f^0]^1,$$

which shows the second equality in (6.20). Substituting  $s=0$  in (3.19) and (6.23), we have

$$f^1(0) = \int_{-h}^0 d\eta^*(s) f^0 + \tilde{w}(0) = \int_{-h}^0 e^{\lambda s} \psi^1(s) ds + \int_{-h}^0 e^{\lambda s} d\eta^*(s) f^0.$$

Hence the equality (6.21) is rewritten as

$$(\lambda I - A_0^* - \int_{-h}^0 e^{\lambda s} d\eta^*(s)) f^0 = \psi^0 + \int_{-h}^0 e^{\lambda s} \psi^1(s) ds, \quad f^0 \in D(A_0^*),$$

which is the first equality in (6.20).

**Theorem 6.2.** (i)  $\rho(A^*) = \rho(A_0^*, \eta^*)$  and the resolvent  $R(\lambda; A^*)$  of  $A^*$  is given by

$$(6.24) \quad R(\lambda; A^*) = F^* E_\lambda R(\lambda; A_0^*, \eta^*) H_\lambda + K_\lambda, \quad \lambda \in \rho(A^*).$$

(ii)  $\rho(A^\#) = \rho(A_0, \eta)$  and the resolvent  $R(\lambda; A^\#)$  of  $A^\#$  is given by

$$(6.25) \quad R(\lambda; A^\#) = F E_\lambda R(\lambda; A_0, \eta) H_\lambda + K_\lambda, \quad \lambda \in \rho(A^\#).$$

Proof. Using Lemma 6.1 we can prove (i) by analogous argument as in the proof of Theorem 6.1. The proof of the remaining part (ii) is similar.

Here we give important relations between the operators  $E_\lambda, H_\lambda, T_\lambda$  and  $K_\lambda$ . Taking into account of the relation (2.18), we can verify by direct computations involving the pairing  $\langle \cdot, \cdot \rangle_{M_p}$  that for each  $\lambda \in \mathbf{C}^1$ ,

$$(6.26) \quad E_\lambda^* = H_{\bar{\lambda}}, \quad H_\lambda^* = E_{\bar{\lambda}}, \quad T_\lambda^* = K_{\bar{\lambda}}, \quad K_\lambda^* = T_{\bar{\lambda}}.$$

Consequently, by using the equality (2.17) and (6.26) Theorem 6.2 can be derived as the adjoint version of Theorem 6.1. This may be a simple proof of Theorem 6.2.

**Corollary 6.1.** (i)

$$(6.27) \quad FR(\lambda; A) = R(\lambda; A^\#)F, \quad R(\lambda; A)G = GR(\lambda; A^\#)$$

for  $\lambda \in \rho(A) = \rho(A^\#) = \rho(A_0, \eta)$ .

(ii)

$$(6.28) \quad R(\lambda; A^*)F^* = F^*R(\lambda; A_T), \quad G^*R(\lambda; A^*) = R(\lambda; A_T)G^*$$

for  $\lambda \in \rho(A^*) = \rho(A_T) = \rho(A_0^*, \eta^*)$ .

Proof. (i) follows from Theorems 4.1, 4.2, 6.1 and 6.2. (ii) follows from Proposition 6.1 (i), Theorems 6.1 and 6.2.

**7. Spectral decomposition**

In this and following sections we study the spectral decomposition theory for the FDE's in Banach spaces. The spectral theory for various types of FDE's in  $\mathbf{R}^n$  is further developed by many authors (see [6, 15, 18, 19, 20, 24, 36, 40] for examples). An attempt to extend the spectral theory to retarded FDE's in infinite dimensional spaces was first made by Travis and Webb [44] whose main concern is the stability of mild solutions. Their analysis and investigations have been carried in the space  $C(I_n; X)$ , but seems incomplete compared with those for  $X=\mathbf{R}^n$ . The purpose here is to construct a rather complete spectral decomposition theory for the equation (E) on the space  $M_p$ , which extends the work of [15,29] to general Banach space case. Our analysis, however, is more delicate than those in [15, 29] because of permitting  $X$  being infinite dimensional.

7.1. Classification of spectrum

The retarded spectrum introduced in Section 2 is efficiently used to determine the spectrum of the infinitesimal generators associated with (E) and (E<sup>T</sup>).

**Proposition 7.1.** *Three kinds of spectrum of A and A<sub>T</sub> are given by*

$$(7.1) \quad \sigma_P(A) = \sigma_P(A_0, \eta), \quad \sigma_C(A) = \sigma_C(A_0, \eta), \quad \sigma_R(A) = \sigma_R(A_0, \eta),$$

$$(7.2) \quad \sigma_P(A_T) = \sigma_P(A_0^*, \eta^*), \quad \sigma_C(A_T) = \sigma_C(A_0^*, \eta^*), \quad \sigma_R(A_T) = \sigma_R(A_0^*, \eta^*),$$

respectively.

Proof. First we recall the following fact which is already shown in the proof of Theorem 6.1. That is, the relation  $(\lambda I - A)g = \phi, g \in D(A), \phi \in M_p$  is equivalent to that  $\Delta(\lambda)g^0 = H_\lambda F \phi, g^0 \in D(A_0), g = E_\lambda g^0 + T_\lambda \phi$ . If we substitute  $\phi = 0$  in the above equivalence, then we have that  $\text{Ker}(\lambda I - A) = \{0\}$  is equivalent to that  $\text{Ker} \Delta(\lambda) = \{0\}$ , and hence  $\text{Ker}(\lambda I - A) \neq \{0\}$  if and only if  $\text{Ker} \Delta(\lambda) \neq \{0\}$ . This concludes, by definition,  $\sigma_P(A) = \sigma_P(A_0, \eta)$ . By the same reason, from Lemma 6.1 it follows that  $\text{Ker}(\lambda I - A^*) = \{0\}$  if and only if  $\text{Ker} \Delta_T(\lambda) = \{0\}$ . Then by putting  $\lambda = \bar{\lambda}$  and using the duality theorem, we have that  $\text{C1}(\text{Im}(\bar{\lambda} I - A^*))^* = \text{C1}(\text{Im}(\lambda I - A)) = M_p$  if and only if  $\text{C1}(\text{Im} \Delta_T(\bar{\lambda})^*) = \text{C1}(\text{Im} \Delta(\lambda)) = X$ . This implies, by contradiction, that  $\text{Im} \Delta(\lambda)$  is not dense in  $X$  if and only if  $\text{Im}(\lambda I - A)$  is not dense in  $M_p$ . Now we are ready to prove  $\sigma_R(A_0, \eta) = \sigma_R(A)$ . From the definition of residual spectrum,  $\lambda \in \sigma_P(A_0, \eta)$  if and only if

$$(7.3) \quad \Delta(\lambda)^{-1} \text{ exists (i.e., } \text{Ker } \Delta(\lambda) = \{0\}) \text{ but } \text{Im } \Delta(\lambda) \text{ is not dense in } X.$$

It then follows that (7.3) is equivalent to

$$(7.4) \quad (\lambda I - A)^{-1} \text{ exists but } \text{Im}(\lambda I - A) \text{ is not dense in } M_p.$$

The statement (7.4) is exactly the definition of  $\lambda \in \sigma_R(A)$ . Hence  $\sigma_R(A_0, \eta) = \sigma_R(A)$  is proved. The rest equality  $\sigma_C(A_0, \eta) = \sigma_C(A)$  is now evident. The part for  $A_T$  is proved in a same manner as above.

REMARK 7.1. In the case where  $X = \mathbf{R}^n$ , it is well known that  $\sigma(A) = \sigma_P(A) = \{\lambda : \det \Delta(\lambda) = 0\}$  is countable and isolated. However there exists an operator  $A$  defined by (3.2) and (3.3) such that  $\sigma_C(A) \neq \phi$  or  $\sigma_R(A) \neq \phi$  in our infinite dimensional case.

7.2. Generalized eigenspaces and spectral decomposition

Let  $\lambda \in \sigma_P(A)$ . The generalized eigenspace  $\mathcal{M}_\lambda$  of  $A$  corresponding to  $\lambda$  is defined by

$$(7.5) \quad \mathcal{M}_\lambda = \bigcup_{l=0}^{\infty} \text{Ker}(\lambda I - A)^l.$$

To characterize the structure of  $\text{Ker}(\lambda I - A)^l$ ,  $l = 1, 2, \dots$ , we introduce operator valued matrices  $\mathcal{A}_l = \mathcal{A}_l(\lambda)$  defined by

$$(7.6) \quad \mathcal{A}_l = \begin{pmatrix} D_1 & D_2 & \cdots & D_l \\ 0 & D_1 & \cdots & D_{l-1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & D_1 \end{pmatrix}, \quad l = 1, 2, \dots$$

where

$$(7.7) \quad D_{j+1} = D_{j+1}(\lambda) = \frac{1}{j!} \Delta^{(j)}(\lambda), \quad j = 0, 1, 2, \dots$$

The following result extends the results of [15, Proposition 4.3] and [18, Lemma 3.3, p. 177] to reflexive Banach space case.

**Proposition 7.2.** *Let  $\lambda \in \sigma_P(A)$ . Then  $\text{Ker}(\lambda I - A)^l$  coincides with the space of functions  $\phi \in M_p$  of the form*

$$(7.8) \quad \phi = \sum_{j=0}^{l-1} \frac{1}{j!} (E_\lambda y_{j+1})^{(j)},$$

where  $\mathcal{Q} = \text{col}(y_1, \dots, y_l)$ ,  $y_{j+1} \in D(A_0)$ ,  $j = 0, 1, \dots, l-1$  satisfies  $\mathcal{A}_l \mathcal{Q} = 0$  in  $X^l$ .

Proof. Let  $\phi \in \text{Ker}(\lambda I - A)^l$ . Set  $\phi_0 = \phi$  and  $\phi_j = (\lambda I - A)\phi_{j-1}$ ,  $j = 1, \dots, l$ ; then  $\phi$  is characterized by  $\phi_l = 0$ . The relation  $\phi_j = (\lambda I - A)\phi_{j-1}$  is written as

$$(7.9) \quad \phi_{j-1} = E_\lambda \phi_{j-1}^0 + T_\lambda \phi_j \quad \text{and} \quad \Delta(\lambda) \phi_{j-1}^0 = H_\lambda F \phi_j, \quad \phi_{j-1}^0 \in D(A_0), \\ j = 1, \dots, l.$$

Since  $\phi_l=0$ , we have

$$(7.10) \quad \begin{cases} \phi_{l-1} = E\phi_{l-1}, & \phi_{l-2} = E_\lambda\phi_{l-2}^0 + T_\lambda\phi_{l-1} = E_\lambda\phi_{l-2}^0 + T_\lambda E_\lambda\phi_{l-1}^0, \dots, \\ \phi_j = \sum_{r=j}^{l-1} T_\lambda^{r-j} E_\lambda\phi_r^0, \dots, & \phi = \phi_0 = \sum_{r=0}^{l-1} T_\lambda^r E_\lambda\phi_r^0, \end{cases}$$

so that by (6.6) and a change  $r \rightarrow j$ ,

$$\phi = \sum_{j=0}^{l-1} \frac{1}{j!} (E_\lambda(-1)^j \phi_j^0)^{(j)}.$$

If we put  $y_{i+1} = (-1)^j \phi_j^0 \in D(A_0)$ ,  $j = 0, 1, \dots, l-1$ , we obtain (7.8). From the second equality in (7.9) it follows by (7.10) that

$$(7.11) \quad \begin{aligned} \Delta(\lambda)y_j &= (-1)^{j-1} H_\lambda F \phi_j = (-1)^{j-1} \left( \sum_{r=j}^{l-1} H_\lambda F T_\lambda^{r-j} E_\lambda(-1)^{-r} y_{r+1} \right) \\ &= (-1) \sum_{r=j}^{l-1} (-1)^{j-r} H_\lambda F T_\lambda^{r-j} E_\lambda y_{r+1}. \end{aligned}$$

By virtue of (6.8), the equation (7.11) is written as

$$\Delta(\lambda)y_j = (-1) \sum_{r=j}^{l-1} \frac{1}{(r-j+1)!} \Delta^{(r-j+1)}(\lambda) y_{r+1},$$

and hence (by changing  $r \rightarrow r+1$ ),

$$(7.12) \quad \sum_{r=j}^l \frac{1}{(r-j)!} \Delta^{(r-j)}(\lambda) y_r = 0, \quad j = 1, \dots, l.$$

The system of equations (7.12) is rewritten simply by  $\mathcal{A}_l \mathcal{Q} = 0$  in  $X^l$ , where  $\mathcal{Q} = \text{col}(y_1, \dots, y_l)$ . This completes the proof.

In order to go into a further spectral decomposition theory we have to restrict  $\lambda$  to the isolated spectrum. We now require the following definitions:

$$\begin{aligned} \sigma_I(A_0, \eta) &= \{ \lambda \in \mathbf{C}^1: \lambda \text{ is an isolated singular point of } R(\mu; A_0, \eta) \}, \\ \sigma_0(A_0, \eta) &= \{ \lambda \in \mathbf{C}^1: \lambda \text{ is a pole of } R(\mu; A_0, \eta) \}. \end{aligned}$$

Similarly we define the spectral sets  $\sigma_I(A)$  and  $\sigma_0(A)$ . It is obvious that  $\sigma_0(A_0, \eta) \subset \sigma_I(A_0, \eta)$ ,  $\sigma_0(A) \subset \sigma_I(A)$ . Since all  $E_\lambda$ ,  $H_\lambda$  and  $T_\lambda$  are entire functions, Theorem 6.1 (i) implies that  $\lambda$  is a pole of  $R(\mu; A)$  of order  $k_\lambda$  if and only if  $\lambda$  is a pole of  $R(\mu; A_0, \eta)$  of same order  $k_\lambda$ . Then,  $\sigma_0(A) = \sigma_0(A_0, \eta)$ .

Let  $\lambda \in \sigma_I(A)$  and let  $P_\lambda$  be the projection operator

$$(7.13) \quad P_\lambda = \frac{1}{2\pi i} \int_{\Gamma_\lambda} R(\mu; A) d\mu,$$

where  $\Gamma_\lambda$  is a closed rectifiable curve containing  $\lambda$  inside and all other singular

points of  $R(\mu; A)$  outside. By Yosida [49, p. 228–231] (see also Taylor and Lay [41], Tanabe [43] and Kato [25]), we obtain the following decomposition of the space  $M_p$ .

**Theorem 7.1.** *Let  $\lambda$  be a pole of  $R(\mu; A_0, \eta)$  of order  $k_\lambda$ . Then  $\lambda \in \sigma_p(A)$  and the direct sum decomposition*

$$(7.14) \quad M_p = \text{Ker}(\lambda I - A)^{k_\lambda} \oplus \text{Im}(\lambda I - A)^{k_\lambda}, \quad \mathcal{M}_\lambda = P_\lambda M_p = \text{Ker}(\lambda I - A)^{k_\lambda}$$

holds. Both  $\mathcal{M}_\lambda$  and  $\text{Im}(\lambda I - A)^{k_\lambda} = \text{Ker} P_\lambda$  are closed and invariant under  $S(t)$ . Moreover the resolvent  $R(\mu; A)$  has the Laurent series expansion

$$(7.15) \quad R(\mu; A) = \sum_{n=-k_\lambda}^{\infty} (\mu - \lambda)^n Q_n$$

in a neighbourhood of  $\lambda$ , where  $Q_n$  is given by

$$(7.16) \quad Q_n = \frac{1}{2\pi i} \int_{\Gamma_\lambda} (\mu - \lambda)^{-n-1} R(\mu; A) d\mu.$$

Clearly  $P_\lambda = Q_{-1}$ . Put  $Q_\lambda = Q_{-2}$ , then from the expression (7.16) (cf. Kato [25, p. 180]) it follows that

$$(7.17) \quad Q_{-n} = Q_\lambda^{n-1}, \quad n = 2, \dots, k_\lambda, \quad Q_\lambda^{k_\lambda} = O \text{ (nilpotent),}$$

$$(7.18) \quad P_\lambda Q_\lambda = Q_\lambda P_\lambda = Q_\lambda, \quad AP_\lambda = \lambda P_\lambda + Q_\lambda.$$

The decomposition of  $M_p$  in Theorem 7.1 is slightly generalized as

$$(7.19) \quad M_p = \left( \bigoplus_{\lambda \in \Lambda} \mathcal{M}_\lambda \right) \oplus \mathcal{R}_\lambda, \quad \mathcal{R}_\lambda = \bigcap_{\lambda \in \Lambda} \text{Im}(\lambda I - A)^{k_\lambda},$$

where  $\Lambda \subset \sigma_0(A)$  is a finite set (see e.g., Kato [25, p. 181]).

REMARK 7.2. Proposition 7.2 tells us  $\dim \text{Ker}(\lambda I - A)^l = \dim \text{Ker} \mathcal{A}_l$ , but the dimension may be infinity even if  $\lambda$  is a pole of  $R(\mu; A)$ .

From (7.13), (7.16), (7.18) and (6.11) the following corollary follows at once. Notice that  $T_\lambda$  is entire.

**Corollary 7.1.** *For  $\lambda \in \sigma_0(A_0, \eta)$ ,*

(i)

$$(7.20) \quad P_\lambda = \mathcal{E}_\lambda F, \quad \mathcal{E}_\lambda = \frac{1}{2\pi i} \int_{\Gamma_\lambda} E_\mu R(\mu; A_0, \eta) H_\mu d\mu;$$

$$(7.21) \quad Q_\lambda = \mathcal{F}_\lambda F, \quad \mathcal{F}_\lambda = \frac{1}{2\pi i} \int_{\Gamma_\lambda} (\mu - \lambda) E_\mu R(\mu; A_0, \eta) H_\mu d\mu;$$

(ii)

$$(7.22) \quad \text{Ker} F \subset \text{Ker} P_\lambda \subset \text{Ker} Q_\lambda.$$



Next we consider the case for the transposed operator  $A_T$ . Let  $\mathcal{M}_\lambda^T$  denote the generalized eigenspace of  $A_T$  corresponding to  $\lambda \in \sigma_P(A_T)$ ; let the matrices  $\mathcal{A}_l^T = \mathcal{A}_l^T(\lambda)$ ,  $l=1, 2, \dots$  be defined by (7.6) in which  $D_{j+1}$  is replaced by  $\frac{1}{j!} \Delta_T^{(j)}(\lambda)$ ,  $j=0, 1, 2, \dots$ ; and let  $P_\lambda^T$  denote the spectral projection corresponding to  $\lambda \in \sigma_I(A_T)$ . Then we have:

**Theorem 7.2.** (i) *If  $\lambda$  is a pole of  $R(\mu; A_\delta^*, \eta^*)$  of order  $m_\lambda$ , then  $\lambda \in \sigma_P(A_T)$  and the decomposition*

$$M_p^* = \text{Ker}(\lambda I - A_T)^{m_\lambda} \oplus \text{Im}(\lambda I - A_T)^{m_\lambda}, \quad \mathcal{M}_\lambda^T = P_\lambda^T M_p^* = \text{Ker}(\lambda I - A_T)^{m_\lambda}$$

holds. Moreover the resolvent  $R(\mu; A_T)$  has the Laurent series expansion

$$R(\mu; A_T) = \sum_{n=-m_\lambda}^{\infty} (\mu - \lambda)^n Q_n^T$$

in a neighbourhood of  $\lambda$ , where  $Q_n^T$  is given by (7.16) in which  $R(\mu; A)$  is replaced by  $R(\mu; A_T)$ .

(ii) *For  $\lambda \in \sigma_P(A_T)$ , the space  $\text{Ker}(\lambda I - A_T)^l$  coincides with the set of functions  $\psi \in M_p^*$  of the form*

$$\psi = \sum_{j=0}^{l-1} \frac{1}{j!} (E_\lambda y_{j+1}^T)^{(j)},$$

where  $Q_j^T = \text{col}(y_1^T, \dots, y_l^T)$ ,  $y_{j+1}^T \in D(A_\delta^*)$ ,  $j=0, 1, \dots, l-1$  satisfies  $\mathcal{A}_l^T Q_j^T = 0$  in  $X^{*l}$ .

(iii) *For  $\lambda \in \sigma_0(A_\delta^*, \eta^*)$ ,*

$$P_\lambda^T = \mathcal{E}_\lambda^T F^*; \quad \mathcal{E}_\lambda^T = \frac{1}{2\pi i} \int_{\Gamma_\lambda} E_\mu R(\mu; A_\delta^*, \eta^*) H_\mu d\mu,$$

$$Q_\lambda^T = \mathcal{F}_\lambda^T F^*; \quad \mathcal{F}_\lambda^T = \frac{1}{2\pi i} \int_{\Gamma_\lambda} (\mu - \lambda) E_\mu R(\mu; A_\delta^*, \eta^*) H_\mu d\mu,$$

where  $Q_\lambda^T = Q_{l-2}^T$  and

$$\text{Ker } F^* \subset \text{Ker } P_\lambda^T \subset \text{Ker } Q_\lambda^T.$$

We shall describe a group property of  $S(t)$ . For this the following discrete spectrum  $\sigma_d(A)$  of  $A$  is needed to be defined by

$$\sigma_d(A) = \{\lambda \in \sigma_I(A) : \dim(\text{Im } P_\lambda) < +\infty\}.$$

It is well known (Kato [25, p. 181]) that  $\sigma_d(A) \subset \sigma_0(A) \subset \sigma_P(A)$  and  $\text{Im } P_\lambda = \mathcal{M}_\lambda = \text{Ker}(\lambda I - A)^{k_\lambda}$  for  $\lambda \in \sigma_0(A)$ . Let  $\lambda \in \sigma_d(A)$  and let  $d_\lambda = \dim \mathcal{M}_\lambda$ . We shall write by  $\Phi_\lambda = \{\phi_{\lambda,1}, \dots, \phi_{\lambda,d_\lambda}\}$  a basis of  $\mathcal{M}_\lambda$  of the form (7.8). Since  $A\mathcal{M}_\lambda \subset \mathcal{M}_\lambda$ , there exists a  $d_\lambda \times d_\lambda$  matrix  $B_\lambda$  such that

$$(7.23) \quad A\Phi_\lambda = \Phi_\lambda B_\lambda \quad \text{and} \quad S(t)\Phi_\lambda = \Phi_\lambda e^{B_\lambda t} \quad \text{for } t \geq 0.$$

Hence the only eigenvalue of  $B_\lambda$  is  $\lambda$  and  $S(t)$  can be extended to a holomorphic group on  $\mathcal{M}_\lambda$ . Now we can prove the following result in a similar fashion as in Hale [18, Chapter 7, Theorem 2.2] with suitable modifications to the space  $M_p$ .

**Proposition 7.3.** *Assume that  $\Lambda \subset \sigma_d(A)$  is a finite set  $\{\lambda_1, \dots, \lambda_n\}$ . Let  $\Phi_\Lambda = \{\Phi_{\lambda_1}, \dots, \Phi_{\lambda_n}\}$  and  $B_\Lambda = \text{diag}(B_{\lambda_1}, \dots, B_{\lambda_n})$ . Then for any column vector  $a$  of the same dimension as  $\Phi_\Lambda (= d_{\lambda_1} + \dots + d_{\lambda_n})$ , we have*

(i)  $S(t)\Phi_\Lambda a$  can be defined on  $(-\infty, \infty)$  by the relation

$$S(t)\Phi_\Lambda a = \Phi_\Lambda \exp(tB_\Lambda)a$$

where

$$\begin{cases} [\Phi_\Lambda]^1(s) = [\Phi_\Lambda]^0 \exp(sB_\Lambda), & s \in I_h, \\ [\Phi_\Lambda]^0 = \{\Phi_{\lambda_1}(0), \dots, \Phi_{\lambda_n}(0)\}; \end{cases}$$

(ii)  $\begin{cases} [S(t)\Phi_\Lambda a]^1(s) = [\Phi_\Lambda \exp((t+s)B_\Lambda)a]^0, & s \in I_h, \\ [S(t)\Phi_\Lambda a]^0 = [\Phi_\Lambda]^0 \exp(tB_\Lambda)a, \end{cases}$  for  $t \geq 0$ ;

(iii)  $x(t) = [S(t)\Phi_\Lambda a]^0$  is a mild solution (in fact, a strong solution) of (E) on  $(-\infty, \infty)$  with the initial condition  $g = \Phi_\Lambda a$  and  $u = 0$ ;

(iv)  $M_p$  is decomposed into the direct sum

$$M_p = \mathcal{M}_\Lambda \oplus \mathcal{R}_\Lambda, \quad \mathcal{M}_\Lambda = \bigoplus_{\lambda \in \Lambda} \mathcal{M}_\lambda$$

as in (7.19), where  $\mathcal{M}_\Lambda$  is given by

$$\mathcal{M}_\Lambda = \{\phi \in M_p : \phi = \Phi_\Lambda a \text{ for some } a \in \mathbb{C}^{\dim \Phi_\Lambda}\}.$$

Moreover,  $\begin{cases} S(t)\mathcal{M}_\Lambda \subset \mathcal{M}_\Lambda & \text{for all } t \in (-\infty, \infty) \\ S(t)\mathcal{R}_\Lambda \subset \mathcal{R}_\Lambda & \text{for all } t \geq 0. \end{cases}$

The above proposition gives a precise information on the asymptotic behaviour of the mild solution of (E) on generalized eigenspaces for discrete spectrum. Analogous result to Proposition 7.3 for  $A_T$  holds true.

**REMARK 7.3.** If  $A_0$  has compact resolvent, then the retarded resolvent  $R(\lambda; A_0, \eta)$  is compact. From this and the representation (7.20) the compactness of  $P_\lambda$  follows, so that  $\sigma_I(A) = \sigma_d(A)$  is true in this case.

**Proposition 7.4.** (i) For  $\lambda \in \sigma_0(A)$ ,  $F$  is one to one on  $\mathcal{M}_\lambda$ .

(ii) For  $\lambda \in \sigma_0(A_T)$ ,  $F^*$  is one to one on  $\mathcal{M}_\lambda^T$ .

**Proof.** We prove only (i). Assume first that  $\phi \in \text{Ker}(\lambda I - A)$  and  $F\phi = 0$ . Then by (4.11),  $0 = GF\phi = S(h)\phi = e^{\lambda h}\phi$  and hence  $\phi = 0$ . This shows  $F$  is one to one on  $\text{Ker}(\lambda I - A)$ . Next assume  $\phi \in \mathcal{M}_\lambda = \text{Ker}(\lambda I - A)^{k_\lambda}$  and  $F\phi = 0$ . If we set  $\phi_1 = (\lambda I - A)^{k_\lambda - 1}\phi$ , then  $\phi_1 \in \text{Ker}(\lambda I - A)$  and

$$S(h)\phi_1 = (\lambda I - A)^{k_\lambda - 1} S(h)\phi = (\lambda I - A)^{k_\lambda - 1} GF\phi = 0.$$

So that  $\phi_1 = 0$ . Continuing this procedure  $k_\lambda$  times we have  $\phi = 0$ , i.e.,  $F$  is one to one on  $\mathcal{M}_\lambda$ .

## 8. Adjoint spectral decomposition

In this section we study the spectral decomposition theory for the adjoint operator  $A^*$  of  $A$  in the space  $M_p^*$ , with an emphasis of the relations between  $A^*$  and the transposed operator  $A_T$ . The structural operators  $F^*$  and  $G^*$  will appear to key connections between the generalized eigenspaces of  $A^*$  and  $A_T$ .

### 8.1. Generalized eigenspaces and structural operators

Let  $\mathcal{M}_\lambda^*$  (not the adjoint space of  $\mathcal{M}_\lambda$ !) denote the generalized eigenspace of  $A^*$  corresponding to  $\lambda \in \sigma_p(A^*)$ . Similarly we denote by  $\mathcal{M}_\lambda^{T*}$  the generalized eigenspace of  $A_T^*$  corresponding to  $\lambda \in \sigma_p(A_T^*)$ .

**Theorem 8.1.** (i) *Three kinds of spectrum of  $A^*$  and  $A_T$  are identical and are given by*

$$(8.1) \quad \begin{aligned} \sigma_p(A^*) &= \sigma_p(A_T) = \sigma_p(A_0^*, \eta^*), & \sigma_R(A^*) &= \sigma_R(A_T) = \sigma_R(A_0^*, \eta^*), \\ \sigma_C(A^*) &= \sigma_C(A_T) = \sigma_C(A_0^*, \eta^*), \end{aligned}$$

respectively.

(ii) *For each  $\lambda \in \sigma_p(A^*) = \sigma_p(A_T)$ ,*

$$(8.2) \quad \text{Ker}(\lambda I - A^*)^l = F^* \text{Ker}(\lambda I - A_T)^l, \quad l = 1, 2, \dots.$$

*In particular*

$$(8.3) \quad \mathcal{M}_\lambda^* = F^* \mathcal{M}_\lambda^{T*}.$$

Proof. (i). Using Lemma 6.1 we can prove this part by similar arguments as in the proof of Proposition 7.1.

(ii). By (4.30),  $(\lambda I - A^*)F^* = F^*(\lambda I - A_T)$  on  $D(A_T)$ . Hence, by induction, we have

$$(\lambda I - A^*)^l F^* = F^*(\lambda I - A_T)^l \quad \text{on } D(A_T^l).$$

Thus

$$(8.4) \quad F^* \text{Ker}(\lambda I - A_T)^l \subset \text{Ker}(\lambda I - A^*)^l, \quad l = 1, 2, \dots.$$

The reverse inclusions in (8.4) were proved by Delfour and Manitius [15] by using mathematical induction. Here we give a direct proof based on Lemma 6.1. Let  $\psi \in \text{Ker}(\lambda I - A^*)^l$  and put

$$(8.5) \quad \psi_0 = \psi \quad \text{and} \quad \psi_j = (\lambda I - A^*)\psi_{j-1}, \quad j = 1, \dots, l.$$

Then  $\psi \in \text{Ker}(\lambda I - A^*)^l$  is equivalent to  $\psi_l = 0$ . In view of Lemma 6.1 we see that  $\psi_j = (\lambda I - A^*)\psi_{j-1}$  is equivalent to

$$(8.6) \quad \Delta_T(\lambda)\psi_{j-1}^0 = H_\lambda\psi_j, \quad \psi_{j-1}^0 \in D(A_0^*) \quad \text{and} \quad \psi_{j-1} = K_\lambda\psi_j + F^*E_\lambda\psi_{j-1}^0.$$

Since  $\psi_l = 0$ , it follows from the last equality in (8.6) that

$$\begin{aligned} \psi_{l-1} &= F^*E_\lambda\psi_{l-1}^0 \\ \psi_{l-2} &= F^*E_\lambda\psi_{l-2}^0 + K_\lambda\psi_{l-1} = F^*E_\lambda\psi_{l-2}^0 + K_\lambda F^*E_\lambda\psi_{l-1}^0 \\ &= F^*E_\lambda\psi_{l-2}^0 + F^*T_\lambda E_\lambda\psi_{l-1}^0 = F^*(E_\lambda\psi_{l-2}^0 + T_\lambda E_\lambda\psi_{l-1}^0), \quad (\text{by (6.5)}) \\ &\dots\dots\dots, \\ \psi &= \psi_0 = F^*\left(\sum_{j=0}^{l-1} T_\lambda^j E_\lambda\psi_j^0\right). \end{aligned}$$

If we set  $y_{j+1}^T = (-1)^j \psi_j^0, j=0, 1, \dots, l-1$ , then by (6.6)  $\psi$  can be written as  $\psi = F^*\left(\sum_{j=0}^{l-1} \frac{1}{j!} (E_\lambda y_{j+1}^T)^{(j)}\right)$ , where  $Q^T = \text{col}(y_1^T, \dots, y_l^T)$  satisfies  $\mathcal{A}_l^T Q^T = 0$  in  $X^{*l}$ .

Hence by Theorem 7.2 (ii),  $\psi \in F^* \text{Ker}(\lambda I - A_T)^l$ , i.e., the reverse inclusions of (8.4) are proved. Therefore (8.2) is shown. The rest equality (8.3) is clear from (8.2) and the definition of generalized eigenspaces.

The statement (ii) of Proposition 8.1 has concluded that the null space  $\text{Ker}(\lambda I - A^*)^l$  is obtained by the application of  $F^*$  to  $\text{Ker}(\lambda I - A_T)^l$ , whose elements are straightforwardly computed as given in Theorem 7.2 (ii).

If we notice  $(A_T)_T = A$ , we have the following result.

**Corollary 8.1.** (i)  $\sigma_P(A_\#) = \sigma_P(A) = \sigma_P(A_0, \eta)$ ,

$$\sigma_R(A_\#) = \sigma_R(A) = \sigma_R(A_0, \eta), \quad \sigma_C(A_\#) = \sigma_C(A) = \sigma_C(A_0, \eta).$$

(ii) For each  $\lambda \in \sigma_P(A_\#) = \sigma_P(A)$ ,

$$\text{Ker}(\lambda I - A_\#)^l = F \text{Ker}(\lambda I - A)^l, \quad l = 1, 2, \dots.$$

*In particular*

$$\mathcal{M}_\lambda^{T*} = F \mathcal{M}_\lambda.$$

Analogous inclusions to (8.4) involving the operator  $G$  are:

**Lemma 8.1.** For each  $\lambda \in \sigma_P(A^*) = \sigma_P(A_T)$  (resp.  $\lambda \in \sigma_P(A) = \sigma_P(A_\#)$ ),

$$\begin{aligned} G^* \text{Ker}(\lambda I - A^*)^l &\subset \text{Ker}(\lambda I - A_T)^l, \quad l = 1, 2, \dots \\ (\text{resp. } G \text{Ker}(\lambda I - A_\#)^l &\subset \text{Ker}(\lambda I - A)^l, \quad l = 1, 2, \dots). \end{aligned}$$

*Proof.* This lemma can be proved similarly as in the proof of (8.4) by using (4.26) and (4.27) instead of (4.30).

**Theorem 8.2.** (i)

$$(8.7) \quad \sigma_d(A^*) = \sigma_d(A_T).$$

(ii)

$$(8.8) \quad G^* \text{Ker}(\lambda I - A^*)^l = \text{Ker}(\lambda I - A_T)^l,$$

$$(8.9) \quad \dim \text{Ker}(\lambda I - A^*)^l = \dim \text{Ker}(\lambda I - A_T)^l = \dim \text{Ker } \mathcal{A}_l^T(\lambda) < \infty, \\ l = 1, 2, \dots, \lambda \in \sigma_d(A_T).$$

*In particular*

$$(8.10) \quad \dim \mathcal{M}_\lambda^* = \dim \mathcal{M}_\lambda^T \quad \text{for } \lambda \in \sigma_d(A^*) = \sigma_d(A_T).$$

Proof. (i). Let  $\lambda \in \sigma_d(A^*)$ . Then  $\lambda \in \sigma_0(A^*)$  and from Kato [25, p. 184] and Theorem 7.1, we have

$$(8.11) \quad \bar{\lambda} \in \sigma_d(A) \text{ is a pole of } R(\mu; A) \text{ of order } k_{\bar{\lambda}} \text{ at } \mu = \bar{\lambda}, \\ \mathcal{M}_{\bar{\lambda}} = \text{Ker}(\bar{\lambda} I - A)^{k_{\bar{\lambda}}}, \quad \dim \mathcal{M}_{\bar{\lambda}}^* = \dim \mathcal{M}_{\bar{\lambda}} < \infty,$$

$$(8.12) \quad \sup_{l \geq 1} \dim \text{Ker}(\bar{\lambda} I - A)^l = \dim \mathcal{M}_{\bar{\lambda}} < \infty.$$

Since  $G$  is one to one and

$$(8.13) \quad G \text{Ker}(\bar{\lambda} I - A^*)^l \subset \text{Ker}(\bar{\lambda} I - A)^l, \quad l = 1, 2, \dots$$

for  $\bar{\lambda} \in \sigma_d(A) \subset \sigma_p(A) = \sigma_p(A^*)$  (by Lemma 8.1), it follows from (8.11)–(8.13) that  $\sup_{l \geq 1} \dim \text{Ker}(\bar{\lambda} I - A^*)^l < \infty$ . This implies  $\dim \mathcal{M}_{\bar{\lambda}}^{T*} < \infty$ , and hence  $\bar{\lambda} \in \sigma_d(A^*)$ .

Again by using Kato [25, p. 184] we have  $\lambda = \bar{\bar{\lambda}} \in \sigma_d(A_T)$ , which proves  $\sigma_d(A^*) \subset \sigma_d(A_T)$ . The reverse inclusion  $\sigma_d(A_T) \subset \sigma_d(A^*)$  is obtained similarly as above. Thus, (8.7) is shown.

(ii). For  $\lambda \in \sigma_d(A_T)$ , the space  $\text{Ker}(\lambda I - A_T)^l$  is finite dimensional and is invariant under the semigroup  $S_T(t)$ . Then the operator  $S_T(h) = G^* F^*$  in (4.13) is bijective on  $\text{Ker}(\lambda I - A_T)^l$  (cf. Proposition 7.4). Consequently by (8.2),

$$\text{Ker}(\lambda I - A_T)^l = G^* F^* \text{Ker}(\lambda I - A_T)^l = G^* \text{Ker}(\lambda I - A^*)^l.$$

Since  $G^*$  is one to one and  $\dim \text{Ker}(\lambda I - A_T)^l < \infty$ , we have (8.9) by (8.8) and Theorem 7.2 (ii).

**Corollary 8.2.** (i)  $\sigma_d(A^*) = \sigma_d(A)$ .

(ii)

$$G \text{Ker}(\lambda I - A^*)^l = \text{Ker}(\lambda I - A)^l, \\ \dim \text{Ker}(\lambda I - A^*)^l = \dim \text{Ker}(\lambda I - A)^l = \dim \text{Ker } \mathcal{A}_l(\lambda) < \infty, \\ l = 1, 2, \dots, \lambda \in \sigma_d(A).$$

Lastly in this subsection we give an  $M_p$ -adjoint result for  $A$  which is an

immediate consequence from Kato [25, p. 184].

**Theorem 8.3.** *Let  $\lambda$  be a pole of  $R(\mu; A_0, \eta)$  of order  $k_\lambda$  at  $\mu=\lambda$ . Then  $\bar{\lambda}$  is a pole of  $R(\mu; A_0^*, \eta^*)$  of same order  $k_\lambda$  at  $\mu=\bar{\lambda}$ . Furthermore*

$$\mathcal{M}_\lambda^* = \text{Ker } (\bar{\lambda}I - A^*)^{k_\lambda} = (P_\lambda)^* M_p^*, \quad \dim \mathcal{M}_\lambda = \dim \mathcal{M}_\lambda^* \text{ (may be infinity),}$$

where the adjoint  $(P_\lambda)^*$  of  $P_\lambda$  in (7.13) is given explicitly by

$$(P_\lambda)^* = \frac{1}{2\pi i} \int_{\bar{\Gamma}_\lambda} R(\mu; A^*) d\mu = P_{\bar{\lambda}}^*$$

with  $\bar{\Gamma}_\lambda$  the miller image of  $\Gamma_\lambda$ .

The same result for  $A_T$  holds, but we omit to give such a representation.

8.2. Representations of spectral projections

It was shown in the proof of Theorem 8.2 that if  $\lambda \in \sigma_d(A)$ , then  $\bar{\lambda} \in \sigma_d(A^*) = \sigma_d(A_T)$  and

$$(8.14) \quad \dim \mathcal{M}_\lambda = \dim \mathcal{M}_\lambda^* = \dim \mathcal{M}_\lambda^T = d_\lambda < \infty .$$

Let  $\Phi = \{\phi_1, \dots, \phi_{d_\lambda}\}$  and  $\Psi = \{\psi_1, \dots, \psi_{d_\lambda}\}$  be the bases of  $\mathcal{M}_\lambda$  and  $\mathcal{M}_\lambda^T$  respectively. Let  $M$  be a  $d_\lambda \times d_\lambda$  matrix of element  $m_{ij} = \langle \phi_i, F^* \psi_j \rangle_{M_p}$ . Then by (8.14) and (8.3),  $M$  is nonsingular. Hence we can suppose

$$(8.15) \quad \langle \phi_j, F^* \psi_i \rangle_{M_p} = \langle \phi_i, \psi_j \rangle = \delta_{ij}, \quad i, j = 1, \dots, d_\lambda,$$

where  $\langle \cdot, \cdot \rangle$  denotes the hereditary pairing in (4.35). Now we introduce the continuous projection operator

$$\tilde{P}_\lambda g = \sum_{i=1}^{d_\lambda} \langle g, F^* \psi_i \rangle_{M_p} \phi_i, \quad g \in M_p .$$

It is easily verified that  $\text{Im } \tilde{P}_\lambda = \mathcal{M}_\lambda$  and  $\text{Ker } \tilde{P}_\lambda = \text{Im } (\lambda I - A)^{k_\lambda}$ , so that  $\tilde{P}_\lambda = P_\lambda$ . Thus, we obtain the following desired result.

**Theorem 8.4.** *Let  $\lambda \in \sigma_d(A)$ . Then the spectral projection  $P_\lambda$  in (7.13) has the following equivalent representation*

$$(8.16) \quad \tilde{P}_\lambda g = \sum_{i=1}^{d_\lambda} \langle g, \psi_i \rangle \phi_i, \quad g \in M_p,$$

where  $\{\phi_1, \dots, \phi_{d_\lambda}\}$  is a basis of  $\mathcal{M}_\lambda$  and  $\{\psi_1, \dots, \psi_{d_\lambda}\}$  is a basis of  $\mathcal{M}_\lambda^T$  satisfying (8.15).

**Corollary 8.4.** *Let  $\lambda \in \sigma_d(A_T)$ . Then the spectral projection  $P_\lambda^T$  associated with  $A_T$  has the following equivalent representation*

$$\bar{P}_\lambda^T f = \sum_{i=1}^{d'_\lambda} \langle \phi_i, f \rangle \psi_i, \quad f \in M_p^*,$$

where  $d'_\lambda = \dim \mathcal{M}_\lambda^T = \dim \mathcal{M}_\lambda$ ,  $\{\psi_1, \dots, \psi_{d'_\lambda}\}$  is a basis of  $\mathcal{M}_\lambda^T$  and  $\{\phi_1, \dots, \phi_{d'_\lambda}\}$  is a basis of  $\mathcal{M}_\lambda$  satisfying the same condition in (8.15).

**9. Completeness of generalized eigenfunctions**

The problem of completeness of generalized eigenfunctions of retarded FDE's has been studied by Delfour and Manitius [15] and Manitius [29] for  $n$ -dimensional equations. The purpose of this section is to extend some of their results to infinite dimensional case.

First we give characterizations of the kernels of spectral projections  $P_\lambda$  and others in terms of  $F, H_\lambda$  and the retarded resolvent. Let  $\lambda \in \sigma_0(A)$  and  $P_\lambda$  be the projection in (7.13). Then by (7.15), (7.16) and (7.18), we have that

$$\begin{aligned} g \in \text{Ker } P_\lambda & \quad \text{if and only if} \quad Q_\lambda^n g = ((\lambda I - A)P_\lambda)^n g = 0, \quad n = 1, \dots, k_\lambda \\ & \quad \text{if and only if} \quad R(\mu; A)g \text{ is holomorphic (h.l.) at } \mu = \lambda. \end{aligned}$$

Since  $E_\lambda, H_\lambda$  and  $T_\lambda$  are operator valued entire functions, the equality

$$(9.1) \quad \begin{aligned} \{g \in M_p; R(\mu; A)g \text{ is h.l. at } \mu = \lambda\} \\ = \{g \in M_p; R(\mu; A_0, \eta)H_\mu Fg \text{ is h.l. at } \mu = \lambda\} \end{aligned}$$

follows from the representation (6.11) of  $R(\mu; A)$ . Hence  $\text{Ker } P_\lambda$  is given by the right hand side of (9.1). Let  $P_\lambda^*$  and  $P_\lambda^{T*}$  denote the spectral projections associated with  $A^*$  and  $A_\tau^*$ , respectively. Then the next proposition follows from the representations of  $R(\mu; A^*)$  and other resolvents given in Theorems 6.1 and 6.2 as above. Note that  $[R(\mu; A^*)f]^0 = R(\mu; A_\tau^*, \eta^*)H_\mu f$ .

**Proposition 9.1.**

- (i)  $\text{Ker } P_\lambda = \{g \in M_p; R(\mu; A_0, \eta)H_\mu Fg \text{ is h.l. at } \mu = \lambda\}, \lambda \in \sigma_0(A)$ .
- (ii)  $\text{Ker } P_\lambda^T = \{f \in M_p^*; R(\mu; A_\tau^*, \eta^*)H_\mu F^*f \text{ is h.l. at } \mu = \lambda\}, \lambda \in \sigma_0(A_\tau)$ .
- (iii)  $\text{Ker } P_\lambda^* = \{f \in M_p^*; R(\mu; A_\delta^*, \eta^*)H_\mu f \text{ is h.l. at } \mu = \lambda\}, \lambda \in \sigma_0(A^*)$ .
- (iv)  $\text{Ker } P_\lambda^{T*} = \{g \in M_p; R(\mu; A_0, \eta)H_\mu g \text{ is h.l. at } \mu = \lambda\}, \lambda \in \sigma_0(A_\tau^*)$ .

For notational brevity we set

$$\mathcal{M} = \bigcup_{\lambda \in \sigma_p(A)} \mathcal{M}_\lambda, \quad \mathcal{M}^T = \bigcup_{\lambda \in \sigma_p(A_\tau)} \mathcal{M}_\lambda^T, \quad \mathcal{M}^* = \bigcup_{\lambda \in \sigma_p(A^*)} \mathcal{M}_\lambda^*, \quad \mathcal{M}^{T*} = \bigcup_{\lambda \in \sigma_p(A_\tau^*)} \mathcal{M}_\lambda^{T*}.$$

DEFINITION 9.1. The systems of generalized eigenfunctions of  $A, A^*, A_\tau$  and  $A_\tau^*$  are said to be complete if

$$C1(\mathcal{M}) = M_p, \quad C1(\mathcal{M}^*) = M_p^*, \quad C1(\mathcal{M}^T) = M_p^*, \quad C1(\mathcal{M}^{T*}) = M_p,$$

respectively.

For a set  $E \subset C^1$ ,  $\bar{E}$  denotes the miller image of  $E$ . Following the consideration in preceding sections, we know that

$$(9.2) \quad \begin{aligned} \overline{\sigma_0(A)} &= \sigma_0(A^*) = \sigma_0(A_T) = \sigma_0(A_\delta^*, \eta^*), \\ \overline{\sigma_0(A_T)} &= \sigma_0(A_\delta^*) = \sigma_0(A) = \sigma_0(A_0, \eta). \end{aligned}$$

**Proposition 9.2.** (i) *If  $\sigma_P(A) = \sigma_0(A_0, \eta)$ , then*

$$(9.3) \quad \mathcal{M}^\perp = \{f \in M_p^*: R(\lambda; A_\delta^*, \eta^*)H_\lambda f \text{ is h.l. on } \rho(A_\delta^*, \eta^*) \cup \sigma_0(A_\delta^*, \eta^*)\},$$

$$(9.4) \quad (\mathcal{M}^{T*})^\perp = \{f \in M_p^*: R(\lambda; A_\delta^*, \eta^*)H_\lambda F^*f \text{ is h.l. on } \rho(A_\delta^*, \eta^*) \cup \sigma_0(A_\delta^*, \eta^*)\},$$

(ii) *If  $\sigma_P(A_T) = \sigma_0(A_\delta^*, \eta^*)$ , then*

$$(9.5) \quad (\mathcal{M}^T)^\perp = \{g \in M_p; R(\lambda; A_0, \eta)H_\lambda g \text{ is h.l. on } \rho(A_0, \eta) \cup \sigma_0(A_0, \eta)\},$$

$$(9.6) \quad (\mathcal{M}^*)^\perp = \{g \in M_p; R(\lambda; A_0, \eta)H_\lambda Fg \text{ is h.l. on } \rho(A_0, \eta) \cup \sigma_0(A_0, \eta)\},$$

Proof. We shall prove only (9.3). Other equalities are proved quite analogously. Using the duality theorem and assumption, we have by the first relation in (9.2) that

$$(9.7) \quad \begin{aligned} \mathcal{M}^\perp &= \bigcap_{\lambda \in \sigma_P(A)} \mathcal{M}_\lambda^\perp = \bigcap_{\lambda \in \sigma_0(A)} (\text{Im } P_\lambda)^\perp = \bigcap_{\lambda \in \sigma_0(A)} \text{Ker}(P_\lambda)^* \\ &= \bigcap_{\lambda \in \sigma_0(A)} \text{Ker } P_\lambda^* = \bigcap_{\lambda \in \sigma_0(A)} \text{Ker } P_\lambda^* = \bigcap_{\lambda \in \sigma_0(A^*)} \text{Ker } P_\lambda^*. \end{aligned}$$

Hence the equality (9.3) follows immediately from Proposition 9.1 (iii). See Theorem 8.3 for completeness in deriving (9.7). Hence the equality (9.3) follows immediately from Proposition 9.1 (iii).

Since  $C1(\mathcal{M}) = M_p$  if and only if  $\mathcal{M}^\perp = \{0\}$ , from Proposition 9.2 we have the following criteria for the completeness of generalized eigenfunctions.

**Theorem 9.1.** (i) *Assume that  $\sigma(A) = \sigma_0(A_0, \eta)$ . Then the system of generalized eigenfunctions of  $A$  (resp.  $A_\delta^*$ ) is complete if and only if*

$$(9.8) \quad \{f \in M_p^*: R(\lambda; A_\delta^*, \eta^*)H_\lambda f \text{ is entire}\} = \{0\}$$

$$(9.9) \quad (\text{resp. } \{f \in M_p^*: R(\lambda; A_\delta^*, \eta^*)H_\lambda F^*f \text{ is entire}\} = \{0\}).$$

(ii) *Assume that  $\sigma(A_T) = \sigma_0(A_\delta^*, \eta^*)$ . Then the system of generalized eigenfunctions of  $A_T$  (resp.  $A^*$ ) is complete if and only if*

$$(9.10) \quad \{g \in M_p; R(\lambda; A_0, \eta)H_\lambda g \text{ is entire}\} = \{0\}$$

$$(9.11) \quad (\text{resp. } \{g \in M_p; R(\lambda; A_0, \eta)H_\lambda Fg \text{ is entire}\} = \{0\}).$$

We now recall the definition of  $H_\lambda$  appearing in the conditions (9.8) and (9.10):



$$(9.12) \quad H_\lambda \phi = \phi^0 + \int_{-h}^0 e^{\lambda s} \phi^1(s) ds = c + q(\lambda),$$

for  $\phi \in M_p^*$  in (9.8) or for  $\phi \in M_p$  in (9.10), where  $c = \phi^0$ . The last term  $q(\lambda)$  in (9.12) is a finite Laplace transform of  $\phi^1 \in L_p(I_h; X^*)$  in the case (9.8) or of  $\phi^1 \in L_p(I_h; X)$  in the case (9.10). We denote these sets of all such functions by  $FLT_p^*$  and by  $FLT_p$ , respectively.

An additional property of  $H_\lambda$  is given by

$$\text{Lemma 9.1.} \quad \bigcap_{\lambda \in \mathbb{C}^1} \text{Ker } H_\lambda = \{0\}.$$

Proof. We shall show the case  $H_\lambda: M_p \rightarrow X$ . Let  $\phi \in \bigcap_{\lambda \in \mathbb{C}^1} \text{Ker } H_\lambda$ . Then by (9.12),

$$(9.13) \quad \phi^0 + \int_{-h}^0 e^{\lambda s} \phi^1(s) ds = 0 \quad \text{for all } \lambda \in \mathbb{C}^1.$$

Tending  $\text{Re } \lambda \rightarrow \infty$  in (9.13) we have  $\phi^0 = 0$ . Then  $\phi^1 = 0$  follows from (9.13) and the bijectivity of Laplace transform.

**Corollary 9.1.** *Assume that  $\sigma(A) = \sigma_0(A_0, \eta)$  (resp.  $\sigma(A_T) = \sigma_0(A_0^*, \eta^*)$ ). The system of generalized eigenfunctions of  $A$  (resp.  $A_T$ ) is complete if and only if for  $c \in X^*$  and  $q(\lambda) \in FLT_p^*$ ,*

$$R(\lambda; A_0^*, \eta^*)(c + q(\lambda)) \text{ is entire} \Rightarrow c + q(\lambda) \equiv 0$$

(resp. for  $c \in X$  and  $q(\lambda) \in FLT_p$ ,

$$R(\lambda; A_0, \eta)(c + q(\lambda)) \text{ is entire} \Rightarrow c + q(\lambda) \equiv 0).$$

Proof. Obvious from Theorem 9.1 and Lemma 9.1.

Corollary 9.1 is interpreted as that the completeness for  $A$  is equivalent to the nonexistence of nontrivial entire function in the class  $c + q(\lambda)$ ,  $c \in X^*$ ,  $q(\lambda) \in FLT_p^*$ , which completely cancel all poles of  $R(\lambda; A_0^*, \eta^*)$  in the form  $R(\lambda; A_0^*, \eta^*)(c + q(\lambda))$ , provided that  $R(\lambda; A_0^*, \eta^*)$  has poles only.

## 10. Illustrative examples

We shall give some applications of the abstract results of preceding sections to practical partial FDE's in the following examples.

EXAMPLE 10.1. Consider the parabolic partial FDE

$$(10.1) \quad \frac{\partial x(t, \xi)}{\partial t} = \frac{\partial}{\partial \xi} \left( a(\xi) \frac{\partial x(t, \xi)}{\partial \xi} \right) + b(\xi)x(t, \xi) + \sum_{r=1}^m a_r(\xi)x(t-h_r, \xi) \\ + \int_{-h}^0 a_I(s, \xi)x(t+s, \xi) ds, \quad t > 0, \xi \in (0, 1)$$

with boundary and initial conditions

$$(10.2) \quad x(t, 0) = x(t, 1) = 0, \quad t > 0,$$

$$(10.3) \quad x(0, \xi) = g^0(\xi), \quad x(s, \xi) = g^1(s, \xi) \quad \text{a.e. } (s, \xi) \in I_h \times [0, 1].$$

For the system (11.1)–(11.3) we assume

(i)

$$(10.4) \quad a(\xi) > 0 \quad \text{for } \xi \in [0, 1], \quad a(\cdot) \in C^1[0, 1], \quad b(\cdot) \in C[0, 1];$$

(ii)

$$(10.5) \quad a_r(\cdot) \in L_2[0, 1], \quad r = 1, \dots, m, \quad 0 \leq h_1 < \dots < h_m = h, \\ a_r(\cdot) \in L_2(I_h \times [0, 1]);$$

(iii)

$$(10.6) \quad g = (g^0(\cdot), g^1(\cdot)) \in L_2[0, 1] \times L_2(I_h \times [0, 1]).$$

The product space  $L_2[0, 1] \times L_2(I_h \times [0, 1])$  in (10.6) can be identified with  $L_2[0, 1] \times L_2(I_h; L_2[0, 1])$ , so we denote this space simply by  $M_2$ . Let  $A_0$  be the realization in  $L_2[0, 1]$  of the Sturm-Liouville operator  $\partial/\partial\xi(a(\xi)\partial/\partial\xi) + b(\xi)$  with Dirichlet boundary condition (10.2). In what follows we shall write  $L_2$  instead of  $L_2[0, 1]$  for brevity. Since  $L_2$  is a Hilbert space, we identify  $L_2$  and  $L_2^*$  as usual. Now we define the operator  $A_r \in B(L_2)$ ,  $r = 1, \dots, m$  and  $A_r \in L_2(I_h; B(L_2))$  by

$$(A_r z)(\xi) = a_r(\xi)z(\xi) \quad \text{a.e. } \xi \in [0, 1], \quad r = 1, \dots, m$$

and

$$(A_r(s)z)(\xi) = a_r(s, \xi)z(\xi) \quad \text{a.e. } \xi \in [0, 1] \quad \text{for a.e. } s \in I_h,$$

respectively. By the condition (10.5) and the use of Schwartz inequality the above operators are well defined. Then the system (10.1)–(10.3) can be written in the same form as of (E) on the space  $X = L_2$ . The (weak) solution  $x(t, \xi; g)$  of (10.1)–(10.3) is interpreted as the mild solution  $x(t; g)(\xi)$  of (E) at the point  $\xi \in [0, 1]$ . So, for each  $t > 0$ ,  $x(t, \xi; g)$  has sense for a.e.  $\xi \in [0, 1]$ . Since  $A_0$  is selfadjoint with compact resolvent in  $L_2$  by (10.4), there exists a set of eigenvalues and eigenfunctions  $\{\mu_n, \Psi_n: n = 1, 2, \dots\}$  of  $A_0$  such that

(iv)  $\{\Psi_n\}$  is a complete orthonormal system in  $L_2$ ;

(v)  $\sqrt{-\mu_n} = Cn + O\left(\frac{1}{n}\right)$  as  $n \rightarrow \infty$ , where  $C$  is a constant depending only on the coefficient  $a(\xi)$  (cf. Kato [25, p. 277], Ince [23, p. 270–273]).

Consequently,  $A_0$  generates an analytic semigroup  $T(t)$  given by

$$T(t)z = \sum_{n=1}^{\infty} e^{\mu_n t} \langle z, \Psi_n \rangle_{L_2} \Psi_n, \quad z \in L_2,$$

where  $\langle z, \Psi_n \rangle_{L_2} = \int_0^1 z(\xi) \Psi_n(\xi) d\xi$ . Using the asymptotics of  $\{\mu_n\}$  in (v), we can verify that  $T(t)$  is compact for all  $t > 0$ . Then by Proposition 3.1 (ii),  $S(t)$  is compact for  $t > h$ . This implies by spectral mapping theorem (cf. Yosida [49, p. 277]) that  $\sigma(A) = \sigma_p(A) = \sigma_d(A)$  is countable, bounded from below and  $\sigma(A) \cap \{z: \alpha \leq \text{Re } z\}$  is a finite set for each  $\alpha \in \mathbf{R}^1$ . Now following the line of Hale's proof in [18, Chapter 7, Section 4] with some obvious modifications, we have the following result on the asymptotic stability.

The zero solution of (10.1)–(10.3) is exponentially asymptotic stable in  $L_2$ , i.e., there exist constants  $K \geq 1$  and  $\varepsilon_0 > 0$  such that

$$(10.7) \quad \int_0^1 |x(t, \xi; g)|^2 d\xi \leq K e^{-\varepsilon_0 t} \left( \int_0^1 |g^0(\xi)|^2 d\xi + \int_0^1 \int_{-h}^0 |g^1(s, \xi)|^2 d\xi ds \right),$$

$$t \geq 0, g \in M_2$$

provided that

$$(10.8) \quad \sup \{ \text{Re } \lambda : \lambda \in \sigma_d(A) \} < 0.$$

For the system (10.1)–(10.3), in view of Proposition 7.1, the condition (10.8) is replaced by that, for  $\text{Re } \lambda \geq 0$ ,

$$(10.9) \quad \lambda g^0 - \frac{\partial}{\partial \xi} \left( a(\xi) \frac{\partial g^0}{\partial \xi} \right) - b(\xi) g^0 - \sum_{r=1}^m e^{-\lambda h_r} \cdot a_r(\xi) g^0 - \int_{-h}^0 e^{\lambda s} a_I(s, \xi) g^0 ds = 0,$$

$$g^0 \in D(A_0) \text{ implies } g^0 = 0 \text{ in } L_2.$$

EXAMPLE 10.2. In this example, we consider the special equation

$$(10.10) \quad \frac{\partial x(t, \xi)}{\partial t} = a \frac{\partial^2 x(t, \xi)}{\partial \xi^2} + b x(t, \xi) + \sum_{r=1}^m a_r x(t - h_r, \xi), \quad t > 0, \xi \in (0, 1)$$

of (10.1) with the same mixed conditions (10.2) and (10.3), where  $a > 0$ ,  $b$ ,  $a_r$  are real constants. For the system (10.10), (10.2), (10.3) we have easily that

$$\mu_n = -an^2\pi^2 + b, \quad \Psi_n = \sqrt{2} \sin n\pi\xi, \quad n = 1, 2, \dots,$$

so that the spectrum  $\sigma(A)$  is given by

$$\sigma(A) = \{ \lambda \in \mathbf{C}^1 : \lambda + an^2\pi^2 - b - \sum_{r=1}^m a_r e^{-\lambda h_r} = 0 \text{ for some } n = 1, 2, \dots \}.$$

It is evident that  $\sigma(A)$  is countable and each  $\lambda \in \sigma(A)$  has finite multiplicity (may be  $\neq 1$ ). The asymptotic stability condition (10.9) is now reduced to a verifiable condition that

$$(10.11) \quad \left\{ \begin{array}{l} \text{all roots of the transcendental equations} \\ \lambda = -a n^2 \pi^2 + b + \sum_{r=1}^m a_r e^{-\lambda h_r}, \quad n = 1, 2, \dots \\ \text{have negative real parts.} \end{array} \right.$$

A simple sufficient condition for (10.11) is  $\sum_{r=1}^m |a_r| < a\pi^2 - b$ , which is shown by direct calculations using contradiction. Recently Lenhart and Travis [28, Corollary 1.2] have proved that (10.11) holds for all  $h_r \geq 0$  if and only if  $\sum_{r=1}^m |a_r| < a\pi^2 - b$  and  $\sum_{r=1}^m a_r < a\pi^2 - b$ . Set  $\Delta_n(\lambda) = \lambda + a n^2 \pi^2 - b - \sum_{r=1}^m a_r e^{-\lambda h_r}$ . Let  $\{\lambda_{nj}\}_{j=1}^{\infty}$  be the set of roots of  $\Delta_n(\lambda) = 0$  and let  $k_{nj}$  be the multiplicity of  $\lambda_{nj}$ . Then the retarded resolvent  $R(\lambda; A_0, \eta)$  is given by

$$R(\lambda; A_0, \eta)z = \sum_{n=1}^{\infty} \frac{2}{\Delta_n(\lambda)} \langle z, \sin n\pi\xi \rangle_{L_2} \sin n\pi\xi, \quad \lambda \in \mathbf{C}^1 - \{\lambda_{nj} : n, j = 1, 2, \dots\};$$

the basis of the generalized eigenspace  $\mathcal{M}_{\lambda_{nj}}$  corresponding to  $\lambda_{nj} \in \sigma(A)$  is given by

$$\{e^{\lambda_{nj}s} \sin n\pi\xi, \dots, s^{k_{nj}-1} e^{\lambda_{nj}s} \sin n\pi\xi\}.$$

If  $a_m \neq 0$ , this system of generalized eigenfunctions is complete in  $M_2$ . But if  $a_m = 0$ , the completeness does not hold in general (cf. [29, 35]).

#### Acknowledgments

I would like to express my sincere gratitude to Professor H. Tanabe for his helpful comments and suggestions in accomplishing this work. I have learned much about functional analysis and partial differential equations from his excellent books [42,43]. I also express my gratitude to the referee for his comments.

#### References

- [1] N.U. Ahmed and K.L. Teo: *Optimal control of distributed parameter systems*, North-Holland, New York, 1981.
- [2] H.T. Banks and J.A. Burns: *Hereditary control problems: numerical methods based on averaging approximations*, SIAM J. Control Optim. **16** (1978), 169-208.
- [3] V. Barbu, *Nonlinear semigroups and differential equations in Banach spaces*, Noordhoff International Publishing, Leyden, the Netherlands, 1976.
- [4] C. Bernier and A. Manitius: *On semigroups in  $R^n \times L^p$  corresponding to differential equations with delay*, Canad. J. Math. **30** (1978), 897-914.
- [5] J.B. Borisovic and A.S. Turbabin: *On the Cauchy problem for linear non-homo-*

- geneous differential equations with retarded argument*, Soviet Math. Doklady **10** (1969), 401–405.
- [6] J.A. Burns, T.L. Herdman and H.W. Stech: *Linear functional differential equations as semigroups on product spaces*, *SIAM J. Math. Anal.* **14** (1983), 98–116.
- [7] J.A. Burns and T.L. Herdman: *Adjoint semigroup theory for a class of linear functional differential equations*, *SIAM J. Math. Anal.* **7** (1976), 729–745.
- [8] R.F. Curtain and A.J. Pritchard: *Infinite dimensional linear systems theory*, Lec. Notes in Control & Inform. Sci. 8, Springer-Verlag, Berlin, Heidelberg, New York, 1978.
- [9] R. Datko: *Linear autonomous neutral differential equations in a Banach space*, *J. Differential Equations* **25** (1977), 258–274.
- [10] R. Datko: *Representation of solutions and stability of linear differential-difference equations in a Banach space*, *J. Differential Equations* **29** (1978), 105–166.
- [11] M.C. Delfour: *The linear quadratic optimal control problem for hereditary differential systems: Theory and numerical solutions*, *Appl. Math. Optim.* **3** (1977), 101–162.
- [12] M.C. Delfour: *Linear optimal control of systems with state and control variable delays*, *Automatica* **20** (1984), 69–77.
- [13] M.C. Delfour: *State theory of linear hereditary differential systems*, *J. Math. Anal. Appl.* **60** (1977), 8–35.
- [14] M.C. Delfour, E.B. Lee and A. Manitius: *F-reduction of the operator Riccati equations for hereditary differential systems*, *Automatica* **14** (1978), 385–395.
- [15] M.C. Delfour and A. Manitius: *The structural operator F and its role in the theory of retarded systems, Part I*, *J. Math. Anal. Appl.* **73** (1980), 466–490; *Part II*, *ibid.*, **74** (1980), 359–381.
- [16] P.A. Fuhrmann: *Linear systems and operators in Hilbert space*, McGraw-Hill International Book Company, New York, 1981.
- [17] J.S. Gibson: *Linear-quadratic optimal control of hereditary differential systems: Infinite dimensional Riccati equations and numerical approximations*, *SIAM J. Control. Optim.* **21** (1983), 95–139.
- [18] J.K. Hale: *Theory of functional differential equations*, Springer-Verlag, New York, Heidelberg, Berlin, 1977.
- [19] J.K. Hale and R. Meyer: *A class of functional equations of neutral type*, *Memoirs Amer. Math. Soc.* No. 76, 1967.
- [20] J.K. Hale and J. Kato: *Phase space for retarded equations with infinite delay*, *Funk. Ekvac.* **21** (1978), 11–41.
- [21] E. Hille and R.S. Phillips: *Functional analysis and semigroup*, *Amer. Math. Soc. Colloq. Publ.* XXXI, Providence, R.I., 1957.
- [22] C.S. Hönl: *Volterra Stieltjes-integral equations*, in “*Functional differential equations and bifurcation*”, A.F. Ise ed., 173–216, Springer-Verlag, New York, Heidelberg, Berlin, 1980.
- [23] E.L. Ince: *Ordinary differential equations*, Dover, New York, 1944.
- [24] F. Kappel and H.K. Wimmer: *An elementary divisor theory for autonomous linear functional differential equations*, *J. Differential Equations* **21** (1976), 134–147.
- [25] T. Kato: *Perturbation theory for linear operators*, Second Edi., Springer-Verlag, Berlin, Heidelberg, New York, 1976.

- [26] K. Kunisch and W. Schappacher: *Mild and strong solutions for partial differential equations with delay*, Ann. Mat. Pure Appl. **75** (1980), 193–219.
- [27] K. Kunisch and W. Schappacher: *Necessary conditions for partial differential equations with delay to generate  $C_0$ -semigroups*, J. Differential Equations **50** (1983), 49–79.
- [28] S. Lenhart and C.C. Travis: *Stability of functional partial differential equations*, J. Differential Equations **58** (1985), 212–227.
- [29] A. Manitius: *Completeness and  $F$ -completeness of eigenfunctions associated with retarded functional differential equations*, J. Differential Equations **35** (1980), 1–29.
- [30] A. Manitius: *Necessary and sufficient conditions of approximate controllability for linear retarded systems*, SIAM J. Control Optim. **19** (1981), 516–532.
- [31] A. Manitius:  *$F$ -controllability and observability of linear retarded systems*, Appl. Math. Optim. **9** (1982), 73–95.
- [32] R.K. Miller: *Nonlinear Volterra integral equations*, Benjamin, New York, 1971.
- [33] S. Nakagiri: *On the fundamental solution of delay-differential equations in Banach spaces*, J. Differential Equations **41** (1981), 349–368.
- [34] S. Nakagiri: *Optimal control of linear retarded systems in Banach spaces*, J. Math. Anal. Appl. **120** (1986), 169–210.
- [35] S. Nakagiri: *Pointwise completeness and degeneracy of functional differential equations in Banach spaces, I: General time delays*, J. Math. Anal. Appl. **127** (1987), 492–529.
- [36] T. Naito: *On autonomous linear functional differential equations with infinite retardations*, J. Differential Equations **21** (1976), 297–315.
- [37] R.S. Phillips: *The adjoint semi-group*, Pacific J. Math. **5** (1955), 269–283.
- [38] H. Royden: *Real analysis*, Macmillan, New York, 1963.
- [39] D. Salamon: *On controllability and observability of time delay systems*, IEEE Trans. Automat. Control **AC-29** (1984), 432–438.
- [40] D. Salamon: *Control and observation of neutral systems*, Research Notes in Mathematics **91**, Pitman, London, 1984.
- [41] A.E. Taylor and D.C. Lay: *Introduction to functional analysis*, Second Edi., John Wiley, New York, 1980.
- [42] H. Tanabe: *Equations of evolution*, Pitman, London, 1979.
- [43] H. Tanabe: *Functional analysis, I. II*, Jikkyo Shuppan Publ. Company, Tokyo, 1977, 1981 (in Japanese).
- [44] C.C. Travis and G.F. Webb: *Existence and stability for partial functional differential equations*, Trans. Amer. Math. Soc. **200** (1974), 395–418.
- [45] C.C. Travis and G.F. Webb: *Existence, stability and compactness in the  $\alpha$ -norm for partial functional differential equations*, Trans. Amer. Math. Soc. **240** (1978), 129–143.
- [46] R.B. Vinter: *On the evolution of the state of linear differential delay equations in  $M^2$ : Properties of a generator*, J. Inst. Math. Appl. **21** (1978), 13–23.
- [47] R.B. Vinter and R.H. Kwong: *The finite time quadratic control problem for linear systems with state and control delays: An evolution equation approach*, SIAM J. Control Optim. **19** (1981), 139–153.
- [48] G.F. Webb: *Functional differential equations and nonlinear semigroups in  $L^p$ -space*, J. Differential Equations **20** (1976), 71–89.

- [49] K. Yosida: Functional analysis, Third Edi., Springer-Verlag, Berlin, Heidelberg, New York, 1971.

Department of Applied Mathematics  
Faculty of Engineering  
Kobe University  
Kobe 657  
Japan