



Title	Treatment of anterior open bite in a patient with dentinogenesis imperfecta
Author(s)	Tsujimoto, Takayuki; Kurosaka, Hiroshi; Yamashiro, Takashi
Citation	AJO-D0 Clinical Companion. 2024, 4(4), p. 303-313
Version Type	AM
URL	https://hdl.handle.net/11094/97807
rights	This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Table I: Cephalometric measurements before and after treatment

Measurements	Japanese norms (Female Adult) $\pm$ SD	Pretreatment (24 years, 1 months)	Posttreatment (27 years, 3 months)
Angular (degrees)			
SNA	80.8 $\pm$ 3.6	75.5	75.5
SNB	77.9 $\pm$ 4.5	72.3	72.6
ANB	2.8 $\pm$ 2.4	3.3	2.9
FMA	30.5 $\pm$ 2.1	40.0	38.6
Gonial angle	122.1 $\pm$ 5.3	138.8	138.8
U1-FH	112.3 $\pm$ 8.3	121.7	118.0
IMPA	93.4 $\pm$ 6.8	89.5	92.2
IIA	123.6 $\pm$ 10.6	108.8	111.2
Linear (mm)			
N-Me	125.8 $\pm$ 5.0	132.0	129.5
N-ANS	58.7 $\pm$ 3.5	57.4	57.4
ANS-Me	72.7 $\pm$ 4.4	76.1	73.3
Overjet	3.1 $\pm$ 1.1	6.3	3.4
Overbite	3.3 $\pm$ 1.9	-2.8	2.0