

Title	On representations of direct products of finite solvable groups
Author(s)	Hikari, Michitaka
Citation	Osaka Journal of Mathematics. 1971, 8(2), p. 299-307
Version Type	VoR
URL	https://doi.org/10.18910/9792
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ON REPRESENTATIONS OF DIRECT PRODUCTS OF FINITE SOLVABLE GROUPS

MICHITAKA HIKARI

(Received January 28, 1971)

Let K be a field and π a finite group. We denote by $G_0(K\pi)$ the Grothendieck ring of $K\pi$. Let π_i be a finite group and M_i be finitely generated $K\pi_i$ -module, $i=1, 2$. Let us denote by $M_1 \# M_2$ the outer tensor product of M_1 and M_2 . We can define the natural ring homomorphism $\varphi: G_0(K\pi_1) \otimes G_0(K\pi_2) \rightarrow G_0(K(\pi_1 \times \pi_2))$ by putting $\varphi([M_1] \otimes [M_2]) = [M_1 \# M_2]$. In this paper we study the kernel and cokernel of φ .

1. Let π be a finite group, E a finite normal separable extension of K which is a splitting field of π , and $\mathcal{G}(E/K)$ the Galois group of E over K . Let N be an $E\pi$ -module with character χ and $\sigma \in \mathcal{G}(E/K)$. Then we define an $E\pi$ -module σN , the conjugate of N , as usual and denote its character by $\sigma\chi$. We denote the Schur index of N over K by $m_K(N)$.

Now, let π be the direct product of finite groups π_1 and π_2 , $\pi = \pi_1 \times \pi_2$. Let M_i be an irreducible $K\pi_i$ -module, $i=1, 2$, and denote an irreducible $E\pi_i$ -component of $M_i^E = M_i \otimes_K E$ by N_i , the character of N_i by ψ_i and the Galois group E over $K(\psi_i)$ by $\mathcal{H}_i = \mathcal{G}(E/K(\psi_i))$. Then, the following results can be found in [3].

(1) If $\sigma, \tau \in \mathcal{G}(E/K)$, then $\sigma N_1 \# \tau N_2$ is an irreducible $E[\pi_1 \times \pi_2]$ -module also and $m_K(N_1 \# N_2) = m_K(\sigma N_1 \# \tau N_2)$.

(2) $M_1 \# M_2$ is completely reducible. $M_1 \# M_2 = k(T_1 \oplus \cdots \oplus T_r)$, where the $\{T_i\}$ are nonisomorphic irreducible $K\pi$ -modules and $k = m_K(N_1)m_K(N_2)/m_K(N_1 \# N_2)$. The $\{T_i\}$ have common K -dimension s , where $s = m_K(N_1 \# N_2)(K(\psi_1, \psi_2): K(N_1 \# N_2: E))$.

(3) $M_1 \# M_2$ is an irreducible $K\pi$ -module if and only if the following conditions are satisfied:

(a) $m_K(N_1)m_K(N_2) = m_K(N_1 \# N_2)$.

(b) $\mathcal{G}(E/K) = \mathcal{H}_1\mathcal{H}_2$.

(c) $(K(\psi_1): K)(K(\psi_2): K) = (K(\psi_1, \psi_2): K)$.

(4) Let $\pi_1 = \pi_2$, $\pi = \pi_1 \times \pi_1$. Let M_1 be an irreducible $K\pi_1$ -module. Then $M_1 \# M_1$ is irreducible if and only if M_1 is an absolutely irreducible $K\pi_1$ -module.

Since for any irreducible $K[\pi_1 \times \pi_2]$ -module M we can find a unique irreducible $K\pi_i$ -module M_i , $i=1, 2$, satisfying $M_1 \# M_2 \oplus \cdots \oplus M$, the following is an immediate corollary to (3).

(5) We denote the order of a group π by $|\pi|$. Let Q be the field of rational numbers. If $(|\pi_1|, |\pi_2|)=1$, then

$$\varphi: G_0(Q\pi_1) \otimes G_0(Q\pi_2) \xrightarrow{\sim} G_0(Q[\pi_1 \times \pi_2]).$$

One aim of this paper is to study the converse to (5).

2. Hereafter we assume $\text{char. } K=0$.

Lemma 1. *If π_1 and π_2 are finite abelian groups, then $\text{Ker } \varphi=0$ and $\text{Coker } \varphi$ is torsion free.*

Proof. Since the Schur index of abelian groups is 1, then φ is a split map by (2). Q.E.D.

Let $j: \pi' \rightarrow \pi$ be a group homomorphism. Then we have the induction and restriction functors

$$\text{mod} - K\pi' \xrightleftharpoons[j_* = \text{res}]{j^* = (\cdot \otimes_{K\pi'} K\pi)} \text{mod} - K\pi,$$

and these functors induce the additive homomorphisms of Grothendieck rings,

$$G_0(K\pi') \xrightleftharpoons[j_*]{j^*} G_0(K\pi).$$

Let π'_i be a subgroup of π_i . Then the following diagram is commutative.

$$\begin{array}{ccccccc} \text{Ker } \varphi & \longrightarrow & G_0(K\pi_1) \otimes G_0(K\pi_2) & \xrightarrow{\varphi} & G_0(K[\pi_1 \times \pi_2]) & \longrightarrow & \text{Coker } \varphi \\ \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow \\ \text{Ker } \psi & \longrightarrow & G_0(K\pi'_1) \otimes G_0(K\pi'_2) & \xrightarrow{\psi} & G_0(K[\pi'_1 \times \pi'_2]) & \longrightarrow & \text{Coker } \psi \end{array}$$

Proposition 2. *For any finite groups π_1, π_2 , we have $\text{Ker } \varphi=0$.*

Proof. Since $\text{Ker } \psi=0$ for cyclic groups π'_1 and π'_2 , by the commutativity of the above diagram and the Artin's induction theorem, $\text{Ker } \varphi=0$. Q.E.D.
(But we can prove this proposition without the induction theorem.)

Now let π'_i be a normal subgroup of π_i . Then we have the exact sequence $1 \rightarrow \pi'_i \xrightarrow{j} \pi_i \xrightarrow{p} \pi'_i \rightarrow 1, i=1, 2$. From this we obtain the following commutative diagram.

$$\begin{array}{ccccccc} G_0(K\pi'_1) \otimes G_0(K\pi'_2) & \xrightarrow{\varphi_1} & G_0(K[\pi'_1 \times \pi'_2]) & \xrightarrow{\varphi'_1} & \text{Coker } \varphi_1 \\ \updownarrow p^* \updownarrow p_* & & \updownarrow p^* \updownarrow p_* & & \updownarrow p^* \updownarrow p_* \\ G_0(K\pi_1) \otimes G_0(K\pi_2) & \xrightarrow{\varphi_2} & G_0(K[\pi_1 \times \pi_2]) & \xrightarrow{\varphi'_2} & \text{Coker } \varphi_2 \\ \updownarrow j^* \updownarrow j_* & & \updownarrow j^* \updownarrow j_* & & \updownarrow j^* \updownarrow j_* \\ G_0(K\pi'_1) \otimes G_0(K\pi'_2) & \xrightarrow{\varphi_3} & G_0(K[\pi'_1 \times \pi'_2]) & \xrightarrow{\varphi'_3} & \text{Coker } \varphi_3. \end{array}$$

Let M be an irreducible $K[\pi'_1 \times \pi'_2]$ -module, E a finite normal separable extension of K which is a splitting field of $\pi'_1 \times \pi'_2$ and $N_1 \# N_2$ an $E[\pi'_1 \times \pi'_2]$ -irreducible component of M^E , where N_i is the $E\pi'_i$ -irreducible module, $i=1, 2$. Denote the characters of M, N_i by χ, ψ_i respectively and put $m = |\pi'_1 \times \pi'_2|$.

Lemma 3. (a) *If there exists an irreducible $K[\pi'_1 \times \pi'_2]$ -module M such that $\varphi'_3([M]) \neq 0$ and*

$$m_K(N_1)m_K(N_2)(K(\psi_1):K)(K(\psi_2):K)/m_K(N_1 \# N_2)(K(\psi_1, \psi_2):K) \nmid m,$$

then $\text{Coker } \varphi_2 \neq 0$.

(b) *If there exists an irreducible $K[\pi'_1 \times \pi'_2]$ -module M such that $\varphi'_3([M]) \neq 0$ and the inertial group of $\chi, I(\chi) = \{g \mid g \in \pi_1 \times \pi_2, \chi^g = \chi\}$, coincides with $\pi_1 \times \pi_2$ and if $\text{Coker } \varphi_3$ is torsion free, then $\text{Coker } \varphi_2 \neq 0$.*

(c) *Let $K = \mathbb{Q}$. Let π'_i be an elementary abelian p -group and $|\pi'_i| = p^{n_i}, i=1, 2$, where p is an odd prime. Denoting by c_i the centralizer of π'_i in π_i , then we can regard π_i/c_i as a group of morphisms of the module π'_i . This identification induces the natural map*

$$\psi: \pi_1 \times \pi_2 \longrightarrow \pi_1/c_1 \times \pi_2/c_2 \longrightarrow PGL(n_1 + n_2, p).$$

Then $j_ j^* \text{Coker } \varphi_3 = 0$ if and only if*

$$\begin{array}{c} \psi(\pi_1 \times \pi_2) \text{ contains} \\ \sigma = \begin{pmatrix} \overbrace{1 \dots 1}^{n_1} & & \overbrace{1 \dots 1}^{n_2} \\ & \ddots & \\ & & 1 \\ & & & r \\ & & & & \ddots \\ & & & & & r \end{pmatrix}, \end{array}$$

where r is a primitive root modulo p and the order of σ is $p-1$.

(d) *If $\text{Coker } \varphi_1 \neq 0$, then $\text{Coker } \varphi_2 \neq 0$.*

Proof. (a) Assume $j_* j^* [M] = [M \otimes_{K[\pi'_1 \times \pi'_2]} K[\pi_1 \times \pi_2]] \in \text{Im } \varphi_3$. Then

$$M \otimes_{K[\pi'_1 \times \pi'_2]} K[\pi_1 \times \pi_2] = M_{11} \# M_{21} \oplus M_{12} \# M_{22} \oplus \dots \oplus M_{1s} \# M_{2s}$$

where each M_{ij} is a $K\pi'_i$ -irreducible module, $i=1, 2, j=1, 2, \dots, s$.

$$(*) \quad M^E \otimes_{E[\pi'_1 \times \pi'_2]} E[\pi_1 \times \pi_2] = M_{11}^E \# M_{21}^E \oplus M_{12}^E \# M_{22}^E \oplus \dots \oplus M_{1s}^E \# M_{2s}^E.$$

Let N_{ij} be an $E\pi'_i$ -irreducible component of $M_{ij}^E, i=1, 2, j=1, 2, \dots, s$. Since $N_1 \# N_2$ is an irreducible component of M^E , there exists an element g_{ij} of π_i and $\sigma_i \in \mathcal{G}(E/K)$ such that $N_{ij} = (\sigma_i N_i) g_{ij}$. Let ψ_{ij} be the character of N_{ij} . Then $m_K(N_{ij}) = m_K((\sigma_i N_i) g_{ij}) = m_K(N_i)$ and $K(\psi_{ij}) = K(\psi_i)$. Comparing the E -dimensions of both sides in $(*)$, we obtain

$$m_K(N_1 \# N_2)(K(\psi_1, \psi_2): K)m(N_1 \# N_2: E) \\ = s \cdot m_K(N_1)m_K(N_2)(K(\psi_1): K)(K(\psi_2): K)(N_1 \# N_2: E).$$

Hence

$$m = s \cdot m_K(N_1)m_K(N_2)(K(\psi_1): K)(K(\psi_2): K)/m_K(N_1 \# N_2)(K(\psi_1, \psi_2): K).$$

This contradicts the assumption. Therefore $\text{Coker } \varphi_2$ is not zero.

(b) Since $I(X) = \pi_1 \times \pi_2$, $M \otimes_{K[\pi_1' \times \pi_2'] } K[\pi_1 \times \pi_2] \cong M^m$ as $K[\pi_1' \times \pi_2']$ -modules. Since $\text{Coker } \varphi_3$ is torsion free, we have $\text{Coker } \varphi_2 \neq 0$.

(c) First, assume $j_* j^* \text{Coker } \varphi_3 = 0$. We have $Q\pi_1' \cong Q[X_1, \dots, X_{n_1}]/(X_1^p - 1, \dots, X_{n_1}^p - 1)$ and $Q\pi_2' \cong Q[Y_1, \dots, Y_{n_2}]/(Y_1^p - 1, \dots, Y_{n_2}^p - 1)$. Let ζ be a primitive p -th root of unity and put $G = \mathcal{G}(Q(\zeta)/Q)$. Further put $M_1 = Q[X_1, \dots, X_{n_1}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta)^G$ and $M_2 = Q[Y_1, \dots, Y_{n_2}]/(Y_1 - \zeta, \dots, Y_{n_2} - \zeta)^G$ where $()^G$ is the set of all G -invariant elements of $()$. Then each M_i is an irreducible $Q\pi_i'$ -module.

$$M_1 \# M_2 \cong Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta, Y_1 - \zeta, \dots, Y_{n_2} - \zeta)^G \\ \oplus Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta, Y_1 - \zeta^2, \dots, Y_{n_2} - \zeta^2)^G \\ \oplus \dots \\ \oplus Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta, Y_1 - \zeta^{p-1}, \dots, Y_{n_2} - \zeta^{p-1})^G$$

as $Q[\pi_1' \times \pi_2']$ -modules. If we put

$$M = Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta, Y_1 - \zeta, \dots, Y_{n_2} - \zeta)^G,$$

we have $\varphi_3([M]) \neq 0$ and so, by the assumption, $j_* j^* M \oplus > M_1 \# M_2$. Therefore we can find an element c of $\pi_1 \times \pi_2$ such that

$$M \otimes c = Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta, Y_1 - \zeta, \dots, Y_{n_2} - \zeta)^G \otimes c \\ \cong Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta, \dots, X_{n_1} - \zeta, Y_1 - \zeta^r, \dots, Y_{n_2} - \zeta^r)^G.$$

Then we have $\psi(c) = \sigma$.

Conversely, assume $\psi(\pi_1 \times \pi_2) \ni \sigma$. Let c be a representative of σ in $\pi_1 \times \pi_2$, $\{g_i, g_i c, g_i c^2, g_i c^3, \dots, g_i c^{p-2}\}$ representatives of $\pi_1' \times \pi_2'$ in $\pi_1 \times \pi_2$ and M an irreducible $Q[\pi_1' \times \pi_2']$ -module. (We can find representatives of above type.) Then $j_* j^* M = \sum_i^{\oplus} (M \otimes g_i \oplus M \otimes g_i c \oplus \dots \oplus M \otimes g_i c^{p-2})$ and there exist integers $r_1, \dots, r_{n_1}, t_1, \dots, t_{n_2}$ such that $M \otimes g_i \cong Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta^{r_1}, \dots, X_{n_1} - \zeta^{r_{n_1}}, Y_1 - \zeta^{t_1}, \dots, Y_{n_2} - \zeta^{t_{n_2}})^G$. By the assumption, $\sum_{j=0}^{p-2} M \otimes g_i c^j \cong \sum_{j=1}^{p-1} Q[X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}]/(X_1 - \zeta^{r_1}, \dots, X_{n_1} - \zeta^{r_{n_1}}, Y_1 - \zeta^{j t_1}, \dots, Y_{n_2} - \zeta^{j t_{n_2}})^G \cong [Q[X_1, \dots, X_{n_1}]/(X_1 - \zeta^{r_1}, \dots, X_{n_1} - \zeta^{r_{n_1}})^G \# Q[Y_1, \dots, Y_{n_2}]/(Y_1 - \zeta^{t_1}, \dots, Y_{n_2} - \zeta^{t_{n_2}})^G]^u$ where u is a positive integer. Therefore $[j_* j^* M] \in \text{Im } \varphi_3$ and

$$\varphi'_3(j_* j^*[M])=0.$$

(d) Since $p^* p_* = 1$, it is trivial. Q.E.D.

Denote by $e(\pi)$ the exponent of a group π and by ζ_n a primitive n -th root of unity for any integer n .

Lemma 4. *Let π_i be an abelian group, $i=1, 2$, and $G.C.D.(e(\pi_1), e(\pi_2)) = \prod p^{h_p}$. Let $s_p = \max \{s \mid \zeta_p^s \in K\}$ for each prime p . If there exists at least one prime p such that $h_p > s_p$, then $\varphi: G_0(K\pi_1) \otimes G_0(K\pi_2) \not\rightarrow G_0(K[\pi_1 \times \pi_2])$.*

Proof. $K(\zeta_{p^{h_p}})$ is an irreducible $K\pi_i$ -module. Let us consider the underlying abelian group of $K(\zeta_{p^{h_p}}) \# K(\zeta_{p^{h_p}})$. There exists an integer n such that $K(\zeta_{p^{h_p}}) \otimes_K K(\zeta_{p^{h_p}}) \cong K(\zeta_{p^{h_p}})^n$. Since $(K(\zeta_{p^{h_p}}): K) \neq 1$, we have $n \neq 1$ and so $\text{Coker } \varphi \neq 0$. Q.E.D.

3. (I) We can determine $\text{Coker } \varphi$ when π_1 and π_2 are abelian groups. Let π_1 be an abelian group with invariants l_1, \dots, l_n and π_2 an abelian group with invariants l_{n+1}, \dots, l_{n+m} . Then

$$\begin{aligned} \text{rank Coker } \varphi = & \sum_{d_i \mid l_i} [\eta(d_1) \times \dots \times \eta(d_{n+m}) \times \{(K(\zeta_{\substack{L.C.M.(d_i) \\ 1 \leq i \leq n+m}}}: K)^{-1} \\ & - (K(\zeta_{\substack{L.C.M.(d_i) \\ 1 \leq i \leq n}}}: K)^{-1} (K(\zeta_{\substack{L.C.M.(d_{n+j}) \\ 1 \leq j \leq m}}}: K)^{-1}\}] \end{aligned}$$

where η is the Euler's function.

(II) We denote the center of a group π by $Z(\pi)$.

Theorem 5. *Let $L.C.M.(e(Z(\pi_1/\pi'_1))) = \prod p^{m_p}$, $L.C.M.(e(Z(\pi_2/\pi'_2))) = \prod p^{n_p}$ and $s_p = \max \{s \mid \zeta_p^s \in K\}$. If there exists a prime p such that $\min(m_p, n_p) > s_p$, then $G_0(K\pi_1) \otimes G_0(K\pi_2) \not\rightarrow G_0(K[\pi_1 \times \pi_2])$.*

Proof. By assumption, there exists a normal subgroup π'_i of π_i such that $p^{m_p} \mid e(Z(\pi_1/\pi'_1))$ and $p^{n_p} \mid e(Z(\pi_2/\pi'_2))$. Put $\pi''_i = \pi_i/\pi'_i$ and consider the following commutative diagram;

$$\begin{array}{ccccccc} G_0(K\pi_1) \otimes G_0(K\pi_2) & \xrightarrow{\varphi_1} & G_0(K[\pi_1 \times \pi_2]) & \longrightarrow & \text{Coker } \varphi_1 \\ \updownarrow & & \updownarrow & & \updownarrow \\ G_0(K\pi'_1) \otimes G_0(K\pi'_2) & \xrightarrow{\varphi_2} & G_0(K[\pi'_1 \times \pi'_2]) & \longrightarrow & \text{Coker } \varphi_2 \\ \updownarrow & & \updownarrow & & \updownarrow \\ G_0(K[Z(\pi''_1)]) \otimes G_0(K[Z(\pi''_2)]) & \xrightarrow{\varphi_3} & G_0(K[Z(\pi''_1 \times \pi''_2)]) & \longrightarrow & \text{Coker } \varphi_3 \end{array}$$

Let $G.C.D.(e(Z(\pi''_1)), e(Z(\pi''_2))) = \prod p^{h_p}$. Since $h_p > s_p$, $\text{Coker } \varphi_3 \neq 0$ by Lemma 4 and since $\text{Coker } \varphi_3$ is torsion free by Lemma 1, then $\text{Coker } \varphi_2 \neq 0$ by Lemma 3 (b), and therefore $\text{Coker } \varphi_1 \neq 0$ by Lemma 3 (d). Q.E.D.

Corollary 6. *Let L.C.M. $(e(Z(\pi/\pi'))) = \prod_{\pi' \triangleleft \pi} p^{m_p} = h$. Then any splitting field of π contains the primitive h -th root of unity.*

Proof. By (4) $G_0(K\pi) \otimes G_0(K\pi) \xrightarrow{\sim} G_0(K[\pi \times \pi])$ if and only if K is a splitting field of π . So this corollary is trivial. Q.E.D.

(III) **Theorem 7.** *Let π_i be a group of odd order. Assume that there exists an odd prime p such that $p \mid (|\pi_1|, |\pi_2|)$ and $2 \mid (K(\zeta_p): K)$ where ζ_p is a primitive p -th root of unity. Then $\varphi: G_0(K\pi_1) \otimes G_0(K\pi_2) \xrightarrow{\sim} G_0(K[\pi_1 \times \pi_2])$.*

Proof. Since π_i is a group of odd order, each π_i is solvable. We can consider a principal series $\pi_i = \pi_i^{(0)} \supset \pi_i^{(1)} \supset \dots \supset \pi_i^{(n_i)} \supset \dots \supset (1)$ and find integers n_i, r_i such that $|\pi_i^{(n_i)} : \pi_i^{(n_i+1)}| = p^{r_i}, r_i > 0$, for each $i=1, 2$. And consider the following commutative diagram;

$$\begin{array}{ccc}
 G_0(K\pi_1) \otimes G_0(K\pi_2) & \xrightarrow{\varphi_1} & \\
 \updownarrow & & \\
 G_0(K[\pi_1/\pi_1^{(n_1+1)}]) \otimes G_0(K[\pi_2/\pi_2^{(n_2+1)}]) & \xrightarrow{\varphi_2} & \\
 \updownarrow & & \\
 G_0(K[\pi_1^{(n_1)}/\pi_1^{(n_1+1)}]) \otimes G_0(K[\pi_2^{(n_2)}/\pi_2^{(n_2+1)}]) & \xrightarrow{\varphi_3} & \\
 \updownarrow & & \\
 G_0(K[\pi_1 \times \pi_2]) & \xrightarrow{\quad} & \text{Coker } \varphi_1 \\
 \updownarrow & & \updownarrow \\
 G_0(K[\pi_1/\pi_1^{(n_1+1)}] \times \pi_2/\pi_2^{(n_2+1)}) & \xrightarrow{\quad} & \text{Coker } \varphi_2 \\
 \updownarrow & & \updownarrow \\
 G_0(K[\pi_1^{(n_1)}/\pi_1^{(n_1+1)}] \times \pi_2^{(n_2)}/\pi_2^{(n_2+1)}) & \xrightarrow{\quad} & \text{Coker } \varphi_3.
 \end{array}$$

By Lemma 4, $\text{Coker } \varphi_3 \neq 0$. Since

$$(K(\zeta_p): K)(K(\zeta_p): K)/(K(\zeta_p): K) \nmid \prod_{i=1, 2} |\pi_i : \pi_i^{(n_i)}|,$$

from Lemma 3 (a) it follows that $\text{Coker } \varphi_2 \neq 0$ and so by Lemma 3 (d) we have $\text{Coker } \varphi_1 \neq 0$. Q.E.D.

In case $2 \nmid |\pi_1| \cdot |\pi_2|$, we can prove the converse to (5) by putting $K=Q$ in Theorem 7.

Corollary 8 *Assume $2 \nmid |\pi_1| \cdot |\pi_2|$. Then*

$$\varphi: G_0(Q\pi_1) \otimes G_0(Q\pi_2) \xrightarrow{\sim} G_0(Q[\pi_1 \times \pi_2]) \text{ if and only if } (|\pi_1|, |\pi_2|) = 1.$$

Corollary 9. *Put $|\pi| = \prod_{i=1}^m p_i^{e_i}$ and suppose that $p_i \nmid p_j - 1$ for any indices $1 \leq i, j \leq m$. Then any splitting field of π contains the primitive $p_1 \cdots p_m$ -th root of unity.*

Proof. We can show this corollary by the same method as in Theorem 7. Q.E.D.

REMARK. If π is a nilpotent group, this result has been seen. For a given integer $n=p_1^{n_1} \cdots p_m^{n_m}$ all of groups of order n are nilpotent if and only if $p_j \nmid p_i^{n_i-t} - 1$ for all t such that $n_i > t \geq 0$ and all i, j .


(IV) Here we consider 2-groups. In this case the groups with a cyclic subgroup of index 2 are important. For any character of 2-groups is induced by the character of such groups. (See [4] p. 73 (14.3).) Such groups can be classified as follows. Put $|\pi| = 2^{n+1}$,

- I $\pi = \langle s \mid s^{2^{n+1}} = 1 \rangle$.
- II $\pi = \langle s, t \mid s^{2^n} = 1, t^2 = 1, tst^{-1} = s \rangle$
- III $\pi = \langle s, t \mid s^{2^n} = 1, t^2 = s^{2^{n-1}}, tst^{-1} = s^{-1} \rangle, \quad n \geq 2$.
- IV $\pi = \langle s, t \mid s^{2^n} = 1, t^2 = 1, tst^{-1} = s^{-1} \rangle, \quad n \geq 2$.
- V $\pi = \langle s, t \mid s^{2^n} = 1, t^2 = 1, tst^{-1} = s^{1+2^{n-1}} \rangle, \quad n \geq 3$.
- VI $\pi = \langle s, t \mid s^{2^n} = 1, t^2 = 1, tst^{-1} = s^{-1+2^{n-1}} \rangle, \quad n \geq 3$.

Theorem 10. Let π_1 and π_2 be arbitrary two groups of the above types. Then $\varphi: G_0(Q\pi_1) \otimes G_0(Q\pi_2) \xrightarrow{\sim} G_0(Q[\pi_1 \times \pi_2])$ if and only if

- (a) π_1 is a group of type (I, $n=0$), (II, $n=1$) or (IV, $n=2$) and π_2 is any,
- (b) π_1 is of type (I, $n=1$), (II, $n=2$), (III, $n=2$), (V, $n=3$) or (VI, $n=3$) and π_2 is of type IV,
- (c) π_1 is of type (I, $n=1$), (II, $n=2$) or (V, $n=3$) and π_2 is of type VI.

Let $Q_k = Q(\cos \pi/2^{k-1} + i \sin \pi/2^{k-1})$,
 $R_k = Q(\cos \pi/2^{k-1})$ and $S_k = Q(i \sin \pi/2^{k-1})$.



First, we shall write out the division algebras which are contained within $Q\pi$. (See, Feit [4] p. 63-p. 66.)

If π is of type I, $\{Q_i\}_{1 \leq i \leq n+1}$ are all of the division algebras of $Q\pi$. When π is of type II, $\{Q_i\}_{1 \leq i \leq n}$ are all of the division algebras. If π is of type III, then $\{D, R_i\}_{1 \leq i \leq n-1}$ are all of the division algebras where D is the division algebra of a faithful irreducible representation of π . Hence the center of D is R_n . If π

is of type IV, $\{R_i\}_{1 \leq i \leq n}$ are all of the division algebras. When π is of type V and $n=3$, then Q_1 and Q_2 are only division algebras of π . If $n>3$, Q_3 is one of the division algebras of π . And if π is of type VI, $\{S_n, R_i\}_{1 \leq i \leq n-1}$ are all of the division algebras.

Lemma 11. *Let χ be a faithful irreducible character of the group of type III. Then $m_{S_k}(\chi)=1$ for $k \geq 2$.*

In case $k=2$, we can see the proof of Lemma 11, for example, in Feit [4]. In case $k>2$, we can prove it similarly.

Proof of Theorem 10. a) When π_1 is of type (I, $n=0$), (II, $n=1$) or (IV, $n=2$), Q is a splitting field of π_1 . Therefore φ is an isomorphism.

(b) If π_i is of type I, II or V, Q_2 is one of the division algebras of π_i , $i=1, 2$. Then $\text{Coker } \varphi \neq 0$, because $Q_2 \otimes_Q Q_2 \cong Q_2 \oplus Q_2$.

(c) If π_1 is of type I, II or V and π_2 of type III, then $\text{Coker } \varphi \neq 0$ because $Q_2 \otimes_Q D \cong (Q_2)_2$.

(d) If π_1 is of type (I, $n=1$), (II, $n=2$) or (V, $n=3$) and π_2 is of type IV, the division algebra of π_1 is Q_1 or Q_2 and the division algebra of π_2 is one of $\{R_i\}_{1 \leq i \leq n}$. Since $Q_2 \otimes_Q R_i \cong Q_i$ for $3 \leq i \leq n$ and $Q_2 \otimes_Q R_i \cong Q_2$ for $i < 3$, we obtain $\text{Coker } \varphi = 0$. If π_1 is of type (I, $n>1$), (II, $n>2$) or (V, $n>3$) and π_2 of type IV, $\text{Coker } \varphi \neq 0$, because Q_3 is one of the division algebras of π_1 and $Q_3 \otimes_Q R_3 \cong Q_3 \oplus Q_3$.

(e) If π_1 is of type (I, $n=1$), (II, $n=2$) or (V, $n=3$) and π_2 is of type VI, $\text{Coker } \varphi = 0$. For the division algebra of π_2 is one of $\{S_n, R_i\}_{1 \leq i \leq n-1}$, $n \geq 3$ and $Q_2 \otimes_Q S_n \cong Q_n$ for $n \geq 3$, $Q_2 \otimes_Q R_i \cong Q_i$ for $3 \leq i \leq n-1$ and $Q_2 \otimes_Q R_i \cong Q_2$ for $i < 3$. If π_1 is of type (I, $n>1$), (II, $n>2$) or (V, $n>3$) and π_2 is of type VI, then $\text{Coker } \varphi \neq 0$ because $Q_3 \otimes_Q S_3 \cong Q_3 \oplus Q_3$ and $Q_3 \otimes_Q R_3 \cong Q_3 \oplus Q_3$.

(f) If π_1 is of type (III, $n=2$), the division algebra of π_1 is Q_1 or D with center Q . Since $D \otimes_Q R_i$ is a division algebra also for all i , we have $\text{Coker } \varphi = 0$ if π_2 is of type IV. If $n>2$, there exists a division algebra of π_1 with center R_3 . From the fact that $R_3 \otimes_Q R_3 \cong R_3 \oplus R_3$, it follows that $\text{Coker } \varphi \neq 0$ for the group π_2 of type IV.

(g) Assume that π_1 is of type III and π_2 of type VI. Since $D \otimes_Q S_n \cong (S_n)_2$ by Lemma 11, we obtain $\text{Coker } \varphi \neq 0$.

(h) Suppose that π_1 is of type IV. If π_2 is of type (VI, $n=3$), the division algebra of π_2 is Q_1 or S_3 . Since $R_i \otimes_Q S_3 \cong Q_i$ for $3 \leq i \leq n$ or S_3 for $i < 3$, $\text{Coker } \varphi = 0$. If π_2 is of type (VI, $n=4$), then S_4 is a division algebra of π_2 and if π_2 is of type (VI, $n>4$), R_4 is a division algebra of π_2 . Since $R_3 \otimes_Q S_4 \cong S_4 \oplus S_4$ and $R_3 \otimes_Q R_4 \cong R_4 \oplus R_4$, in both cases $\text{Coker } \varphi \neq 0$. Q.E.D.

References

- [1] H. Bass: *Algebraic K-Theory*, Benjamin, 1968.
- [2] C. W. Curtis and I. Reiner: *Representation Theory of Finite Groups and Associative Algebras*, Interscience, 1962.
- [3] B. Fein: *Representations of direct products of finite groups*, Pacific J. Math. **20** (1967), 45-58.
- [4] W. Feit: *Characters of Finite Groups*, Benjamin, 1967.
- [5] W. Feit and J. G. Thompson: *Solvability of groups of odd order*, Pacific J. Math. **13** (1963), 775-1029.
- [6] L. Solomon: *The representation of finite groups in algebraic number fields*, J. Math. Soc. Japan **13** (1961), 144-164.

