



Title	Kikuchi Disease After SARS-CoV-2 Vaccination: A Case Report With Immunohistochemical Analyses
Author(s)	Hamamoto, Yuichiro; Kawamura, Michihiro; Mori, Hideo et al.
Citation	International Journal of Surgical Pathology. 2024, 32(6), p. 1123-1128
Version Type	AM
URL	https://hdl.handle.net/11094/98830
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Tables

Table 1 Eight cases of Kikuchi disease after SARS-CoV-2 vaccination

No.	Reference	Age	Sex	Symptoms	Detail of vaccine	Period until onset
1	Betancur et al. ⁴	40s	F	axillary lymphadenopathy	SARS-CoV-2 BNT162b2 mRNA vaccine	3 months
2	Ikeda et al. ⁵	20s	F	fever and painful cervical lymphadenopathy	not available	0 day
3	Kashiwada et al. ⁶	27	F	fever and axillary lymphadenitis	SARS-CoV-2 BNT162b2 mRNA vaccine	3 months
4	Tan et al. ⁷	18	F	fever and axillary lymphadenopathy	SARS-CoV-2 BNT162b2 mRNA vaccine	34 days
5	Tan et al. ⁷	34	M	fever and axillary lymphadenopathy	SARS-CoV-2 BNT162b2 mRNA vaccine	16 days
6	Soub et al. ⁸	18	M	fever, neck swelling, anorexia, and nausea	SARS-CoV-2 BNT162b2 mRNA vaccine	10 days
7	Daghri et al. ⁹	24	F	fever and cervical lymphadenopathy	not available	1 month
8	Guan et al. ¹⁰	36	M	fever and cervical lymphadenopathy	Sinopharm-inactivated SARS-CoV-2 vaccine	6 days

F, female; M, male