

Title	Artinian rings related to relative almost projectivity. II
Author(s)	Harada, Manabu
Citation	Osaka Journal of Mathematics. 1995, 32(3), p. 577-589
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9910">https://doi.org/10.18910/9910</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## ARTINIAN RINGS RELATED TO RELATIVE ALMOST PROJECTIVITY II

MANABU HARADA

(Received November 8, 1993)

Let  $R$  be an artinian ring. In [10], we have studied  $R$  on which the following condition holds: for  $R$ -modules  $M$  and  $N$ , if  $M$  is  $N$ -projective, then  $M'$  is always almost  $N$ -projective for every submodule  $M'$  of  $M$ . If  $M'$  is always  $N$ -projective in the above, then this property characterizes hereditary rings with  $J^2=0$  [2] and [6], where  $J$  is the Jacobson radical of  $R$ .

We have investigated the above condition in [10], when i);  $M$  and  $N$  are local and ii):  $M$  is local and  $N$  is a direct sum of local modules. In this paper we give a characterization of  $R$  over which the above condition is satisfied for any  $R$ -modules  $M$  and  $N$ .

### 1. Preliminaries

In this paper  $R$  is always an artinian ring with identity, and every module is a finitely generated  $R$ -module. We shall use the same notations given in [10].

We have studied rings  $R$  over which the following condition is satisfied in [10]:

For any  $R$ -modules  $M$  and  $N$

- (#) if  $M$  is  $N$ -projective, then  $M'$  is always almost  $N$ -projective for every submodule  $M'$  of  $M$ .

We denote primitive idempotents in  $R$  by  $e, f, g$ , and so on. Assume that (#) holds whenever  $M$  and  $N$  are local. Then we have shown in [10] that  $R$  has the following structure:

- (0)  $J^3=0$  and for a primitive idempotent  $e$  with  $eJ^2 \neq 0$   
 $eR \supset eJ \simeq \sum_k \oplus (f_k R)^{(n_k)} \oplus \sum_j \oplus S_j$ ,  
 where the  $f_i R$  is a uniserial and projective module with  $f_i J^2=0$ ,  $f_i J \neq 0$   
 and the  $S_j$  is simple.

(If necessary, we use the following decomposition:

$$e_i R \supset e_i J \simeq \sum_{k(1)} \oplus (f_{ik} R)^{(n_{ik})} \oplus \sum_{j(1)} \oplus S_{ij}.)$$

We shall use frequently the following theorem: [10], Theorem 1.

**Theorem 0.** *Let  $R$  be artinian. Then (#) holds whenever  $M$  and  $N$  are local*

if and only if i)  $J^3=0$  and  $(eR \supset) eJ$  has the decomposition (0), ii) if  $e_1R \not\approx e_2R$ , then  $f_{1i}R \not\approx f_{2j}$  for any  $i$  and  $j$ , iii)  $fR/fJ$  is never isomorphic to any simple component of  $\text{Soc}(R)$  and iv) for any simple submodule  $S$  in  $\sum_K \oplus (f_k R)^{(n_k)}$ ,  $eReS = \sum_{K'} \oplus (f_k J)^{(n_k)}$  for some  $K' \subset K$ , where  $e$  and  $f$  are in (0).

## 2. Condition(#)

In this section we study (#) where  $M$  and  $N$  are any finitely generated  $R$ -modules. In this case  $gJ$  is almost projective for any primitive idempotent  $g$ , and hence  $R$  is a right almost hereditary ring [7], i.e.  $J$  is almost projective as a right  $R$ -module.

First we give

**Proposition 1.** Let  $R$  be right almost hereditary. Then the following are equivalent :

- 1) (#) holds whenever  $M$  and  $N$  are local.
- 2) (#) holds whenever  $M$  is local and  $N$  is a finite direct sum of local modules.

Proof. 1)  $\rightarrow$  2). Since  $R$  is almost hereditary,  $\text{Soc}(R)$  is almost projective by definition and Theorem 0. If  $gR/gJ$  is monomorphic to  $\text{Soc}(R)$ , then  $gR$  is uniserial by [9], Theorem 1 and we have 2) by [10], Theorem 2.

Next we study (#) when  $M$  is local and  $N$  is any  $R$ -module. We use the decomposition (0) of  $eJ$ . Put  $f_1R = fR$  and  $S = \text{Soc}(fR) = fJ$ .

**Lemma 1.** Assume that (#) holds whenever  $M$  is local and let  $eR \supset fR \supset S$  be as above. Then 1) every element in  $\text{Hom}_R(S, S)$  is extensible to an element in  $\text{Hom}_R(fR, fR)$ , and 2)  $S$  is neither isomorphic to any simple component of  $\text{Soc}(gR)$  nor any  $S_j$  in (0), where  $gR \not\approx eR$ ,  $gR \not\approx fR$  and  $gJ \neq 0$ .

Proof. Assume that  $\text{Soc}(gR)$  contains a simple component isomorphic to  $S$  via  $\theta$  for some primitive idempotent  $g$ . Take  $fR \oplus gR$  and its submodule  $\tilde{S} = \{s + \theta(s) | s \in S\}$  and put  $N = (fR \oplus gR) / \tilde{S}$ . If  $gR \not\approx eR$ ,  $eR/S$  is  $N$ -projective by [1], p. 22, Exercise 4 and [10], Lemma 6. Hence  $fR/S$  is almost  $N$ -projective by (#). However  $fR/S$  is not  $N$ -projective by [1], p. 22, Exercise 4. Therefore  $N$  is decomposable by [3], Theorem 1. Let  $N = N_1 \oplus N_2$ . Suppose  $gR \not\approx fR$ . Then we can assume  $N_1/J(N_1) \approx fR/fJ$  and  $N_2/J(N_2) \approx gR/gJ$ . Further  $N = \tilde{f}R + \tilde{g}R$ , where  $\tilde{f}R \approx fR$  and  $\tilde{g}R \approx gR$ . Since  $|N| = |\tilde{f}R| + |\tilde{g}R| - 1$ , we obtain a)  $N_1 \approx \tilde{f}R$  or b)  $N_2 \approx \tilde{g}R$  via the projections. In a)  $N = \tilde{f}R \oplus N_2$ , and hence  $\theta^{-1}$  is extensible to an element in  $\text{Hom}_R(gR, fR)$ , and in b)  $N = N_1 \oplus \tilde{g}R$ , and hence  $\theta$  is extensible to an element in  $\text{Hom}_R(fR, gR)$ . We obtain the similar result even if  $gR \approx fR$ . Hence from the above observation we obtain 1) and that  $S$  is never isomorphic to

any simple component of  $\text{Soc}(e'R)$  and  $\text{Soc}(f'R)$  form Theorem 0, where  $e'R \not\approx eR$ ,  $f'R \not\approx fR$  and  $e'J^2 \neq 0$ ,  $f'J \neq 0$ . Finally assume  $S \approx S_1 \subset eR$ , where  $S_1$  is a simple module in (0). Take  $(fR \oplus eR/(fJ)^{(n_1)})/\tilde{S}$ . Since  $eRe(fJ) \subset (fJ)^{(n_1)}$ ,  $eR/S$  is  $(fR \oplus eR)/(\tilde{S} \oplus (fJ)^{(n_1)})$ -projective. Similarly to the above we obtain an extension of  $\theta$  (or  $\theta^{-1}$ ) in  $\text{Hom}_R(fR, eR/(fJ)^{(n_1)})$  (or in  $\text{Hom}_R(eR/(fJ)^{(n_1)}, fR)$ ). However there are no extensions of  $\theta$  by Theorem 0, a contradiction.

If  $R$  is left QF-2 in the above, then any element in  $\text{Hom}_R(S, S)$  is extensible to an element in  $\text{Hom}_R(eR, eR)$  by [10], Lemma 13, however this fact is not true in lemma 1 (see Proposition 3 and Example 2 below). Under the assumption (#) we can state the content of Lemma 1 as follows :

*let  $S'$  be a simple submodule in  $gR$ , then any element in  $\text{Hom}_R(S, S')$  is extensible to an element in  $\text{Hom}_R(fR, gR)$ , where  $g$  is any primitive idempotent.*

Finally we study (#) for any  $R$ -modules  $M$  and  $N$ . We start with studying a structure of  $N$ . Let  $eR \supset eJ \approx \sum \oplus (f_i R)^{(n_i)} \oplus \dots$  be as in (0). We consider the condition :

(\*) the properties in Lemma 1 and Theorem 0 hold.

We fix primitive idempotents  $e$  and  $f = f_i$  above. Take a projective module  $T$  and put  $T = (eR)^{(p)} \oplus (fR)^{(q)} \oplus \sum g_j R$ , where  $g_j R \not\approx eR$  and  $g_j R \not\approx fR$  for all  $j$ .

**Lemma 2.** *Assume (\*). Let  $T$ ,  $e$  and  $f$  be as above. If  $T/C$  is indecomposable, then  $T = fR$  or  $q = 0$ .*

*Proof.* Assume  $q \geq 1$ . Consider the decomposition  $T = (fR) \oplus T'$ , where  $T' = (eR)^{(p)} \oplus (fR)^{(q-1)} \oplus \sum \oplus g_j R$ , and use the same notations as in [10], Lemma 10 for this decomposition. We may assume  $C \subset J(T)$  (note that  $T$  is a lifting module). Suppose  $C_1 = 0$  and  $C^1 = fJ$ . Put  $C^1 = xR$ ;  $xk = x$  for a primitive idempotent  $k$  and let  $\theta: C^1 = C^1/C_1 \approx C^2/C_2 \subset J(T')/C_2$  be the isomorphism. Then  $\theta(x) = x_1 + x_2 + x_3 + C_2$ , where  $x_1 \in (eJ)^{(p)}$ ,  $x_2 \in (fJ)^{(q-1)}$ ,  $x_3 \in \sum \oplus g_j J$  and  $x_i k = x_i$  for all  $i$ . Since  $x_1 J = 0$  for all  $i$  from iii) in Theorem 0,  $\theta: xR \approx (x_1 + x_2 + x_3)R \subset T'$ , i.e.,  $(x_1 + x_2 + x_3)R \oplus C_2 = C^2$ . Therefore  $\theta$  is extensible to an element  $\theta'$  in  $\text{Hom}_R(fR, T')$  by the properties in Lemma 1 (note  $(x_1 + x_2 + x_3)R \subset x_1 R \oplus x_2 R x_3 R$ ). As a consequence  $T = T' \oplus fR(\theta') \supset C = C \cap T' \oplus C \cap fR(\theta')$ , provided  $C_1 = 0$ . If  $C_1 \neq 0$ ,  $C_1 = C^1$ , and hence  $T = fR$ , since  $T/C$  is indecomposable. Accordingly we know that if  $T/C$  is indecomposable,  $T = fR$  or  $q = 0$ .

We consider following modules :  $Z = (fR)^{(m)} \oplus fJ$  and  $U = V \oplus fJ$ , where  $V$  is a submodule of  $(fR)^{(m)}$ . Similarly to Lemma 2 we have

**Lemma 3.** *Assume iii) in Theorem 0. Let  $U$  be as above and  $X$  a*

submodule of  $U$ . Then  $X \subset V$  or  $U = V \oplus fJ(\theta)$  and  $X \supset fJ(\theta)$ , where  $\theta \in \text{Hom}_R(fJ, V)$ .

More generally we consider  $Z^* = (fR)^{(p)} \oplus (fJ)^{(q)}$  and  $U^* = V^* \oplus (fJ)^{(q)}$ , where  $V^*$  is a submodule of  $(fR)^{(p)}$ .

**Corollary.** Assume iii) in Theorem 0. Let  $U^*$  be as above and  $X^*$  a submodule of  $U^*$ . Then we obtain the following decomposition of  $U^*$ :  $U^* = V^* \oplus (fJ)^{(q')} \oplus Y_1 \oplus \cdots \oplus Y_{q-q'}$ , and  $X^* \supset Y_1 \oplus \cdots \oplus Y_{q-q'}$ , where  $Y_i \approx fJ$  for all  $i$ ,  $X^* \subset V^* \oplus Y_1 \oplus \cdots \oplus Y_{q-q'}$ ,  $Z^* = (fR)^{(p)} \oplus (fJ)^{(q')} \oplus Y_1 \oplus \cdots \oplus Y_{q-q'}$  and  $U^* = V^* \oplus (fJ)^{(q')} \oplus Y_1 \oplus \cdots \oplus Y_{q-q'}$ .

This corollary means that there exists an automorphism  $\sigma$  of  $Z^*$  such that  $Z^* = (fR)^{(p)} \oplus (fJ)^{(q')} \oplus (fJ)^{(q-q')} \supset \sigma(U^*) = V^* \oplus (fJ)^{(q')} \oplus (fJ)^{(q-q')} \supset \sigma(X^*) \oplus (fJ)^{(q-q')}$ .

We shall denote the above situation by the diagram :

$$(1) \quad \begin{array}{ccccccc} & & & p & & q-q' & & q' \\ Z^* & & fR & \xrightarrow{\quad} & fJ & \xrightarrow{\quad} & fJ & \\ & \cup & & & & & & \\ U^* & & V^* & \xrightarrow{\quad} & fJ & \xrightarrow{\quad} & fJ & \\ & \cup & & & & & & \\ X^* & & X' & \xrightarrow{\quad} & fJ & \xrightarrow{\quad} & 0 & \end{array}$$

Next we study a structure of a submodule  $M'$  of an  $R$ -module  $M$  under  $(*)$ . Let  $P$  be a projective cover of  $M$ , i.e.,

$$(2) \quad M \approx P/Q \text{ and } M' \approx P'/Q \text{ for some submodule } P' \text{ of } P.$$

Then we have a decomposition of  $P$  such that  $P = P_1 \oplus P_2$ ,  $P' = P_1 \oplus P_2 \cap P'$  and  $P'_2 = P_2 \cap P' \subset J(P_2)$ .

Let

$$(2') \quad P_2 = (e_1 R)^{(a_1)} \oplus (e_2 R)^{(a_2)} \oplus \cdots \oplus (f_{11} R)^{(b_{11})} \oplus \cdots \oplus (f_{1s_1} R) \oplus (f_{21} R)^{(b_{21})} \oplus \cdots \oplus (f_{2s_2} R)^{(b_{2s_2})} \oplus \cdots \oplus \Sigma \oplus gR, \text{ where the } e_i, \text{ the } f_i \text{ are given in (0) and } gJ^2 = 0 \text{ (} e_i R \not\approx e_j R, f_{ik} R \not\approx f_{js} R \text{ if } i \neq j \text{ and } gR \not\approx f_{ik} R \text{ for all } i).$$

Consider  $J(P_2)$  and rearrange it as follows :

$$J(P_2) = (D_{11})^{(a_1)} \oplus (f_{11} J)^{(b_{11})} \oplus (D_{12})^{(a_1)} \oplus (f_{12} J)^{(b_{12})} \oplus \cdots \oplus (D_{1s_1})^{(a_1)} \oplus (f_{1s_1} J)^{(b_{1s_1})} \oplus (D_{21})^{(a_2)} \oplus (f_{21} J)^{(b_{21})} \oplus \cdots \oplus \Sigma \oplus gJ, \text{ where } D_{ij} = (f_{ij} R)^{(n_{ij})}.$$

Put  $E_{ij'} = D_{ij}^{(a_i)} \oplus (f_{ij})^{(b_{ij})}$  and  $F = \Sigma \oplus gJ$ . Then from  $(*)$  we know that any simple sub-factor modules of  $E_{ij'}$  are not isomorphic to any ones of  $E_{st'}$  and  $F$  for

$\{i, j\} \neq \{s, t\}$ . Hence we obtain

$$(3) \quad \begin{aligned} P'_2 &= \Sigma \oplus P'_{2ij} \oplus P_0 \\ Q^2 &= \Sigma \oplus Q^2_{ij'} \oplus Q^2_0 \text{ and} \\ Q_2 &= \Sigma \oplus Q^2_{2ij} \oplus Q_{20} \text{ (see [10], Lemma 10 for } Q_1 \text{ and } Q^2), \end{aligned}$$

where  $E'_{ij} \supset P'_{2ij} \supset Q^2_{ij'} \supset Q^2_{2ij}$  and  $F \supset P_0 \supset Q^2_0 \supset Q_{20}$ . We may observe  $E'_{11} \supset P'_{211} \supset Q^2_{11'} \supset Q^2_{211}$  for the fixed (1,1) without loss of generality. From Corollary to Lemma 3 we have

$$(4) \quad \begin{array}{c} E'_{11} \\ E'_{211} \\ Q^2_{11'} \\ Q_{211'} \end{array} \quad \begin{array}{c} D_{11} \\ P_{211} \\ Q^2_{11} \\ Q_{211} \end{array} \quad \begin{array}{ccccccc} a_1 & d_1 & c_1 & b_1 & a_0 & & \\ \hline f_{11}J & f_{11}J & f_{11}J & f_{11}J & f_{11}J & & \\ \cup & & & & & & \\ f_{11}J & f_{11}J & f_{11}J & 0 & & & \\ \cup & & & & & & \\ f_{11}J & f_{11}J & 0 & 0 & 0 & & \end{array} ,$$

where  $b_{11} = a_0 + b_1 + c_1 + d_1$ .

Next we observe  $D_{11} \supset P_{211} \supset Q^2_{11} \supset Q_{211}$ . We put  $f_{11} = f$ . Then from [4], Lemma 5

$$P_{211} \quad \begin{array}{ccccccc} & e_1 & & & e_2 & e_3 & \\ \hline fR & & fJ & & 0 & & \end{array}$$

where  $e_1 + e_2 + e_3 = a_{11}$ .

Further from Corollary to Lemma 3 we have

$$(5) \quad \begin{array}{c} P_{211} \\ Q^2_{11} \\ Q_{211} \end{array} \quad \begin{array}{c} fR \\ H \\ I \end{array} \quad \begin{array}{ccccccc} e_1 & h_1 & g_1 & e'_2 & & & \\ \hline fJ & fJ & fJ & fJ & 0 & & \\ \cup & & & & & & \\ fJ & 0 & 0 & & & & \end{array} ,$$

where  $e_2 = h_1 + g_1 + e'_2$ .

We observe the left side of the above diagram :  $(fR)^{(e_1)} \supset H \supset I$ . From (\*) and [4], Lemma 5 we have

$$\begin{array}{c} (fR)^{(e_1)} (=E) \\ H \end{array} \quad \begin{array}{ccccccc} i_1 & i_2 & e'_1 & & & & \\ \hline fR & fR & fR & & & & \\ & fR & fJ & 0 & & & \end{array} ,$$

where  $e_1 = i_1 + i_2 + e'_1$ .

We apply Lemma 3 to  $I$  and  $H = H_1 \oplus fJ$ , where  $H_1 = (fR)^{(i_1)} \oplus (fJ)^{(i_2-1)}$ . Then  $I \subset H_1$  of  $I \supset fJ(h)$  for some  $h \in \text{Hom}_R(fJ, H_1) \subset \text{Hom}_R(fJ, E_1)$ , where  $E_1 =$

$((fR)^{(i_1)} \oplus (fR)^{(i_2-1)} \oplus fR \subset E$ . From  $(*)$   $h$  is extensible to  $\tilde{h} \in \text{Hom}_R(fR, E)$ . Hence  $E = E_1 \oplus fR(\tilde{h}) \supset H_1 \oplus fJ(h)$  and  $I \supset fJ(h)$ . Repeating this argument we may assume

$$(6) \quad \begin{array}{ccccccc} & i_1 & j_2' & i_2' & e_1' & & \\ E & \underline{fR} & \underline{fR} & \underline{fR} & \underline{fR} & & \\ H & \underline{fR} & \underline{fJ} & \underline{fJ} & \underline{0} & & \\ & \cup & & & & & \\ I & \underline{I'} & \underline{fJ} & \underline{0} & \underline{0} & & \end{array},$$

where  $i_2 = i_1 + i_2'$ .

Applying again [4], Lemma 5 to the left corner of the above diagram we have finally

$$(7) \quad \begin{array}{ccccccccccc} & k_1 & k_2 & i_1' & j_1 & i_2' & e_1' & h_1 & g_1 & & \\ P_2' & \underline{fR} & \underline{fR} & \underline{fR} & \underline{fR} & \underline{fR} & \underline{fR} & \underline{fJ} & \underline{fJ} & & \\ Q^2 & \underline{fR} & \underline{fR} & \underline{fR} & \underline{fJ} & \underline{fJ} & \underline{0} & \underline{fJ} & \underline{fJ} & & \\ Q_2 & \underline{fR} & \underline{fJ} & \underline{0} & \underline{fJ} & \underline{0} & \underline{0} & \underline{fJ} & \underline{0} & & \\ & & & & & & & & & & \\ & e_2' & d_1 & c_1 & b_1 & & & & & & \\ & \underline{fJ} & \underline{fJ} & \underline{fJ} & \underline{fJ} & & & & & & \\ & & & & & & P_0 & & & & \\ & & & & & & \cup & & & & \\ & & & & & & Q_0^2 & & & & \\ & & & & & & \cup & & & & \\ & & & & & & Q_{20} & & & & \end{array},$$

where  $i_1 = k_1 + k_2 + i_1'$  and  $f$  runs over all the idempotents in (0). From the above we have

$$(8) \quad \begin{array}{ccccccccccc} & k_1 & k_2 & i_1' & j_1 & i_2' & e_1' & h_1 & g_1 & & \\ P_2'/Q_2 & \underline{0} & \underline{fR/fJ} & \underline{fR} & \underline{fR/fJ} & \underline{fR} & \underline{fR} & \underline{0} & \underline{fJ} & & \\ Q^2/Q_2 & \underline{0} & \underline{fR/J} & \underline{fR} & \underline{0} & \underline{fJ} & \underline{0} & \underline{0} & \underline{fJ} & & \\ & & & & & & & & & & \\ & e_2' & d_1 & c_1 & b_1 & & & & & & \\ & \underline{fJ} & \underline{0} & \underline{fJ} & \underline{fJ} & & & & & & \\ & & & & & & P_0/Q_{20} & & & & \\ & & & & & & \cup & & & & \\ & & & & & & Q_0^2/Q_{20} & & & & \end{array}$$

Now we come back to (2).  $M' = P'/Q$  and  $P'Q = (P_1/Q_1 \oplus P'_2/Q_2)/(Q/Q_1 \oplus Q_2)$  and  $Q/(Q_1 \oplus Q_2) = Q^2/Q_2(\theta)$  for some  $\theta \in \text{Hom}_R(Q^2/Q_2, Q^1/Q_1)$ . Since  $P_0$  is semisimple,  $Q_0^2/Q_{20}$  is a direct summand of  $P_0/Q_{20}$ . Now  $P'_2/Q_2 = \sum_f ((fR/fJ)^{(k_{2,f})} \oplus \dots \oplus (fR)^{(i_{2',f})} \oplus \dots) \oplus P_0/Q_{20} \supset Q^2/Q_2 = \sum_f ((fR/fJ)^{(k_{2,f})} \oplus \dots \oplus (fJ)^{(i_{2',f})} \oplus \dots) \oplus Q_0^2/Q_{20}$ . We compare direct summands of  $P'_2/Q_2$  and  $Q^2/Q_2$ . Then we know that only one summand  $(fJ)^{(i_{2',f})}$  of  $Q^2/Q_2$  is a proper submodule of  $(fR)^{(i_{2',f})}$ , which is a direct summand of  $P'_2/Q_2$  for each  $f$ . Consider  $\theta|(fJ)^{(i_{2',f})}$ . Since  $Q \subset J(P)$ , we know from a similar argument in the proof of Lemma 2 that  $\theta|(fJ)^{(i_{2',f})}$  is induced from an  $\theta' \in \text{Hom}_R(fJ^{(i_{2',f})}, P_1)$ , and hence  $\theta|(fJ)^{(i_{2',f})}$  is extensible to  $\Theta \in \text{Hom}_R((fR)^{(i_{2',f})}, P_1/Q_1)$ . Therefore

**Lemma 4.** *Let  $M'$  be as above and assume (\*). Then*

$M' = P'/Q \approx P_1/Q_1 \oplus \sum_f \oplus ((fR)^{(e'_{f1})} \oplus (fR/fJ)^{((j'_{f1} + i'_{f2})_f)} \oplus (fJ)^{((b_1 + e'_{f2})_f)}) \oplus \tilde{S}$ ,  
where  $\tilde{S}$  is a direct sum of simple components of  $\text{Soc}(R)$ .

**Theorem.** *Let  $R$  be artinian. Then the following are equivalent :*

- 1) (#) holds whenever  $M$  is local.
- 2) (#) holds for any finitely generated  $R$ -modules.
- 3)  $R$  is a right almost hereditary ring with (\*).

Proof. 1)  $\rightarrow$  3) This is given by Lemma 1 and Theorem 0.

3)  $\rightarrow$  2). Assume that  $M$  is  $N$ -projective. Put  $M = P/Q$ , where  $P$  is a projective cover of  $M$ . For any submodule  $M'$  of  $M$  we can suppose  $M' = P'/Q$  for some  $P' \subset P$ . From Lemma 4,  $M'$  is a direct sum of the following modules :

1)  $P_1/Q_1$ , 2) projective module, 3) simple component of  $\text{Soc}(R)$  and 4)  $fR/fJ$ , where  $fR$  is given in (0).

From the proof of Theorem 1 in [6], p.813 we know that  $P_1/Q_1$  is  $N$ -projective in cases 2) and 3) from (\*). We assume 4), i.e.  $M' = fR/fJ$ . First we suppose that  $N$  is indecomposable. For the fixed  $f$  above (and hence e) we apply Lemma 2. Let  $N = T/C$ ;  $T$  is a projective cover of  $N$ . We use the same notations as in Lemma 2. If  $T = fR$ , then  $M'$  is trivially almost  $N$ -projective (cf. Theorem 0). Hence we assume  $q \neq 0$  from Lemma 2. Take any element  $\theta$  in  $\text{Hom}_R(fR, T)$ . Then  $\theta = \theta_1 + \theta_2$  where  $\theta_1 \in \text{Hom}_R(fR, (eR)^{(p)})$ , and  $\theta_2 \in \text{Hom}_R(fR, \sum \oplus g_j R)$ . Here we recall the proof of Lemma 4. First we consider the decomposition:  $e_i R \supset e_i J \approx \sum_k \oplus (f_{ik} R)^{(n_{ik})} \oplus \sum \oplus S_{ij}$  as in (0). Let

$$\mu_k^i: e_i J \rightarrow (f_{ik} R)^{(n_{ik})}$$

be the projection of  $e_i J$  onto the  $k$ th component  $(f_{ik} R)^{(n_{ik})}$ . Next we take the decomposition of  $P_2$  in (2'). Let

$$\xi_q^p: J(P_2) \rightarrow e_p J$$

be the projection of  $J(P_2)$  onto the radical  $e_p J$  of the  $q$ th component of  $(e_p R)^{(a_p)}$



in (2'). we recall the situation where the case 4) occurs. If we carefully observe it, then we know that it comes from  $P_{2jk}$  and (6), i.e.,  $f_{jk}=f$ ,  $e=e_j$  and  $e_jR \supseteq e_jJ \approx (f_{jk}R)^{(n_{jk})} \oplus \dots$ , and  $0 \neq \mu_k^j \xi_x^j(Q^2) \subset (f_{jk}R)^{(n_{jk})}$  for some  $x$  (note  $Q^2 \subset J(P_2)$ ). Since  $Q^2 \subset J(P_2)$ , there exists a simple submodule  $S$  in  $Q^2$  such that  $\mu_k^j \xi_k^j(S) \neq 0$  from Theorem 0 and [10], Corollary to Lemma 2. Further since  $S$  is simple,  $\xi_x^j(S) \subset \sum_q \oplus (f_{jq}R)^{(n_{jq})} \oplus \sum_t \oplus S_{jt}$  from (\*). To the above  $e$  and  $f$  we consider a homomorphism

$$(9) \quad \Theta: P \xrightarrow{\pi} P_2 \xrightarrow{\lambda} eR \xrightarrow{\theta} eR \subset T,$$

where  $\pi$  is the projection,  $\lambda$  is the projection onto  $eR$  such that  $\lambda|J(P_2) = \xi_k^j$ ,  $\theta$  is any homomorphism and the last  $eR$  is the any direct component of  $(eR)^{(p)}$  in  $T$ . Since  $P/Q$  is  $N=T/C$ -projective,  $\Theta(Q) \subset C$ . Further since  $\xi_k^j(S)$  is non-zero and simple and  $\mu_k^j \xi_k^j(S) \neq 0$ ,  $eRe \xi_k^j(S) \supset (f_{jk}J)^{(n_{jk})}$  by Theorem 0. Moreover  $\xi_x^j(S) = \lambda\pi(S)$ , and hence  $(f_{jk}J)^{(pn_{jk})} \subset \sum_{\theta \in (eRe)^{(p)}} \theta \xi_x^j(S) = \sum_{\theta} \theta \lambda\pi(S) = \sum_{\theta} \theta \Theta(S) \subset \sum_{\theta} \theta \Theta(Q) \subset C$ . As a consequence  $\theta_1(fJ) \subset (f_{jk}J)^{(pn_{jk})} \subset C$ , and clearly  $\theta_2(fJ) = 0$ . Accordingly  $M'$  is  $T/C$ -projective. Finally let  $N = \sum \oplus N_i$ ; the  $N_i$  are indecomposable. Then  $M'$  is almost  $N_i$ -projective as above. If  $M' = fR/fJ$  is not  $N_i$ -projective,  $N_i = fR/A$  from [3], Theorem 1. Hence  $M'$  is almost  $N$ -projective by [5], Theorem. Thus we have shown the implication.

2)  $\rightarrow$  1). This is trivial.

Here we apply Theorem to special hereditary algebras. Let  $R$  be a hereditary algebra over a field  $K$ . Assume

$$(10) \quad eRe = eK \text{ for any primitive idempotent } e.$$

**Corollary.** *Let  $R$  be a basic hereditary algebra as above. Then the following are equivalent:*

- 1) (#) holds when  $M$  and  $N$  are local.
- 2) (#) holds when  $M$  is local and  $N$  is a direct sum of local modules.
- 3) i)  $J^3=0$ , ii)  $J$  is a direct sum of uniserial modules, and iii)  $R/\text{Soc}(R)$  is left serial.

Furthermore the following are equivalent:

- 4) (#) holds for any  $R$ -modules.
- 5) i) 3) holds, ii)  $J^2$  is square-free and iii) any simple component ( $\approx fJ$ ) of  $J^2$  is never isomorphic to any simple ones which are not contained in  $J^2$ , except  $fJ$  in  $fR$ , where  $f$  is a primitive idempotent given in (0).  
In this case (H) in [6] holds.

**Proof.** 1)  $\longleftrightarrow$  2) Since  $\text{Soc}(R)$  is projective, this is clear from [10], Theorem 2.

1)  $\longleftrightarrow$  3) Since  $R$  is hereditary, iii) in Theorem 0 always holds and i), ii) in the

proposition are equivalent to i) in Theorem 0. Further iii) in the proposition is equivalent to ii), iv) in Theorem 0.

4)  $\longleftrightarrow$  5) This is clear from the assumption (10), Lemma 1 and Theorem 0.

The last statement is clear from [6], Theorem 2.

### 3. QF-2 rings

In this section we study a left QF-2 ring with (#) as right  $R$ -modules (cf. [10], Proposition 3).

**Lemma 5.** *Let  $R$  be left QF-2. Further assume that (#) holds as right  $R$ -modules when  $M$  is local and  $N$  is a direct sum of local modules. Then  $\text{Soc}(R)$  is almost projective, and hence  $R$  is right almost hereditary, (cf. Example 4 below).*

*Proof.* Let  $eR \supset eJ$  be as (0). Then for any submodule  $X$  of  $eJ$  we have  $X = \sum_i \oplus X_i \oplus X'$  by Theorem 0 and [10], Lemma 13, where  $X_i = X \cap (f_i R)^{(n_i)}$  and  $X' = X \cap (\sum \oplus S_j)$ . Further  $X_i \approx (f_i R)^{(m_i)} \oplus (f_i J)^{(m'_i)}$  by [4], Lemma 5, where  $n_i \geq m_i + m'_i$ . Let  $Y$  be a submodule of  $X_i$ . Then after changing direct decomposition of  $(f_i R)^{(m_i)} \oplus (f_i J)^{(m'_i)}$ , we can assume  $Y = \sum_i \oplus f_i R \cap Y \oplus \sum_j \oplus f_j J \cap Y$  again by [4], Lemma 5. Now we prove the lemmas. Let  $gR/gJ$  be monomorphic to  $\text{Soc}(R)$  for a primitive idempotent  $g$ . Then  $gR$  is uniserial by [10], Lemma 9. First we shall show that  $gR$  is injective if  $gJ \neq 0$ . Let  $k$  be any primitive idempotent and take any diagram

$$\begin{array}{ccc} 0 \rightarrow & K & \rightarrow kR \\ & \downarrow \rho & \\ & gR & \end{array}$$

In order to show that  $gR$  is injective, we may assume by [8], Lemma 1<sup>#</sup> that  $\rho(K)$  is simple and  $K \subset kJ$ .

a)  $kJ^2 \neq 0$ .

Then  $kR \supset kJ$  have the structure (0). Then from the initial observation and ([4], Lemma 5),  $K/\rho^{-1}(0)$  is isomorphic to one of  $S_j$ ,  $f_i J$  and  $f_i R/f_i J$  for some  $i$  and  $j$  in (0). However the last case does not occur by assumption. Hence  $gR \subsetneq kR$  or  $gR \approx kR$  by [10], Corollary to Lemma 13, provided  $\rho \neq 0$ . In the former case  $gR \approx f_j R$  in (0) for some  $j$ . On the other hand  $\bar{f}_i R \approx gR/gJ$  is not isomorphic to any simple component of  $\text{Soc}(R)$ , a contradiction. Therefore  $\rho = 0$  in this case. Assume  $gR \approx kR$ . Then  $kR$  is uniserial, and hence  $\rho$  is a monomorphism by assumption and  $K$  is simple. Accordingly  $\rho$  is extensible to an element in  $\text{Hom}_R(kR, gR)$  by [10], Lemma 13.

b)  $kJ^2 = 0$ .

Then  $kR \subseteq gR$  or  $kR \approx gR$  by [10], Corollary to Lemma 13, provided  $\rho \neq 0$ , and hence  $kR$  is uniserial. Then  $\rho$  is extensible to an element in  $\text{Hom}_R(kR, gR)$  by [10], Lemma 13.

Thus we have shown that  $gR$  is injective. Finally we shall show that  $gR/gJ$  is injective if  $gJ^2 \neq 0$ . In the above diagram we replace  $gR$  with  $gR/gJ^2$ .

a')  $kJ^2 \neq 0$ .

Then since  $K/\rho^{-1}(0) \approx \{S_j, f_k J, f_k R/f_k J\}$  as the initial observation and  $gJ$  is projective,  $f_i R/f_i J \approx gJ/gJ^2$  for some  $i$  by Theorem 0. Hence  $gR \approx kR$  by Theorem 0. As a consequence we may assume  $gR = kR$ . Since  $gJ$  is projective,  $\rho$  is given by an element  $\theta'$  in  $\text{Hom}_R(gJ, gJ)$  (which induces  $\text{Hom}_{RR}(gJ^2, gJ^2)$ ). Then  $\theta' \in \text{Hom}_R(gJ^2, gJ^2)$  is extensible to  $\theta$  in  $\text{Hom}_R(gR, gR)$  by [10], Lemma 13. Now consider  $(\theta - \theta')|gJ$ . Since  $(\theta - \theta')(gJ^2) = 0$ ,  $(\theta - \theta')|gJ = 0$  by Theorem 0. Hence  $\rho$  is extensible to  $\nu\theta : gR \rightarrow gR/gJ^2$ , where  $\nu : gR \rightarrow gR/gJ^2$  is the natural epimorphism.

b')  $kJ^2 = 0$ .

Then  $\rho = 0$  by assumption. Therefore  $gR/gJ$  is almost projective by [9], Theorem 1.

Thus  $J$  is almost projective from  $(*)$ , and hence  $R$  is right almost hereditary.

**Proposition 2.** *Let  $R$  be a left QF-2 ring. Then the following are equivalent :*

- 1)  *$R$  is a right almost hereditary ring such that  $J^3 = 0$  and if  $eJ^2 \neq 0$  for a primitive idempotent  $e$ , then  $eJ$  has the decomposition (0).*
- 2)  *$R$  is right almost hereditary and  $(\#)$  holds when  $M$  and  $N$  are local.*
- 3)  *$(\#)$  holds when  $M$  is local and  $N$  is a direct sum of local modules.*
- 4)  *$(\#)$  holds for any  $R$ -modules  $M$  and  $N$ , (cf. Example 4 below).*

Proof. 1)  $\longleftrightarrow$  2). This is given in [10], Proposition 3.

2)  $\rightarrow$  3). This is clear from Proposition 1.

3)  $\rightarrow$  4). Since  $R$  is right almost hereditary by Lemma 5, we obtain 4) by Theorem and [10], Lemma 13.

4)  $\rightarrow$  2). This is clear from Theorem.

We shall add one more property when  $R$  is left QF-2. Let  $eR \supset eJ \approx (fR)^{(n)} \oplus \cdots$  as in (0), and put  $eJ \supset \sum_{i \leq n} u_i fR \approx fR^{(n)}$ , where  $u_i fR \approx fR$ . We identify  $(fR)^{(n)}$  with  $\sum \oplus u_i fR$ .

**Lemma 6.** *Assume 1) and 2) in [10], Lemma 13 and (0). Let  $N_1$  and  $N_2$  be submodules in  $(fR)^{(n)}$ , which are isomorphic to  $fR$  and hence  $\theta : N_1 \rightarrow N_2$  be an isomorphism. Then  $\theta$  is given by an element  $z$  in  $eRe$ .*

Proof. Let  $eR \supset fR \supset S = \text{Soc}(fR)$ . Then from 1), 2) and [10], Lemma 6 we obtain

a) every automorphism of  $S$  is extensible uniquely to an automorphism of  $fR$  (cf. a') in the proof of Lemma 5).

Put  $S_i = \text{Soc}(N_i)$  for  $i=1, 2$ .

b) Assume  $N_1 = u_1 fR$  and  $S_1 = S_2$ . Let  $S_1 = xR$  and  $N_2 = yR$ ;  $y = eyf \in (fR)^{(n)}$ . Then  $y = u_1 w_1 + \dots + u_n w_n$ ; the  $w_i$  are units or zero in  $fRf$  by 2). Then  $x = yr = u_1 w_1 + \dots + u_n w_n r$  for some  $r \in R$ . On the other hand  $x = u_1 r'$  for some  $r' \in fR$ . Hence  $w_2 = \dots = w_n = 0$  (cf. the proof of [10], Lemma 13), and  $N_1 = N_2$ .

c) Assume  $N_1 = u_1 fR$  and  $\theta' : N_2 \rightarrow N_1$ . Then  $\theta'|_{S_2}$  is extensible to  $z_l \in \text{Hom}_R(eR, eR)$ , the left-sided multiplication of  $z$ , by 1).

Further  $zN_2 = N_1$  from b), and  $z$  is a unit by [10], Lemma 6. Consider  $z_l \theta'^{-1}|_{S_1} = 1_{S_1}(z_l \theta'^{-1} : N_1 \rightarrow N_1)$ . Then from a)  $z_l \theta'^{-1} = 1_{N_1}$ , and  $z_l|_{N_2} = \theta'$ .

Since  $u_1 fR \approx N_1 \approx N_2$ , we obtain a unit  $z_i$  in  $eRe$  such that  $z_{1l} : N_1 \rightarrow u_1 fR$  and  $z_{2l} : N_2 \rightarrow u_1 fR$  from c). Hence again by c)  $z_{2l} \theta z_{1l}^{-1} = z_l$  for some  $z$ , and  $\theta = (z_2^{-1} z z_1)_l$ .

Concerning Proposition 2, we have

**Proposition 3.** *Let  $R$  be artinian. Assume that  $J^3 = 0$  and  $(eR \supset) eJ$  has the demomposition (0). Then the following are equivalent :*

1) i) *Let  $S_i$  be a simple submodule of  $h_i R$  for  $i=1, 2$ . If  $\theta : S_1 \rightarrow S_2$  is isomorphic, then  $\theta$  is extensible to an element in  $\text{Hom}_R(h_1 R, h_2 R)$  or in  $\text{Hom}_R(h_2 R, h_1 R)$ , where  $h_1, h_2$  are primitive idempotents, ii)  $fR/fJ$  is never monomorphic to  $\text{Soc}(R)$ , where  $f$  appears in (0).*

2)  $R$  is left QF-2. (cf. Example 3.)

Proof. 2)  $\rightarrow$  1). This is clear from [10], Lemma 13.

1)  $\rightarrow$  2). Let  $S_1$  and  $S_2$  be simple left  $R$ -modules of  $Rh(Jh \neq 0)$  for a primitive idempotent  $h$ . Suppose  $S_i \approx R\bar{k}_i$  for  $i=1, 2$ , where the  $k_i$  are primitive idempotents. Put  $S_i = Rx_i$  with  $k_i x_i h = x_i \in J$ . Then  $k_i R \supset x_i R$  and there exists a homomorphism  $\phi_i : hR \rightarrow x_i R$ .

a)  $k_i J^2 = 0$  for  $i=1, 2$ . Then  $x_1 R \approx \bar{h} R$  since  $x_i \in k_i J$  and the  $x_i R$  are local. Hence there exists  $z$  in  $k_2 R k_1$  (or in  $k_1 R k_2$ ) such that  $zx_1 = x_2$  (or  $zx_2 = x_1$ ) by assumption. As a consequence  $S_2 = S_1$ .

b)  $k_1 J^2 \neq 0$  and  $k_2 J^2 = 0$ . Then  $x_2 R \approx \bar{h} R$  as above. If  $x_1 R \approx \bar{h} R$ , then  $S_1 = S_2$  as in a). Suppose that  $x_1 R$  is not simple. Since  $x_1 R \subset k_1 J$  and  $x_1 R$  is local,  $x_1 R$  is projective by (0), and hence  $x_1 R \approx hR$ , which is a contradiction to iii) in Theorem 0 for  $x_2 R \approx \bar{h} R$ .

c)  $k_i J^2 \neq 0$ . Since  $x_i R$  is local,  $x_i R$  is simple or projective by iii) in Theorem 0. Hence again from iii) in Theorem 0 we obtain two cases  $\alpha) x_1 R \approx x_2 R \bar{h} R$  and  $\beta) x_1 R \approx x_2 R \approx hR$  (and  $k_1 R \approx k_2 R$ ). Then from Lemma 6 and the argument in a) we obtain  $S_1 = S_2$  in both cases. Hence  $R$  is left QF-2.

We note the following fact :

*the class of rings with (#) for local modules  $M$  and  $N \cong$  the class of rings with (#) for local module  $M$  and any direct sum of local modules  $N \cong$  the class of rings with (#) for any finitely generated  $R$ -modules. See the following examples.*

#### 4. Examples

Let  $L \supset K$  be fields.

1.

$$R_1 = \begin{pmatrix} K & K & K & K \\ 0 & K & K & 0 \\ 0 & 0 & K & P \\ 0 & 0 & 0 & R \end{pmatrix}, \text{ where } P = L, K \text{ or } 0 \text{ and } e_{13}e_{34}P = 0 = e_{23}e_{34}P.$$

If  $P = L$ , then  $R_1$  satisfies the conditions in Theorem 0, but the conditions in [10], Theorem 2. If  $P = K$ ,  $R_1$  satisfies the conditions in [10], Theorem 2, but  $R_1$  is not almost hereditary. If  $P = 0$ , then  $R_1$  satisfies the conditions in Theorem.

2.

$$R_2 = \begin{pmatrix} K & L & L \\ 0 & L & L \\ 0 & 0 & L \end{pmatrix}.$$

$R_2$  satisfies the condition in Theorem, but not left QF-2.

3.  $R_3 = eK \oplus fK \oplus aK \oplus bK \oplus cK \oplus caK$ , where  $\{e, f\}$  is the set of mutually orthogonal primitive idempotents with  $1 = e + f$ ,  $a = eaf$ ,  $b = ebf$ ,  $c = fce$ , and  $ca = cb$ . Then  $R (= R_3)$  is a left QF-2 ring with  $J^3 = 0$ , but 1) in Proposition 3 does not hold as right  $R$ -modules. However  $R$  satisfies 1) in Proposition 3 as left  $R$ -modules, but not right QF-2.

4. As above  $R_4 = eK \oplus fK \oplus gK \oplus aK \oplus abK \oplus bK \oplus cK$ , where  $a = eaf$ ,  $b = fbe$  and  $c = ecg$ . Then  $R (= R_4)$  is left serial and (#) holds for local modules  $M$  and  $N$ , however  $R$  is not right almost hereditary.

---

#### References

- [1] T. Albu and C. Nastasescu : Relative finiteness in module theory, Monographs Textbooks Pure and Appl. **84**, Marcel Dekker Inc. New York and Basel.
- [2] K.R. Fuller : *Relative projectivity and injectivity classes by simple modules*, J.London Math. Soc. **5** (1972), 423-431.
- [3] M. Harada and T. Mabuchi : *On almost relative projectives*, Osaka J. Math. **26** (1989), 837-848.
- [4] M. Harada and A. Tozaki : *Almost  $M$ -projectives and right Nakayama rings*, J. Algebra **122**

- (1989), 447-474.
- [5] M. Harada : *On almost relative projectives over perfect rings*, Osaka J. Math. **27** (1990), 655-665.
  - [6] ——— : *Hereditary rings and relative projectives*, Osaka J. Math. **28** (1991), 811-827.
  - [7] ——— : *Almost hereditary rings*, Osaka J. Math. **28** (1991), 793-809.
  - [8] ——— : *Note on almost relative projectives and almost relative injectives*, Osaka J. Math. **29** (1992), 435-446.
  - [9] ——— : *Almost projective modules*, J. Algebra **157** (1993), 150-157.
  - [10] ——— : *Artinian rings related with almost relative projectivity I*, to appear.

Department of Mathematics  
Osaka City University  
Sumiyoshi-ku, Osaka 558  
Japan

