

Title	Doubly transitive groups of even degree whose one-point stabilizer has a normal subgroup isomorphic to $PSL(3, 2^n)$
Author(s)	Hiramine, Yutaka
Citation	Osaka Journal of Mathematics. 1979, 16(3), p. 817-830
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9948">https://doi.org/10.18910/9948</a>
rights	
Note	

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## DOUBLY TRANSITIVE GROUPS OF EVEN DEGREE WHOSE ONE POINT STABILIZER HAS A NORMAL SUBGROUP ISOMORPHIC TO $PSL(3,2^n)$

YUTAKA HIRAMINE

(Received October 16, 1978)

### 1. Introduction

Let  $G$  be a doubly transitive permutation group on a finite set  $\Omega$  and  $\alpha \in \Omega$ . By [4], the product of all minimal normal subgroups of  $G_\alpha$  is the direct product  $A \times N$ , where  $A$  is an abelian group and  $N$  is 1 or a nonabelian simple group.

In this paper we consider the case  $N \simeq PSL(3, q)$  with  $q$  even and prove the following:

**Theorem.** *Let  $G$  be a doubly transitive permutation group on  $\Omega$  of even degree and let  $\alpha, \beta \in \Omega$  ( $\alpha \neq \beta$ ). If  $G_\alpha$  has a normal subgroup  $N^\alpha$  isomorphic to  $PSL(3, q)$ ,  $q=2^n$ , then  $N^\alpha$  is transitive on  $\Omega - \{\alpha\}$  and one of the following holds:*

(i)  *$G$  has a regular normal subgroup  $E$  of order  $q^3=2^{3n}$ , where  $n$  is odd and  $G_\alpha$  is isomorphic to a subgroup of  $\Gamma L(3, q)$ . Moreover there exists an element  $g$  in  $Sym(\Omega)$  such that  $\alpha^g = \alpha$ ,  $(G_\alpha)^g$  normalizes  $E$  and  $A\Gamma L(3, q) \geq (G_\alpha)^g E \geq ASL(3, q)$  in their natural doubly transitive permutation representation.*

(ii)  $|\Omega| = 22$ ,  $G^\Omega = M_{22}$  and  $N^\alpha \simeq PSL(3, 4)$ .

(iii)  $|\Omega| = 22$ ,  $G^\Omega = Aut(M_{22})$  and  $N^\alpha \simeq PSL(3, 4)$ .

We introduce some notations.

- $V(n, q)$  : a vector space of dimension  $n$  over  $GF(q)$
- $\Gamma L(n, q)$  : the group of all semilinear automorphism of  $V(n, q)$
- $A\Gamma L(n, q)$  : the semidirect product of  $V(n, q)$  by  $\Gamma L(n, q)$  in its natural action
- $ASL(n, q)$  : the semidirect product of  $V(n, q)$  by  $SL(n, q)$  in its natural action
- $F(X)$  : the set of fixed points of a nonempty subset  $X$  of  $G$
- $X(\Delta)$  : the global stabilizer of a subset  $\Delta$  ( $\subseteq \Omega$ ) in  $X$
- $X_\Delta$  : the pointwise stabilizer of  $\Delta$  in  $X$
- $X^\Delta$  : the restriction of  $X$  on  $\Delta$
- $Sym(\Delta)$  : the symmetric group on  $\Delta$

- $X^H$  : the set of  $H$ -conjugates of  $X$
- $|X|_p$  : the maximal power of a prime  $p$  dividing the order of  $X$
- $I(X)$  : the set of involutions contained in  $X$
- $E_m$  : an elementary abelian group of order  $m$

Other notations are standard and taken from [1].

### 2. Preliminaries

**Lemma 2.1** *Let  $G$  be a doubly transitive permutation group on  $\Omega$  of even degree,  $\alpha \in \Omega$  and  $N^\alpha$  a normal subgroup of  $G_\alpha$  isomorphic to  $PSL(2, q)$ ,  $Sz(q)$  or  $PSU(3, q)$  with  $q (> 2)$  even. Then  $N^\alpha \simeq PSL(2, q)$ ,  $N^\alpha \neq Sz(q)$ ,  $PSU(3, q)$ ,  $N^\alpha$  is transitive on  $\Omega - \{\alpha\}$  and one of the following holds:*

(i)  $G$  has a regular normal subgroup  $E$  of order  $q^2$ ,  $N_\beta^\alpha = N^\alpha \cap N^\beta \simeq E_q$  and  $G_\alpha$  is isomorphic to a subgroup of  $\Gamma L(2, q)$ . Moreover there exists an element  $g$  in  $Sym(\Omega)$  such that  $\alpha^g = \alpha$ ,  $(G_\alpha)^g$  normalizes  $E$  and  $A\Gamma L(2, q) \geq (G_\alpha)^g E \geq ASL(2, q)$  in their natural doubly transitive permutation representation.

(ii)  $|\Omega| = 6$  and  $G^\Omega = A_6$  or  $S_6$ .

*Proof.* By Theorem 2 of [2], it suffices to consider the case that  $N_\beta^\alpha = N^\alpha \cap N^\beta \simeq E_q$  and  $G$  has a regular normal subgroup of order  $q^2$ . Since  $|N^\alpha| = q^2 - 1$ ,  $N^\alpha$  is transitive on  $\Omega - \{\alpha\}$ .

Let  $E$  be the regular normal subgroup of  $G$ . Then we may assume  $\Omega = E$ ,  $\alpha = 0 \in E$  and the semidirect product  $GL(E)E$  is a subgroup of  $Sym(\Omega)$ . There is a subgroup  $H$  of  $GL(E)$  such that  $H \simeq \Gamma L(2, q)$  and  $HE \simeq A\Gamma L(2, q)$ . Let  $L$  be the normal subgroup of  $H$  isomorphic to  $SL(2, q)$ . Then  $L_\beta \simeq E_q$  for  $\beta \in \Omega - \{\alpha\}$ . Hence  $(N^\alpha)^{\Omega - \{\alpha\}} \simeq L^{\Omega - \{\alpha\}}$  and so there are an automorphism  $f$  from  $N^\alpha$  to  $L$  and  $g \in Sym(\Omega)$  satisfying  $\alpha^g = \alpha$  and  $(\beta^x)^g = (\beta^g)^{f(x)}$  for each  $\beta \in \Omega - \{\alpha\}$  and  $x \in N^\alpha$ . From this,  $(\beta^g)^{g^{-1}xg} = (\beta^x)^g = (\beta^g)^{f(x)}$ , so that  $g^{-1}xg = f(x)$ . Hence  $g^{-1}N^\alpha g = L$ .

Set  $S = L_\beta$ ,  $X = Sym(\Omega) \cap N(L)$ ,  $D = C_x(L)$  and  $Y = N_L(S)$ . By the properties of  $A\Gamma L(2, q)$ ,  $L$  is transitive on  $\Omega - \{\alpha\}$ ,  $|F(S)| = q$  and  $Y/S \simeq Z_{q-1}$ . Hence  $D$  is semi-regular on  $\Omega - \{\alpha\}$  and  $Y^{F(S)}$  is regular on  $F(S) - \{\alpha\}$  and so  $D \simeq D^{F(S)} \leq Y^{F(S)}$  because  $[D, N^\alpha] = 1$ . Therefore  $D \leq Z_{q-1}$ . Since  $X/DL$  is isomorphic to a subgroup of the outer automorphism group of  $SL(2, q)$ , we have  $|X| \leq |\Gamma L(2, q)|$ , while  $\Gamma L(2, q) \simeq H \leq X$ . Hence  $X = H$  and  $X$  normalizes  $E$ . Therefore, as  $(G_\alpha)^g \triangleright (N^\alpha)^g = L$ , we have  $(G_\alpha)^g \leq H$ . Thus Lemma 2.1 is proved.

**Lemma 2.2** *Let  $G$  be a doubly transitive permutation group on  $\Omega$  of even degree and  $N^\alpha$  a nonabelian simple normal subgroup of  $G_\alpha$ ,  $\alpha \in \Omega$ . If  $C_G(N^\alpha) \neq 1$ , then  $N_\beta^\alpha = N^\alpha \cap N^\beta$  for  $\alpha \neq \beta \in \Omega$  and  $C_G(N^\alpha)$  is semi-regular on  $\Omega - \{\alpha\}$ . Moreover  $C_G(N^\alpha) = 0(N^\alpha)$ .*

*Proof.* See Lemma 2.1 of [2].

**Lemma 2.3** *Let  $G$  be a transitive permutation group on a finite set  $\Omega$ ,  $H$  a stabilizer of a point of  $\Omega$  and  $M$  a nonempty subset of  $G$ . Then*

$$|F(M)| = |N_G(M)| \times |\{M^g \mid M^g \subseteq H, g \in G\}| / |H|.$$

Proof. See Lemma 2.2 of [2].

**Lemma 2.4** *Let  $H$  be a transitive permutation group on a finite set  $\Delta$  and  $N$  a normal subgroup of  $H$ . Assume that a subgroup  $X$  of  $N$  satisfies  $X^H = X^N$ . Then*

- (i)  $|F(X) \cap \beta^N| = |F(X) \cap \gamma^N|$  for  $\beta, \gamma \in \Delta$ .
- (ii)  $|F(X)| = |F(X) \cap \beta^N| \times r$ , where  $r$  is the number of  $N$ -orbits on  $\Delta$ .

Proof. By the same argument as in the proof of Lemma 2.4 of [3], we obtain Lemma 2.4.

**2.5 Properties of  $PSL(3, q)$ ,  $q=2^n$ .**

Let  $N_1 = SL(3, q)$ ,  $S_1 = \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \mid a, b, c \in GF(q) \right\}$ ,  $A_1 = \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid a, b \in GF(q) \right\}$ ,  $B_1 = \left\{ \begin{pmatrix} 1 & 0 & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \mid b, c \in GF(q) \right\}$  and  $Z = \left\{ \begin{pmatrix} d & 0 & 0 \\ 0 & d & 0 \\ 0 & 0 & d \end{pmatrix} \mid d \in GF(q), d^3 = 1 \right\}$ .

Then  $|Z| = (3, q-1)$  and  $\bar{N}_1 = N_1/Z$  is isomorphic to  $PSL(3, q)$ . Set  $N = \bar{N}_1$ ,  $S = \bar{S}_1$ ,  $A = \bar{A}_1$  and  $B = \bar{B}_1$ . Then the following hold.

(i)  $N$  is a nonabelian simple group of order  $q^3(q-1)^3(q+1)(q^2+q+1)/(3, q-1)$ .

(ii)  $|S| = q^3$ ,  $S' = \Phi(S) = Z(S) = \{x^2 \mid x \in S\} = \left\{ \begin{pmatrix} 1 & 0 & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid b \in GF(q) \right\} \simeq E_q$ ,

$S/S' \simeq E_{q^2}$  and  $S$  is a Sylow 2-subgroup of  $N$ .

(iii)  $S = \langle A, B \rangle$ ,  $A \cap B = Z(S)$ ,  $I(S) \subseteq A \cup B$  and each elementary abelian subgroup of  $S$  is contained in  $A$  or  $B$ . Let  $z \in I(S) - Z(S)$ . Then  $C_S(z) = A$  or  $B$ .

(iv) Set  $M_1 = A^N$ ,  $M_2 = B^N$ . Then  $M_1 \neq M_2$  and  $M_1 \cup M_2$  is the set of all subgroup of  $N$  isomorphic to  $E_{q^2}$ .

(v) Let  $z$  be an involution of  $N$ . Then  $I(N) = z^N$  and  $|C_N(z)| = (q-1)q^3/(3, q-1)$ .

(vi) Let  $E$  denote  $A$  or  $B$ . Then  $|N_N(E)| = (q-1)^2(q+1)q^3/(3, q-1)$ ,  $N_N(E)/E \simeq Z_k \times PSL(2, q)$ , where  $k = (q-1)/(3, q-1)$  and  $N_N(E)$  is a maximal subgroup of  $N$ .

(vii) Set  $M = (N_N(E))'$ . If  $q > 2$ , then  $M = M'$ ,  $M \supseteq E$ ,  $M/E \simeq PSL(2, q)$  and  $M$  acts irreducibly on  $E$ .

(viii) Set  $\Delta = E^N$ . Then  $|\Delta| = q^2 + q + 1$  and by conjugation  $N$  is doubly transitive on  $\Delta$ , which is an usual doubly transitive permutation representation

of  $N$ . If  $C \in \{A, B\} - \{E\}$ ,  $|F(C)| = q + 1$ ,  $C$  is a Sylow 2-subgroup of  $N_{F(C)}$  and  $C$  is semi-regular on  $\Delta - F(C)$ .

**Lemma 2.6** ([6]). *Let notations be as in (2.5) and set  $G = \text{Aut}(N)$ . Then the following hold.*

(i) *There exist in  $G$  a diagonal automorphism  $d$ , a field automorphism  $f$  and a graph automorphism  $g$  and satisfy the following:*

$$G = \langle g, f, d \rangle N \supseteq H_1 = \langle f, d \rangle N \supseteq H_2 = \langle d \rangle N, H_1 = \text{P}\Gamma\text{L}(3, q), H_2 = \text{P}\text{G}\text{L}(3, q)$$

$$H_2/N \simeq Z_r, \text{ where } r = (3, q - 1), G/H_1 \simeq Z_2, H_1/H_2 \simeq Z_n \text{ and } G/H_2 \simeq Z_2 \times Z_n.$$

(ii)  $M_1 = A^{H_1}, M_2 = B^{H_1}$  and  $A^g = B$ .

**Lemma 2.7** *Let  $N = \text{PSL}(3, q)$ , where  $q = 2^n$ . Let  $R$  be a cyclic subgroup of  $N$  of order  $q + 1$  and  $Q$  a nontrivial subgroup of  $R$ . Then  $N_N(Q) = N_N(R) \simeq Z_k \times D_{2(q+1)}$ , where  $k = (q - 1)/(3, q - 1)$  and  $D_{2(q+1)}$  is a dihedral group of order  $2(q + 1)$ .*

Proof. We consider the group  $N$  as a doubly transitive permutation group on  $\Delta = \text{PG}(2, q)$  with  $q^2 + q + 1$  points. By (2.5) (i),  $R$  is a cyclic Hall subgroup of  $N$  and so we may assume  $R \leq N_\alpha$ , where  $\alpha = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \in \text{PG}(2, q)$ . Since  $|N_{\alpha\beta}| = (q - 1)^2 q^2 / (3, q + 1)$  for  $\alpha \neq \beta \in \Delta$  and  $(q + 1, (q - 1)^2 q^2) = 1$ ,  $R$  is semiregular on  $\Delta - \{\alpha\}$ . Hence  $N_N(Q) \leq N_\alpha$ . Put  $E = O_2(N_\alpha)$ . Then  $N_\alpha = N_N(E)$  by (2.5) (viii) and  $N_N(Q)E/E \simeq Z_k \times D_{2(q+1)}$  by (2.5) (vi). Since  $N_N(Q) \cap E = C_E(Q) = 1$  by (2.5) (v). Hence  $N_N(Q) \simeq Z_k \times D_{2(q+1)}$ . As  $R$  is cyclic,  $N_N(R) \leq N_N(Q)$ . Thus  $N_N(Q) = N_N(R) \simeq Z_k \times D_{2(q+1)}$ .

**Lemma 2.8** *Let  $N = \text{PSL}(3, q)$ ,  $q = 2^n$  and let  $H (\neq N)$  be a subgroup of  $N$  of odd index. Then  $H \leq N_N(E)$  for an elementary abelian subgroup  $E$  of  $N$  of order  $q^2$ .*

Proof. Let  $S, A$  and  $B$  be as in (2.5) and let  $\Delta$  be as in Lemma 2.7. Since  $|N : H|$  is odd,  $H$  contains a Sylow 2-subgroup of  $N$  and so we may assume  $S \leq H$ .

Set  $\alpha = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \beta = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \gamma = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ . Then  $S \leq N_\alpha = N_N(A), S_\beta = B, S_\gamma =$

$$\left\{ \begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid a \in \text{GF}(q) \right\} \simeq E_q \text{ and hence } |\alpha^S| = 1, |\beta^S| = q \text{ and } |\gamma^S| = q^2.$$

If  $\alpha^H = \{\alpha\}, H \leq N_\alpha = N_N(A)$  and the lemma holds. By (2.5) (i),  $(q^2 + 1, |N|) = 1$ . Hence  $\alpha^H \neq \{\alpha\} \cup \gamma^S$ , so that we may assume either  $\alpha^H = \{\alpha\} \cup \beta^H$  or  $\alpha^H = \Delta$ .

If  $\alpha^H = \{\alpha\} \cup \beta^H, \alpha^H = F(B)$  and  $B$  is a unique Sylow 2-subgroup of  $H_{F(B)}$  by (2.5) (viii). Hence  $H \supseteq B \simeq E_{q^2}$  and the lemma holds.

If  $\alpha^H = \Delta$ , by (2.5) (iv),  $N_H(A)^{F(A)}$  is transitive and so  $|H|$  is divisible by  $q+1$ . Since  $(q^2+q+1, q+1)=1$ ,  $|H_\alpha|$  is divisible by  $q+1$ . By (2.5) (vi) and by the structure of  $PSL(2, q)$ ,  $Z_m \times PSL(2, q) \simeq H_\alpha/A \leq N_N(A)/A$ , where  $m$  is a divisor of  $(n-1)/(3, n-1)$ . Therefore  $|N:H| \leq q-1$ . We now consider the action of  $N$  on the coset  $\Gamma=N/H$ . As  $|\Gamma| \neq 1$  and  $N$  is a simple group,  $N^\Gamma$  is faithful. But  $N$  has a cyclic subgroup of order  $q+1$  and so  $|\Gamma| > q+1$ , which implies  $|N:H| > q+1$ , a contradiction.

**Lemma 2.9** *Let  $N=PSL(3, q)$ , where  $q=2^{2m}$  and  $t$  a field automorphism of  $N$  of order 2. Let  $S$  be a  $t$ -invariant Sylow 2-subgroup of  $N$ . Then the following hold.*

- (i)  $Z(\langle t \rangle S) \simeq E_{\sqrt{q}}$ .
- (ii) *If  $S_1$  is a subgroup of  $\langle t \rangle S$  isomorphic to  $S$ , then  $S_1=S$ .*

Proof. Since  $C_s(t)$  is isomorphic to a Sylow 2-subgroup of  $PSL(3, \sqrt{q})$ ,  $Z(C_s(t)) \simeq E_{\sqrt{q}}$  and  $Z(C_s(t)) \leq Z(S)$  by (2.5) (ii). Hence  $Z(\langle t \rangle S) = Z(\langle t \rangle S) \cap \langle t \rangle C_s(t) \cap C(Z(S)) = Z(\langle t \rangle S) \cap C_s(t) = Z(C_s(t)) \simeq E_{\sqrt{q}}$ . Thus we have (i).

Suppose  $S_1 \neq S$ . Then  $\langle t \rangle S = S_1 S \supseteq S_1$  and  $[\langle t \rangle S : S] = [S_1 : S_1 \cap S] = 2$ . If  $Z(S_1) \not\leq S$ , we have  $S_1 = \langle z \rangle \times (S_1 \cap S)$  for an involution  $z$  in  $Z(S_1) - S$ . By (2.5) (ii),  $z \in \Phi(S_1)$  and so  $S_1 = \langle z, S_1 \cap S \rangle = S_1 \cap S$ , a contradiction. Hence  $Z(S_1) \leq S$ .

If  $Z(S_1) = Z(S)$ ,  $E_q \simeq Z(S) \leq Z(S_1 S) = Z(\langle t \rangle S) \simeq E_{\sqrt{q}}$  by (i), which is a contradiction. Hence  $Z(S_1) \neq Z(S)$ .

Let  $z$  be an involution in  $Z(S_1) - Z(S)$ . Then  $C_s(z) \simeq E_{q^2}$  by (2.5) (iii). On the other hand,  $S_1 \leq C_{\langle t \rangle S}(z)$  and  $[C_{\langle t \rangle S}(z) : C_s(z)] = 1$  or 2. From this  $S_1$  has an elementary abelian subgroup of index 2. Hence  $q=2$ , a contradiction. Thus we have (ii).

### 3. Proof of the theorem

Throughout the rest of the paper,  $G^\Omega$  always denote a doubly transitive permutation group satisfying the hypotheses of the theorem.

Since  $G_\alpha \supseteq N^\alpha$ ,  $|\beta^{N^\alpha}| = |\gamma^{N^\alpha}|$  for  $\beta, \gamma \in \Omega - \{\alpha\}$  and so  $|\Omega| = 1+r|\beta^{N^\alpha}|$ , where  $r$  is the number of  $N^\alpha$ -orbits on  $\Omega - \{\alpha\}$ . Hence  $r$  is odd and  $N_\beta^\alpha$  is a proper subgroup of  $N^\alpha$  of odd index for  $\alpha \neq \beta \in \Omega$ . Therefore, by Lemma 2.8  $N_\beta^\alpha \supseteq A$  for some elementary abelian subgroup  $A$  of order  $q^2$ . Let  $S$  be a Sylow 2-subgroup of  $N_\beta^\alpha$ . Then, by (2.5) (iii) there exists a unique elementary abelian 2-subgroup  $B$  of  $S$  such that  $A \simeq B \simeq E_{q^2}$  and  $A \neq B$ . Set  $M_1 = A^{N^\alpha}$ ,  $M_2 = B^{N^\alpha}$  and  $K = G_\alpha(M_1) = G_\alpha(M_2)$ . By (2.5) (iv),  $M_1 \cup M_2$  is the set of all elementary abelian 2-subgroup of  $N^\alpha$  of order  $q^2$  and  $G_\alpha$  acts on  $\{M_1, M_2\}$ , so that  $G_\alpha/K \leq Z_2$ . Hence  $K$  is transitive on  $\Omega - \{\alpha\}$ .

$$(3.1) \text{ Let } E=A \text{ or } B. \text{ Then } N_{G_\alpha}(E) \text{ is transitive on } F(E) - \{\alpha\}.$$

Proof. If  $E^h \leq K_\beta$  for some  $h \in K$ ,  $E^h \leq N^\alpha \cap K_\beta = N_\beta^\alpha$ . Since  $E^{N^\alpha} = E^K$  and  $A^K \neq B^K$ ,  $E^h$  is conjugate to  $E$  in  $N_\beta^\alpha$ . By a Witt's theorem  $N_K(E)$  is transitive on  $F(E) - \{\alpha\}$ . Thus  $N_{G_\omega}(E)$  is transitive on  $F(E) - \{\alpha\}$ .

(3.2) *If  $q=2$ ,  $G^\Omega$  is of type (i) of the theorem.*

Proof. Assume  $q=2$ . We note that  $PSL(3,2)$  is isomorphic to  $PSL(2,7)$ . It follows from [3] that  $G$  has a regular normal subgroup  $R$ .

Since  $K$  is transitive on  $\Omega - \{\alpha\}$ , by Lemmas 2.3 and 2.4

$$|F(A)| = 1 + \frac{|N^\alpha \cap N(A)|}{|N_\beta^\alpha|} r = \frac{24r}{|N_\beta^\alpha|} + 1 \quad \text{and}$$

$$|F(B)| = 1 + \frac{|N^\alpha \cap N(B)| |N_\beta^\alpha : N_\beta^\alpha \cap N(B)|}{|N_\beta^\alpha|} r = \frac{24r}{|N_\beta^\alpha \cap N(B)|} + 1.$$

Let  $E=A$  or  $B$ . As  $N_R(E) \neq 1$ ,  $N_G(E)^{F(E)}$  is doubly transitive by (3.1). Hence  $E \leq N^\beta$  and  $|F(A)| = 2^a$ ,  $|F(B)| = 2^b$  for some integers  $a, b$ . From this  $S = \langle A, B \rangle \leq N^\alpha \cap N^\beta$  and  $|N_\beta^\alpha : N^\alpha \cap N^\beta|$  is odd. Hence, if  $S^g \leq G_{\alpha\beta}$ ,  $S^g \leq N_\beta^\gamma \cap N_\beta^\gamma$ , where  $\gamma = \alpha^g$  and so  $S^g \leq N^\alpha \cap N^\beta$ . Since  $S$  and  $S^g$  are Sylow 2-subgroups of  $N^\alpha \cap N^\beta$ ,  $S^g$  is conjugate to  $S$  in  $N^\alpha \cap N^\beta$ . By a Witt's theorem  $N_G(S)^{F(S)}$  is a doubly transitive permutation group with a regular normal subgroup  $N_R(S)$ . Hence  $|F(S)| = 2^c$  for an integer  $c$ . By Lemmas 2.3 and 2.4,

$$|F(S)| = 1 + \frac{8 \times |N_\beta^\alpha : S|}{|N_\beta^\alpha|} r = r + 1 = 2^c.$$

Let  $z$  be an involution of  $Z(S)$  and assume  $z^g \in G_\omega$  for some  $g \in G$ . Then  $z^g \in N_\omega^\gamma$ , where  $\gamma = \alpha^g$ . Since  $|N_\omega^\gamma : N^\gamma \cap N^\omega|$  is odd,  $z^g$  is contained in  $N^\omega$ . By (2.5) (v),  $z^g$  is conjugate to  $z$  in  $N^\omega$ . Hence  $C_G(z)$  is transitive on  $F(z)$  and by Lemmas 2.3 and 2.4,

$$|F(z)| = 1 + \frac{8 \times |I(N_\beta^\alpha)|}{|N_\beta^\alpha|} r.$$

Suppose  $N_\beta^\alpha = S$ . Then  $|F(A)| = 3r + 1 = 2^a = 2^c + 2r$  and  $|F(z)| = 5r + 1$ . Hence  $r = 1$ . Since  $N_R(A) = C_R(A) \leq C_G(z)$  and  $N_R(A) \simeq E_4$ ,  $|F(z)|$  is divisible by 4. But  $|F(z)| = 5r + 1 = 6$ . This is a contradiction.

Suppose  $N_\beta^\alpha \neq S$ . Then  $N_\beta^\alpha = N_{N^\alpha}(A)$  as  $N_{N^\alpha}(A) \simeq S_4$ . From this,  $|F(B)| = 2^b = 2^c + 2r$  and so  $r = 1$ . Hence  $|\Omega| = 1 + |N^\alpha : N_\beta^\alpha| = 8$ . Thus  $|R| = 8$  and  $G_\omega \simeq GL(3,2)$ , hence  $G \simeq AL(2,3)$ .

By (3.2), it suffices to consider the case  $q > 2$  to prove the theorem. From now on we assume the following.

Hypothesis (\*):  $q = 2^n \geq 4$

(3.3) *The following hold.*

- (i)  $|N_\beta^\alpha/N^\alpha \cap N^\beta|$  is odd.
- (ii) Let  $\gamma \in \Omega$  and  $S_0$  a 2-subgroup of  $N^\gamma$ . Then  $F(S_0) = \{\delta \in \Omega \mid S_0 \leq N^\delta\}$ .

Proof. Suppose false and let  $T$  be a Sylow 2-subgroup of  $N_\alpha^\beta N_\beta^\alpha$  such that  $T \geq S$ . Then  $T \neq S$ . Set  $S_1 = T \cap N_\beta^\alpha$  and  $S_2 = T \cap N^\alpha \cap N^\beta$ . Then  $S_1$  is a Sylow 2-subgroup of  $N_\beta^\alpha$ ,  $S_1 \neq S$  and  $S_1, S_2$  and  $S$  are normal subgroups of  $T$ . By Lemma 2.2,  $S_1 N^\alpha / N^\alpha$  is isomorphic to a subgroup of the outer automorphism group of  $N^\alpha$ . It follows from Lemma 2.6 that  $S_1 N^\alpha / N^\alpha$  is abelian of 2-rank at most 2. Since  $S_1 N^\alpha / N^\alpha \simeq S_1 / S_2$  and  $S_1 \simeq S$ , we have  $S_1 / S_2 \leq E_4$  by (2.5) (ii).

Let  $A_1, B_1$  be the subgroups of  $S_1$  such that  $A_1 \simeq B_1 \simeq E_{q^2}$  and  $A_1 \cap S_2 \leq A, B_1 \cap S_2 \leq B$ . Since  $A_1 / A_1 \cap S_2 \simeq A_1 S_2 / S_2 \leq S_1 / S_1 \leq E_4$  and by the hypothesis (\*),  $q \geq 4$ , we have  $|A_1 \cap S_2| \geq q^2/4$ . Therefore, if  $A_1 \cap S_2 \leq Z(S)$ , then  $q = 4$ ,  $A_1 \cap S_2 = Z(S)$  and  $T = A_1 S$  and so  $Z(S) \leq Z(T)$ , contrary to Lemma 2.9. Hence  $A_1 \cap S_2 \not\leq Z(S)$ . Similarly  $B_1 \cap S_2 \not\leq Z(S)$ .

Let  $x \in A_1 \cap S_2 - Z(S)$ . Then  $x \in A' \leq S$  for each  $y \in A_1$  and so  $A_1$  normalizes  $A$ . Hence  $A_1$  normalizes  $B$ . Similarly  $B_1$  normalizes  $A$  and  $B$ . From this  $T = \langle A_1, B_1 \rangle S \supseteq A, B$  and so  $S_1 N^\alpha \leq K$ . Hence  $S_1 N^\alpha / N^\alpha \simeq S_1 / S_2 \simeq Z_2$ , so that there exists a field automorphism  $t$  of order 2 such that  $T = \langle t \rangle S \triangleright S$ . Since  $S_1 \leq T$  and  $S_1 \simeq S$ , we have  $S_1 = S$  by Lemma 2.9, a contradiction. Thus (i) holds.

Let  $\delta \in F(S_0) - \{\gamma\}$ . Then  $S_0 \leq N_\delta^\gamma$ . Since  $N_\delta^\gamma \supseteq N^\gamma \cap N^\delta$  and  $|N_\delta^\gamma / N^\gamma \cap N^\delta|$  is odd by (i),  $S_0 \leq N^\gamma \cap N^\delta \leq N^\delta$ . Hence  $F(S_0) \subseteq \{\delta \in \Omega \mid S_0 \leq N^\delta\}$ . The converse implication is clear. Thus (ii) holds.

(3.4) *The following hold.*

- (i)  $N_G(B)^{F(B)}$  is doubly transitive.
- (ii) If  $F(A) \neq \{\alpha, \beta\}$ ,  $N_G(A)^{F(A)}$  is doubly transitive.

Proof. Let  $E = A$  or  $B$ . By (3.3) (i),  $S$  is a Sylow 2-subgroup of  $N_\beta^\alpha$ . Therefore, by a similar argument as in (3.1),  $N_{G_\beta}(E)$  is transitive on  $F(E) - \{\beta\}$ . Suppose  $N_G(E)^{F(E)}$  is not doubly transitive. Then,  $F(E) = \{\alpha, \beta\}$  by (3.1) and (3.3). Since  $N_{N^\alpha}(E)$  acts on  $F(E)$  and fixes  $\{\alpha\}$ , we have  $N_{N^\alpha}(E) \leq N_\beta^\alpha$ . On the other hand  $N_{N^\alpha}(E)$  is a maximal subgroup of  $N^\alpha$  by (2.5) (vi). Hence  $N_{N^\alpha}(E) = N_\beta^\alpha$ . If  $E = B$ , then  $N_\beta^\alpha \supseteq A$ , a contradiction. Thus  $E = A$  and (3.4) follows.

(3.5) *The following hold.*

- (i) Put  $M = (N_{N^\alpha}(A))'$ . Then  $F(M) = F(A)$ .
- (ii)  $N_\beta^\alpha = N_\gamma^\alpha$  for each  $\gamma \in F(A) - \{\alpha\}$ .

Proof. Suppose  $F(M) \neq F(A)$ . Then  $M \not\leq N_G(A)_{F(A)}$ . It follows from (3.4) that  $F(A) \neq \{\alpha, \beta\}$  and  $N_G(A)^{F(A)}$  is doubly transitive. Moreover by (2.5) (vii)  $N_{G_\alpha}(A)^{F(A)} \supseteq M^{F(A)} \simeq PSL(2, q)$  as  $q > 2$ . By Lemma 2.1,  $r = 1$  and either (1)  $q =$



4 and  $N_G(A)^{F(A)} = A_6$  or  $S_6$  or (2)  $|F(A)| = q^2$ .

If (1) holds,  $|F(A)| = 1 + |N_{N^\alpha}(A) : N_\beta^\alpha| = 1 + 2^6 \cdot 3 \cdot 5 / |N_\beta^\alpha| = 6$  and so  $|N_\beta^\alpha| = 2^6 \cdot 3$ . Hence  $|\Omega| = 1 + |N^\alpha : N_\beta^\alpha| = 1 + 2^6 \cdot 3^2 \cdot 5 \cdot 7 / 2^6 \cdot 3 = 2 \cdot 5 \cdot 3$ . Let  $z$  be an involution of  $N^\alpha \cap N^\beta$ . Then, by (2.5) (v) and (3.3),  $z^G \cap G_\omega = z^{G_\omega}$ , so that  $C_G(z)^{F(z)}$  is transitive by a Witt's theorem. On the other hand  $|F(z)| = 1 + \frac{|C_{N^\alpha}(z)| \times |I(N_\beta^\alpha)|}{|N_\beta^\alpha|} = 1 + 2^6 \cdot 3^3 / 2^6 \cdot 3 = 10$ . In particular  $|C_G(z)|$  is divisible by

5. Let  $R$  be a Sylow 5-subgroup of  $C_G(z)$ . Then  $|\Omega|$ ,  $|G_\omega : N^\alpha|$  and  $|N_\beta^\alpha|$  are not divisible by 5 and so  $F(R) = \{\gamma\}$  and  $R \leq N^\gamma$  for some  $\gamma \in \Omega$ . Therefore  $\langle z \rangle \times R \leq N^\gamma$  by (3.3) (ii). But  $|C_{N^\gamma}(z)| = 2^6$  by (2.5) (v). This is a contradiction.

If (2) holds,  $q^2 = |F(A)| = 1 + |N_{N^\alpha}(A) : N_\beta^\alpha|$ , hence  $|N_\beta^\alpha| = (q-1)q^3 / (3, q-1)$ . From this  $|\Omega| = 1 + |N^\alpha : N_\beta^\alpha| = 1 + (q-1)(q+1)(q^2+q+1) = q(q^3+q^2-1)$ . Hence  $|G|_2 = |\Omega|_2 \times |G_\omega|_2 = q \times |G_\omega : K| \times |K|_2$ . On the other hand  $|N_G(A)|_2 = |F(A)| \times |N_{G_\omega}(A)|_2 = q^2 |K|_2$  because  $K = N_{G_\omega}(A)N^\alpha$ . Therefore  $q^2 |K|_2 = |N_G(A)|_2 \leq |G|_2 = q \times |G_\omega : K| \times |K|_2 \leq 2q |K|_2$  and we obtain  $q=2$ , contrary to the hypothesis (\*). Thus we have (i).

Let  $\gamma \in F(A) - \{\alpha\}$ . By (i) and (3.4) (ii),  $N_\gamma^\alpha \supseteq A$  and  $M \leq N_\gamma^\alpha$ . Since  $N_{N^\alpha}(A)/M \simeq Z_k$ , where  $k = (q-1)/(3, q-1)$  and  $|N_\beta^\alpha/M| = |N_\gamma^\alpha/M|$ , we have  $N_\beta^\alpha = N_\gamma^\alpha$ . Thus (ii) holds.

$$(3.6) \quad B \notin A^G \text{ and } G_\omega = K.$$

Proof If  $B \in A^G$ , by (3.4) (i), there is an element  $g \in G_{\alpha\beta}$  such that  $B = A^g$ . Hence  $N_\beta^\alpha = g^{-1}N_\beta^\alpha g \supseteq g^{-1}Ag = B$  and so  $M$  normalizes  $\langle A, B \rangle = S$ , a contradiction.

(3.7) Set  $L = (N_{N^\alpha}(B))'$ . Then  $r=1$ ,  $L_{F(B)} = B$ ,  $L^{F(B)} = L/B \simeq PSL(2, q)$ ,  $L_\beta = S$  and one of the following holds.

- (i)  $C_G(N^\alpha) = 1$ ,  $|F(B)| = 6$ ,  $q=4$  and  $N_G(B)^{F(B)} = A_6$  or  $S_6$ .
- (ii)  $C_G(N^\alpha) \leq Z_{q-1}$ ,  $|F(B)| = q^2$  and  $N_G(B)^{F(B)}$  has a regular normal subgroup.

Proof. By (3.4) (i),  $N_G(B)^{F(B)}$  is doubly transitive. If  $L \leq G_{\alpha\beta}$ , then  $L \leq N_\beta^\alpha$  and so  $B \leq L = L' \leq (N_\beta^\alpha)' = M$ . Therefore  $L = M$  and  $M \supseteq \langle A, B \rangle = S$ , a contradiction. Hence  $L \not\leq G_{\alpha\beta}$ . From this  $N_{G_\omega}(B)^{F(B)} \supseteq L^{F(B)} \simeq PSL(2, q)$  and (3.7) follows from Lemmas 2.1 and 2.2.

(3.8) If (i) of (3.7) occurs, then we have (ii) or (iii) of the theorem.

Proof. Since  $|F(B)| = 1 + |N_{N^\alpha}(B) : N_{N_\beta^\alpha}(B)| = 6$  and  $|N_\beta^\alpha : N_{N_\beta^\alpha}(B)| = |N_\beta^\alpha : N_{N_\beta^\alpha}(S)| = 5$ , we have  $|N_\beta^\alpha| = 2^6 \cdot 3 \cdot 5$ . Hence  $N_\beta^\alpha = N_{N^\alpha}(A)$  and so  $|\Omega - \{\alpha\}| = |N^\alpha : N_\beta^\alpha| = 21$ . By (3.6),  $PSL(3, 4) \leq (G_\omega)^{\Omega - \{\alpha\}} \leq P\Gamma L(3, 4)$  in their natural doubly transitive permutation representation and hence (3.8) follows from Satz 7 of [7].

In the rest of this paper, we consider the case (ii) of (3.7). From now on

we assume the following.

Hypothesis (\*\*):  $r=1, q=2^n > 2, |F(B)|=q^2$  and  $N_G(B)^{F(B)}$  is a doubly transitive permutation group with a regular normal subgroup.

(3.9) *The following hold.*

- (i)  $N_\beta^\alpha = N^\alpha \cap N^\beta = M$  and  $|N_\beta^\alpha| = (q-1)(q+1)q^3$ .
- (ii)  $n$  is odd.
- (iii)  $|F(A)| = q$ .

Proof. Since  $q^2 = |F(B)| = 1 + |N_{N^\alpha}(B) : N_{N_\beta^\alpha}(B)|$  by (3.7), we have  $|N_{N_\beta^\alpha}(B)| = |N_{N^\alpha}(B)| / (q^2 - 1) = (q-1)q^3 / (3, q-1)$ . As  $N_\beta^\alpha \supseteq A$ ,  $N_{N_\beta^\alpha}(B) = N_{N_\beta^\alpha}(\langle A, B \rangle) = N_{N_\beta^\alpha}(S)$ . On the other hand, from (2.5) (vi)  $|N_{N_\beta^\alpha}(S)| = |N_\beta^\alpha : M| \times |N_M(S)| = |N_\beta^\alpha : M| \times (q-1)q^3$ . Therefore  $(3, q-1) = 1$  and  $|N_\beta^\alpha : M| = 1$ . Hence  $N_\beta^\alpha = M$  and  $n$  is odd. By (3.3) (i) and (2.5) (vii),  $N_\beta^\alpha = N^\alpha \cap N^\beta$ . Hence  $|F(A)| = 1 + |N_{N^\alpha}(A)| / |N^\alpha| = q$ . Thus we have (3.9).

(3.10) *Put  $m = |G_\alpha : N^\alpha|$ . Then the following hold.*

- (i)  $m$  is odd and  $S$  is a Sylow 2-subgroup of  $G_\alpha$ .
- (ii)  $|\Omega| = q^3$  and  $|G| = q^6(q-1)^2(q+1)(q^2+q+1)m$ .

Proof. Set  $C^\alpha = C_G(N^\alpha)$ . By (3.6), (3.9) (ii) and Lemma 2.6,  $|G_\alpha / C^\alpha N^\alpha|$  is odd. Since  $C^\alpha \cap N^\alpha = 1, m = |G_\alpha / C^\alpha N^\alpha| \cdot |C^\alpha|$  and so  $m$  is odd by Lemma 2.2. Therefore  $S$  is a Sylow 2-subgroup of  $G_\alpha$  and so (i) holds.

Since  $|\Omega| = 1 + |N^\alpha : N_\beta^\alpha|, |\Omega| = q^3$  by (3.9). From this  $|G| = |\Omega| \times |G_\alpha| = q^3 m |N^\alpha| = q^6(q-1)^2(q+1)(q^2+q+1)m$ . Thus (ii) holds.

(3.11) *Let  $z$  be an involution of  $G_\alpha$ . Then  $|F(z)| = q^2$ . In particular  $B$  is semi-regular on  $\Omega - F(B)$ .*

Proof. By (3.10) (ii),  $z$  is contained in  $N^\alpha$ . By (2.5) (vii) and (3.9) (ii),  $|I(N_\beta^\alpha)| = |N_\beta^\alpha : N_{N_\beta^\alpha}(S)| \times (q^2 - q) + q^2 - 1 = (q+1)(q^2 - q) + q^2 - 1 = (q-1)(q+1)^2$ , hence  $|F(z)| = 1 + q^3(q-1) \times (q-1)(q+1)^2 / q^3(q-1)(q+1) = q^2$  by Lemma 2.3. As  $|F(B)| = q^2, B$  is semi-regular on  $\Omega - F(B)$ .

(3.12) *Set  $\Delta = F(B)$ . Then the following hold.*

- (i)  $G_\Delta \supseteq B$  and  $B$  is a Sylow 2-subgroup of  $G_\Delta$ .
- (ii)  $G(\Delta) = N_G(B)$  and  $|N_G(B)| = q^5(q-1)^2(q+1)m$ .

Proof. Since  $N_{N^\alpha}(B) \leq N^\alpha(\Delta) \neq N^\alpha$  and  $N_{N^\alpha}(B)$  is a maximal subgroup of  $N^\alpha$ , we have  $N_{N^\alpha}(B) = N^\alpha(\Delta)$ . By (3.7),  $B$  is a normal Sylow 2-subgroup of  $(N^\alpha)_\Delta$  and (i) follows immediately from (3.10) (i).

Since  $G(\Delta) \supseteq G_\Delta$  and  $B$  is a characteristic subgroup of  $G_\Delta$  by (i), we have  $G(\Delta) \leq N_G(B)$ . The converse implication is clear. Thus  $G(\Delta) = N_G(B)$ . By (3.6),  $G_\alpha = N_{G_\alpha}(B)N^\alpha$  and so  $|N_{G_\alpha}(B) : N_{N^\alpha}(B)| = |G_\alpha : N^\alpha| = m$ . Hence  $|N_G(B)|$

$= |F(B)| \times |N_{G_\alpha}(B)| = q^2 m \times |N_{N^\alpha}(B)| = q^5 (q-1)^2 (q+1) m$ . Thus we have (ii).

(3.13) *Let  $T_1$  be a Sylow 2-subgroup of  $N_G(B)$  and  $T_2$  a Sylow 2-subgroup of  $N_G(T_1)$ . Then  $T_1 \neq T_2$ . Let  $x$  be an element of  $T_2 - T_1$  and set  $U = BB^x$ . Then  $U \simeq E_{q^4}$  and for each  $\gamma \in \Omega$ ,  $U_\gamma \simeq E_{q^2}$ ,  $U_\gamma \in B^G$ ,  $\gamma^U = F(U_\gamma)$  and  $|\gamma^U| = q^2$ . Moreover  $U_\gamma = U_\delta$  for all  $\delta \in \gamma^U$ .*

Proof. If  $B \cap B^x \neq 1$ , by (3.11) and (3.12) (i), we have  $B = B^x$  and so  $x \in T_1$ , contrary to the choice of  $x$ . Hence  $B \cap B^x = 1$ . As  $T_1 \supseteq B$  and  $T_1 = T_1^x \supseteq B^x$ ,  $U = B \times B^x$  and  $U \simeq E_{q^4}$ .

Let  $\gamma \in \Omega$  and put  $D = U_\gamma$ . Then  $F(D) \supseteq \gamma^U$  as  $U$  is abelian. Therefore  $|U : D| = |\gamma^U| \leq q^2$  by (3.11), while  $|D| \leq q^2$  because  $D$  is an elementary abelian subgroup of  $N^\gamma$ . Hence  $D \simeq E_{q^2}$  and  $|F(D)| = |\gamma^U| = q^2$ . By (3.6) and (3.9) (iii),  $D \in B^G$ . Since  $U_\gamma \leq U_\delta \simeq E_{q^2}$  for each  $\delta \in \gamma^U$ , we have  $U_\gamma = U_\delta$ .

(3.14) *Let  $U$  be as in (3.13). Let  $\Gamma = \{X_i \mid 1 \leq i \leq s\}$  be the set of  $U$ -orbits on  $\Omega$  and set  $B_i = U_\gamma$  for  $\gamma \in X_i$  with  $1 \leq i \leq s$ . Then the following hold.*

(i)  $s = q, \Omega = \bigcup_{i=1}^q X_i$  and  $|X_i| = q^2$ .

(ii)  $B_i$  is semi-regular on  $\Omega - X_i$  and  $B_i \cap B_j = 1$  for each  $i, j$  with  $i \neq j$ .

Proof. By (3.10) (ii) and (3.13),  $|X_i| = q^2$  and  $|\Omega| = q^3$ , hence  $s = q$ . Clearly  $\Omega = \bigcup_{i=1}^q X_i$ . Thus we have (i).

By (3.13) (ii),  $B_i$  is conjugate to  $B$  for each  $i$ . Hence  $B_i$  is semi-regular on  $\Omega - X_i$  by (3.11). Therefore, if  $B_i \cap B_j \neq 1$ , then  $X_i = F(B_i) = F(B_j) = X_j$ , so that  $i = j$ . Thus we have (ii).

(3.15) *Set  $Y = \{B_i \mid 1 \leq i \leq q\}$  and let  $D \in Y$ . Then  $N_G(D) \leq N_G(U)$  and  $U$  is a unique Sylow 2-subgroup of  $C_G(D)$ .*

Proof. Suppose  $N_G(D) \not\leq N_G(U)$ . Since  $[N_G(D), U] \not\leq U$ , there exist  $g \in N_G(D)$  and  $B_i \in Y - \{D\}$  such that  $(B_i)^g \not\leq U$ . Set  $D_1 = (B_i)^g$ . Since  $[D_1, D] = [B_i, D]^g = 1$ , it follows from (3.10) (i) that  $F(D_1) \cap F(D) = \phi$  and so  $D$  is regular on  $F(D_1)$  by (3.11). Hence  $F(D_1) = \gamma^D = \gamma^U$  for  $\gamma \in F(D_1)$ . By (3.14),  $F(D_1) = F(B_j)$  for some  $B_j \in Y$ . By (3.12) (i),  $D_1 = B_j$ , so that  $D_1 \leq U$ , a contradiction. Thus we have  $N_G(D) \leq N_G(U)$ . Hence  $U \leq O_2(C_G(D))$ . Since  $U \leq C_G(B)$ ,  $C_G(B)$  is transitive on  $F(B)$ . Hence  $|C_G(B)|_2 = |F(B)| \times |C_{G_\alpha}(B)|_2 = q^4$  by (3.10) (i). Therefore  $|C_G(D)|_2 = q^4$  as  $D \in B^G$  and so  $U$  is a unique Sylow 2-subgroup of  $C_G(D)$ .

(3.16)  $|N_G(U)| = q^6 (q-1)^2 (q+1) m$ .

Proof. Let  $S_1$  be a Sylow 2-subgroup of  $N_G(U)$  and  $S_2$  be a Sylow 2-subgroup of  $N_G(S_1)$ . Suppose  $S_1 \neq S_2$  and let  $w$  be an element of  $S_2 - S_1$ .

Set  $\gamma = \alpha^{w^{-1}}$ . Then  $(U_\gamma)^w \in B^G$  by (3.13) and  $(U_\gamma)^w \leq (G_\gamma)^w = G_\alpha$ . Since  $U$  and  $U^w$  are normal subgroups of  $S_1$ ,  $\langle B, (U_\gamma)^w \rangle$  is 2-subgroup of  $G_\alpha \cap S_1 = S$ . Hence  $B = (U_\gamma)^w$  by (2.5) (iii) and (3.6). Therefore  $U, U^w \leq C_G(B)$ , so that  $U = U^w$  by (3.15) and  $w \in S_2 \cap N_G(U) = S_1$ , contrary to the choice of  $w$ . Hence  $S_1 = S_2$  and  $S_1$  is a Sylow 2-subgroup of  $G$ . It follows from (3.10) that  $|S_1| = q^6$ .

We now consider the action of  $N_G(U)$  on  $\Gamma = \{X_i \mid 1 \leq i \leq q\}$ . Set  $\Delta = F(B)$ . By (3.12),  $S_1(\Delta) \leq G(\Delta) = N_G(B)$  and  $|N_G(B)|_2 = q^5$  and so  $|S_1 : S_1(\Delta)|$  is divisible by  $q$ . Hence  $S_1$  is transitive on  $\Gamma$  and so  $N_G(U)$  is transitive on  $\Gamma$ . Therefore  $|N_G(U)| = q \times |N_G(U) \cap N_G(B)| = q \times |N_G(B)| = q^6(q-1)^2(q+1)m$  by (3.12) (ii) and (3.15).

(3.17) *Let  $R$  be a cyclic subgroup of  $N_\beta^\alpha$  of order  $q+1$ . Then  $|F(R)| = q$  and  $R$  is semi-regular on  $\Omega - F(R)$ .*

Proof. Since  $N_\beta^\alpha/A \cong PSL(2, q)$ , there exists a cyclic subgroup  $R$  of  $N_\beta^\alpha$  of order  $q+1$ . Let  $Q \neq 1$  be a subgroup of  $R$ . Then, by Lemma 2.7  $|F(Q)| = 1 + \frac{|N_{N^\alpha}(Q)| \times |N_\beta^\alpha : N_{N_\beta^\alpha}(Q)|}{|N_\beta^\alpha|} = 1 + \frac{2(q-1)(q+1)}{2(q+1)} = q$ . Thus (3.17) holds.

(3.18) *Let  $V \in U^G$ . If  $V \neq U$ , then  $|F(U_\gamma) \cap F(V_\gamma)| = 1$  or  $q$  for  $\gamma \in \Omega$ .*

Proof. Suppose  $\gamma \neq \delta \in F(U_\gamma) \cap F(V_\gamma)$ . By (3.13),  $U_\gamma, V_\gamma \in B^G$  and so by (3.3) (ii),  $U_\gamma, V_\gamma \leq N^\gamma \cap N^\delta$ . Set  $H = O_2(N_\delta^\gamma)$ . Then, by (3.6) and (3.9) (i),  $U_\gamma H$  and  $V_\gamma H$  are Sylow 2-subgroups of  $N_\delta^\gamma$ . If  $U_\gamma H = V_\gamma H$ , then  $U_\gamma = V_\gamma$  and  $U, V \leq C_G(U_\gamma)$ . By (3.15) we have  $U = V$ , a contradiction. Therefore  $U_\gamma H \neq V_\gamma H$ . Set  $X = \langle U_\gamma, V_\gamma \rangle$ . Then  $XH = N_\delta^\gamma$  because  $N_\delta^\gamma/H \cong PSL(2, q)$ ,  $q = 2^n$ , and  $PSL(2, q)$  is generated by its two distinct Sylow 2-subgroups. Hence  $N_\delta^\gamma \geq X \cap H$ . By (2.5) (iii),  $E_q \cong U_\gamma \cap H \leq X \cap H$ . Since  $N_\delta^\gamma$  acts irreducibly on  $H$  by (2.5) (vii),  $X \cap H = H$  and hence  $H \leq X$ . From this  $X = N_\delta^\gamma$ . Thus, by (3.5)(i) and (3.9),  $|F(U_\gamma) \cap F(V_\gamma)| = |F(X)| = |F(N_\delta^\gamma)| = q$ .

(3.19) *Let  $Q$  be a cyclic subgroup of  $N_{N^\alpha}(B)$  of order  $q+1$ ,  $V \in U^G$  and set  $P = N_Q(V)$ . Then the following hold.*

- (i)  $Q$  is semi-regular on  $\Omega - F(Q)$  and  $|F(Q)| = q$ .
- (ii) If  $P \neq 1$  and  $V \geq D \in B^G$ , then  $P$  normalizes  $D$  and  $|F(P) \cap F(D)| = 1$ .

Proof. Since  $N_{N^\alpha}(B)/B \cong PSL(2, q)$ , there exists a cyclic subgroup  $Q$  of  $N_{N^\alpha}(B)$  of order  $q+1$ . Clearly  $Q$  is a cyclic Hall subgroup of  $N^\alpha$ , hence  $Q$  is conjugate to  $R$  defined in (3.17). By (3.17),  $Q$  is semi-regular on  $\Omega - F(Q)$  and  $|F(Q)| = q$ . Thus (i) holds.

Suppose  $P \neq 1$  and let  $\gamma \in F(P)$ . Then, by (3.9) (i),  $P \leq N^\gamma$  and hence  $P$  normalizes  $N^\gamma \cap V$ . By (3.10) (i) and (3.13),  $N^\gamma \cap V = V_\gamma$  and  $V_\gamma \in B^G$  and so  $P \leq N_{N^\gamma}(V_\gamma)$  and  $N_G(V_\gamma)^{F(V_\gamma)} \cong N_G(B)^{F(B)}$ . Hence we have  $F(P) \cap F(V_\gamma) = \{\gamma\}$  by (3.7). As  $|F(P)| = q$  by (i), (ii) holds.

(3.20) *Let  $V \in U^G - \{U\}$  and let  $Q$  be a cyclic subgroup of  $N_{N^a}(B)$  of order  $q+1$ . Then  $N_Q(V)=1$ .*

Proof. Set  $P=N_Q(V)$  and assume  $P \neq 1$ . Let  $\gamma \in \Omega - F(Q)$  and set  $B_1=U_\gamma$ ,  $B_2=V_\gamma$ . By (3.15),  $Q$  normalizes  $U$  and so by (3.19)  $Q$  normalizes  $B_1$ . Similarly  $P$  normalizes  $B_2$ . Therefore  $F(B_1) \cap F(B_2) \geq \gamma^P \neq \{\gamma\}$  as  $P \neq 1$  and  $P$  is semi-regular on  $\Omega - F(Q)$ . By (3.18), we have  $|F(B_1) \cap F(B_2)|=q$ . Since  $P$  acts on  $F(B_1) \cap F(B_2)$  and  $|P|$  divides  $q+1$ ,  $P$  fixes at least two points of  $F(B_1) \cap F(B_2)$ , which contradicts to (3.19).

(3.21) *Let  $T$  be a Sylow 2-subgroup of  $N_G(U)$ . Then, for each  $V \in U^G - \{U\}$ ,  $|T : N_T(V)|$  is divisible by  $q$ .*

Proof. Suppose  $|T : N_T(V)| < q$  and set  $T_1=N_T(V)$ . Then  $|T_1| > q^5$  as  $|T|=q^6$  by (3.16). Hence  $q > |T_1V : T_1| = |V : V \cap T_1|$  and so  $|V \cap T_1| > q^3$ . Therefore, for each  $B_1 \in B^G$  such that  $B_1 \leq V$ ,  $q > |B_1(V \cap T_1) : V \cap T_1| = |B_1 : B_1 \cap T_1| = |B_1 : B_1 \cap T|$ . Hence  $|B_1 \cap T| > q$ . Let  $\gamma \in F(B_1 \cap T)$  and set  $B_2=U_\gamma$ . Then  $\langle B_1 \cap T, B_2 \rangle \leq N^\gamma \cap T$ . As  $|B_1 \cap T| > q$  by (2.5) (iii),  $B_1 \cap T \cap B_2 \neq 1$ . By (3.11),  $\langle B_1 \cap T, B_2 \rangle \leq G_{F(B_2)}$ . By (3.12) (i), we have  $B_1 \cap T \leq B_2$ , so that  $F(B_1) = F(B_1 \cap T) = F(B_2)$ . Again, by (3.12) (i),  $B_1 = B_2$  and so  $U, V \leq C_G(B_2)$ . Therefore  $U=V$  by (3.15), a contradiction.

(3.22) *Put  $W=U^G$ . Then  $|W|=q^2+q+1$  and  $G^W$  is doubly transitive.*

Proof. Set  $H=N_G(U)$ . By (3.10) (ii) and (3.16),  $|W|=|G : H|=q^2+q+1$ . Let  $V \in W - \{U\}$  and let  $Q$  be as defined in (3.20). By (3.15),  $Q \leq H$  and by (3.20),  $Q$  acts semi-regularly on  $W - \{U\}$ . Hence  $|V^H|$  is divisible by  $q+1$ . On the other hand, by (3.21),  $|V^H|$  is divisible by  $q$  and so we have  $|V^H|=q(q+1)$ . Thus (3.22) holds.

(3.23)  $G_W \cap U \neq 1$ .

Proof. Suppose  $G_W \cap U=1$ . Since  $G \triangleright G_W$  and  $H \triangleright U$ ,  $[G_W, U] \leq G_W \cap U = 1$ . Hence  $G_W \leq C_G(U)$ . By (3.15),  $U$  is a unique Sylow 2-subgroup of  $C_G(U)$  and so  $G_W \leq O(G)$ . On the other hand, as  $|\Omega|$  is even and  $G$  is doubly transitive on  $\Omega$ , we have  $O(G)=1$ . Therefore  $G_W=1$  and hence  $G$  acts faithfully on  $W$ . Since  $U$  is not semi-regular on  $W - \{U\}$ , by [4],  $PSL(n_1, q_1) \leq G \leq P\Gamma L(n_1, q_1)$  for some  $n_1 \geq 3$  and  $q_1$  with  $q_1$  even. As  $|W|=q^2+q+1=q_1^{n_1-1} + \dots + q_1 + 1$ ,  $q(q+1)=q_1(q_1^{n_1-2} + \dots + 1)$  and so  $q=q_1$  and  $n_1=3$ . Therefore  $PSL(3, q) \leq G \leq P\Gamma L(3, q)$ . But  $|P\Gamma L(3, q)|_2=q^3$  by (3.9) (ii) and Lemma 2.6. Hence  $q^3=q^6$  by (3.10) (ii). This is a contradiction. Thus  $G_W \cap U \neq 1$ .

(3.24)  $G^\Omega$  has a regular normal subgroup.

Proof. Since  $G_W \leq N_G(U)$ ,  $G_W \cap U$  is a normal subgroup of  $G_W$ . As  $G_W \cap$

$U \leq 0_2(G_w)$  and  $G \geq G_w$ ,  $0_2(G_w)$  is a normal subgroup of  $G$ . Let  $E$  be a minimal normal subgroup of  $G$  contained in  $0_2(G_w)$ . Then  $E$  is an elementary abelian 2-subgroup of  $G$  and acts regularly on  $\Omega$ .

(3.25) *If (ii) of (3.7) occurs, we have (i) of the theorem.*

Proof. By (3.9), (3.10) and (3.24),  $G$  has a regular normal subgroup  $E$  of order  $q^3$ , where  $q=2^n$  and  $n \equiv 1 \pmod{2}$  and  $N^\alpha$  is transitive on  $\Omega - \{\alpha\}$ . Moreover  $G = G_\alpha E$  and  $G_\alpha$  is isomorphic to a subgroup of  $GL(E)$ . As in the proof of Lemma 2.1, we may assume  $\Omega = E$ ,  $\alpha = 0 \in E$  and  $GL(E)E \leq \text{Sym}(\Omega)$ . There exists a subgroup  $H$  of  $GL(E)$  such that  $H \simeq \Gamma L(3, q)$  and  $HE \simeq A\Gamma L(3, q)$ . Let  $L$  be a normal subgroup of  $H$  isomorphic to  $SL(3, q)$ . Since  $q=2^n$  and  $n \equiv 1 \pmod{2}$ ,  $L$  is isomorphic to  $PSL(3, q)$ .

By (3.9) (i) and by the structure of  $A\Gamma L(3, q)$ , there exist an automorphism  $f$  from  $N^\alpha$  to  $L$  and  $g \in \text{Sym}(\Omega)$  such that  $\alpha^g = \alpha$  and  $(\beta^x)^g = (\beta^g)^{f(x)}$  for each  $\beta \in \Omega - \{\alpha\}$  and  $x \in N^\alpha$ . From this  $(\beta^g)^{g^{-1}xg} = (\beta^x) = (\beta^g)^{f(x)}$  for each  $\beta \in \Omega - \{\alpha\}$  and so  $g^{-1}xg = f(x)$ . Hence  $g^{-1}N^\alpha g = L$ .

Set  $X = N(L) \cap \text{Sym}(\Omega)$  and  $D = C_X(L)$ . Then  $D$  is semi-regular on  $\Omega - \{\alpha\}$  as  $L$  is transitive on  $\Omega - \{\alpha\}$ . Put  $T = f(A)$ . Then  $N_L(T)^{F(T)} \simeq Z_{q-1}$  and it is semi-regular on  $F(T) - \{\alpha\}$  by (3.5) (i) and (3.9) (i), (iii). It follows that  $D \leq Z_{q-1}$ . Since  $X/DL$  is isomorphic to a subgroup of the outer automorphism group of  $PSL(3, q)$  and  $f(A)$  and  $f(B)$  are not conjugate in  $\text{Sym}(\Omega)$  by the hypothesis (\*\*\*) and (3.9) (ii), it follows from Lemma 2.6 (i) that  $|X/DL| \leq n$ . Hence  $|X| \leq n(q-1)|L| = |\Gamma L(3, q)|$ . On the other hand  $\Gamma L(3, q) \simeq H < X$  and so  $X = H$ . Therefore  $g^{-1}G_\alpha g \geq g^{-1}N^\alpha g = L$  and  $g^{-1}G_\alpha \leq X = H$ . Thus we have (3.25).

The conclusion of the theorem now follows immediately from steps (3.2), (3.7), (3.8) and (3.25).

OSAKA KYOIKU UNIVERSITY

### References

- [1] D. Gorenstein: *Finite groups*, Harper and Row, New York, 1968.
- [2] Y. Hiramane: *On doubly transitive permutation groups*, Osaka J. Math. **15** (1978), 613-631.
- [3] Y. Hiramane: *On some doubly transitive permutation group in which socle( $G_\alpha$ ) is nonsolvable*, Osaka J. Math. **16** (1979), 797-816.
- [4] M. O'Nan: *A characterization of  $L_n(q)$  as a permutation group*, Math. Z. **127** (1972), 301-314.
- [5] M. O'Nan: *Normal structure of one-point stabilizer of a doubly transitive permutation group II*, Trans. Amer. Math. Soc. **214** (1975), 43-74.
- [6] R. Steinberg: *Automorphism of finite linear groups*, Canad. J. Math. **12** (1960),

606–615.

- [7] H. Zassenhaus: *Über transitive Erweiterungen gewisser Gruppen aus Automorphismen endlicher mehrdimensionaler Geometrien*, Math. Ann. **111** (1935), 748–759.