



Title	Clinical Implications of Serum Mac-2 Binding Protein in Patients After Living Donor Liver Transplantation for Biliary Atresia
Author(s)	Ueno, Takehisa; Takase, Koki; Deguchi, Koichi et al.
Citation	Transplantation Proceedings. 2024, 56(2), p. 343-347
Version Type	AM
URL	<a href="https://hdl.handle.net/11094/99839">https://hdl.handle.net/11094/99839</a>
rights	© 2024. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Table 1 Patient demographic

Variables	n=56
Age, years	4.3 (0.6-18.9)
Observation, years	8.6(1.2-20.6)
Gender, male/female	17/39
AST ,IU/L	28(13-56)
ALT ,IU/L	19(7-35)
Serum albumin, g/dL	3.8(2.5-4.5)
Total bilirubin, mg/dL	0.6(0.3-1.6)
Prothrombin time, INR	1.18(0.98-1.69)
Platelets, 104/mm3	199(43-455)
M2BPGi, COI	0.8(0.2-1.8)
Type IV Collagen 7s, ng/ml (n=52)	5.6(3.1-9.9)
Hyaluronic acid, ng/ml (n=33)	25(9-153)
APRI	0.4(0.1-1.9)
FIB-4	0.4(0.1-1.5)
Histological findings	
F0	7(12%)
F1	36(64%)
F2	13(23%)
Portal vein	
normal	51(91%)
stenosis	3(5%)
thrombosis	2(4%)

**Table 1. Patient demographics:** Data were expressed as median with ranges. Histological findings were based on METAVIR score. Type IV collagen 7s acid in 4 patients and hyaluronic acid in 23 patients were not measured. Therefore, analysis of these data was performed in 52 and 33 patients respectively. APRI: the AST to Platelet Ratio Index; AST: aspartate aminotransferase; ALT: alanine transaminase; COI: Cut Off Index; FIB-4: Fibrosis-4. M2BPGi: Mac-2 binding protein glycosylation-modified isomer