



Title	Clinical Implications of Serum Mac-2 Binding Protein in Patients After Living Donor Liver Transplantation for Biliary Atresia
Author(s)	Ueno, Takehisa; Takase, Koki; Deguchi, Koichi et al.
Citation	Transplantation Proceedings. 2024, 56(2), p. 343-347
Version Type	AM
URL	https://hdl.handle.net/11094/99839
rights	© 2024. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Table 2 Receiver operating characteristic of fibrosis markers

	AUC	Value	Sensitivity, %	Specificity, %
M2BPGi (COI)	0.78	0.94	58	90
AST to ALT ratio	0.63	1.37	67	67
Platelet count($10^4/\text{mm}^3$)	0.54	206	67	51
FIB-4 index	0.6	0.53	42	82
APRI	0.55	0.51	33	82
Type IV Collagen 7s(mg/L)	0.58	5.3	70	49
Hyaluronic acid(ng/ml)	0.49	27	100	21

Table 2. Receiver operating characteristic of fibrosis markers. Summary of AUC, optimal cutoff point, sensitivity, and specificity for each fibrotic marker: APRI: the AST to Platelet Ratio Index; AST: aspartate aminotransferase; ALT: alanine aminotransferase; COI: Cut Off Index, FIB-4: Fibrosis-4 index